RSC Advances



PAPER

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2018, 8, 11813

Identification of metabolites of liquiritin in rats by UHPLC-Q-TOF-MS/MS: metabolic profiling and pathway comparison in vitro and in vivo†

Xia Zhang, Caijuan Liang, Jintuo Yin, Yupeng Sun and Lantong Zhang **D**

Liquiritin (LQ), the main bioactive constituent of licorice, is a common flavoring and sweetening agent in food products and has a wide range of pharmacological properties, including antidepressant-like, neuroprotective, anti-cancer and anti-inflammatory properties. This study investigated the metabolic pathways of LQ *in vitro* (rat liver microsomes) and *in vivo* (rat model) using ultra high-performance liquid chromatography coupled with hybrid triple quadrupole time-of-flight mass spectrometry (UHPLC-Q-TOF-MS/MS). Moreover, supplementary tools such as key product ions (KPIs) were employed to search for and identify compounds. As a result, 56 *in vivo* metabolites and 15 *in vitro* metabolites were structurally characterized. Oxidation, reduction, hydrolysis, methylation, acetylation, and sulfate and glucuronide conjugation were determined to be the major metabolic pathways of LQ, and there were differences in LQ metabolism *in vitro* and *in vivo*. In addition, the *in vitro* and *in vivo* metabolic pathways were compared in this study.

Received 30th December 2017 Accepted 3rd March 2018

DOI: 10.1039/c7ra13760e

rsc.li/rsc-advances

1. Introduction

Licorice root or *Radix glycyrrhizae*, a Chinese materia medica (CMM), is widely used to invigorate the spleen, replenish the Qi, dispel heat and remove toxic substances.¹ In modern medical terms, the biologically active compounds in licorice are triterpene, saponins, flavonoids, polysaccharides and phenolic compounds,² making licorice root a source of medicine and food. These compounds exhibit several important pharmacological activities, including anti-viral,³ anti-oxidant,⁴ anti-bacterial,⁵ anti-inflammatory⁶ and anti-HIV² activities. Moreover, licorice and its extract are widely used in health foods because of the physiological activities of these substances.⁶ Currently, the use of licorice and its extract in grain products, oil products, meat products, beverages, candy, jelly, dried fruit,




Fig. 1 Chemical structure of LQ

Department of Pharmaceutical Analysis, School of Pharmacy, Hebei Medical University, Shijiazhuang 050017, P. R. China. E-mail: zhanglantong@263.net; Fax: +86-311-86266419; Tel: +86-311-86266419

† Electronic supplementary information (ESI) available: Detailed information regarding other metabolites found *in vitro* and *in vivo*. Fig. S1 MS/MS spectra of all metabolites of LQ detected *in vitro* and *in vivo*. See DOI: 10.1039/c7ra13760e

seeds, soy sauce, *etc.*, are being extensively researched.^{9,10} Among the biologically active compounds in licorice, liquiritin (LQ, Fig. 1) is the dominant component and is considered to be the major active ingredient.¹¹

Drug metabolism may lead to detoxification and/or activation reactions, and studies of drug metabolism can aid the identification of active compounds and explain the mechanisms of action of these compounds. It is well known that the liver plays a key role in the metabolism of orally administered drugs. The rat liver microsome system is often considered as a reasonable model in which to study drug metabolism. On the other hand, *in vivo* metabolic studies could comprehensively reveal the metabolic pathways of drugs.

In recent years, the use of liquid chromatography coupled with tandem mass spectrometry (LC-MS/MS) has been routinely used to detect and identify metabolites^{13,14} and has been used to study drug metabolism (flavones and flavonoids,¹⁵ phenylpropanoids,^{16–18} terpenes,¹⁹ alkaloids,²⁰ saponins,²¹ stilbenes²² and traditional Chinese medicinal extracts²³), pharmacokinetics and toxicokinetics of metabolites,^{24,25} and tissue distribution and excretion of metabolites.²⁶ In addition, this method has applications in lipidomics,²⁷ proteomics²⁸ and metabolomics.^{29,30} A primary advantage of tandem mass spectrometry (MS/MS) is the ability of this method to detect a broad range of drugs with high sensitivity and specificity in a single analytical run.^{31–34} In addition, high-resolution mass spectrometry confirms structures by comparing the exact measured mass of a compound with the exact theoretical mass.^{35–37}

To our knowledge, there has been one report of the metabolic profile of LQ China;³⁸ however, this report was incomplete Open Access Article. Published on 27 March 2018. Downloaded on 11/5/2025 10:00:54 PM

RSC Advances Paper

and only identified 7 metabolites. In this research contribution, a simple and rapid UHPLC-Q-TOF-MS/MS approach combined with pattern recognition analysis was first employed to rapidly screen and characterize metabolites of LQ *in vitro* and *in vivo*, which was the first systematic study of the metabolism of LQ *in vitro* and *in vivo*. The characterization of 56 *in vivo* metabolites and 15 *in vitro* metabolites was achieved by UHPLC-Q-TOF-MS/MS analysis based on the MS/MS spectra and clog *P* values. In addition, the metabolic pathways of LQ were summarized. These results provide insight into the metabolic mechanism of LQ and lay the foundation for novel drug design.

Materials and methods

2.1. Chemicals and materials

LQ (CAS No: 551-15-5) was purchased from Chengdu Desert Biotechnology Co., Ltd. (Chengdu, China). β-NADP (β-nicotin-amine adenine dinucleotide phosphate), glucose-6-phosphate (G-6-P) and glucose-6-phosphatedehydrogenase (G-6-PD) were purchased from Sigma Chemical Co. (St. Louis, MO, USA). MgCl₂, UDPGA (uridine-5′-diphosphoglucuronic acid trisodium salt), Tris-HCl and alamethicin were purchased from BD Biosciences (Woburn, MA, USA). Phosphate-buffered saline (PBS) was purchased from Sangon Biotech Co., Ltd. (Shanghai, China). HPLC-grade methanol was purchased from J.T. Baker Chemical Company (Phillipsburg, NJ, USA). Formic acid (HPLC grade) was provided by Diamond Technology (Dikma Technologies Inc., Lake Forest, CA, USA). Purified water was obtained from Hangzhou Wahaha Group Co., Ltd. (Hangzhou, China).

2.2. Instrumentation and conditions

UHPLC-Q-TOF-MS/MS analysis was performed on a Shimadzu UHPLC system (Shimadzu Corp., Kyoto, Japan) coupled with a triple TOFTM 5600⁺ MS/MS system (AB Sciex, CA, USA). Information-dependent acquisition (IDA) was carried out. Chromatographic separation was conducted on a Poroshell 120 EC-C₁₈ (2.1 \times 100 mm, 2.7 μ m) column with a SecurityGuard® UHPLC C₁₈ pre-column (Agilent Corp, Santa Clara, CA, USA). The column temperature was maintained at 25 °C. The mobile phase was consisted of water containing 0.1% formic acid (A) and methanol (B). The gradient elution program was optimized for the separation, and the program was as follows: 0-2 min, 10-15% B; 2-15 min, 15-45% B; 15-20 min, 45-95% B; 20-25 min, 95-95% B. After maintaining the column at 95% solvent B for 5 min, the column was returned to its starting conditions over 1 min and was equilibrated in 10% solvent B for 5 min. The flow rate of the mobile phase was set to 0.3 mL min⁻¹, and the injection volume was 3 μL.

A Triple TOFTM 5600 system with DuoSprayTM ion sources (AB Sciex Triple TOFTM 5600+, Concord, Ontario, Canada) operating in the negative electrospray ionization mode was used for detection. The following MS/MS conditions were used: ion spray voltage, -4.5 kV; the turbo spray temperature, 550 °C; and declustering potential (DP), -60 V. Nitrogen was used as the nebulizer and auxiliary gas. Furthermore, the flows of the nebulizer gas (gas 1), heater gas (gas 2) and curtain gas were set

to 55, 55 and 35 L min⁻¹, respectively. The collision energy (CE) was set to -35 eV, and the collision energy spread (CES) was 15 eV.

Metabolite identification was performed with MetabolitePilot 1.5 (AB Sciex, CA, USA) based on accurate measurements of *m/z* values and on the processing of the data obtained from the XIC (extracted ion chromatography), MDF (mass defect filtering), PIF (product ion filtering) and NLF (neutral loss filtering) screening of putative metabolites. In addition, elemental compositions and chemical formulas were calculated.

2.3. Animals and drug administration

Thirty male Sprague-Dawley (SD) rats, 10-12 weeks in age and weighing 200-230 g, were obtained from the Experimental Animal Center of Hebei (Shijiazhuang, China). Rats were housed under standard temperature, humidity and light conditions. The animals were kept in an environmentally controlled breeding room for 7 days and fasted for 12 h before experiments. LQ was dissolved in a 0.5% carboxymethyl cellulose sodium (CMC-Na) aqueous solution. Thirty male SD rats were divided into six groups of five rats per group, which included experimental blood, urine and feces, and bile groups as well as blank blood, urine and feces, and bile groups. The prepared LQ suspension was orally administered to 15 rats from the experimental blood, urine and feces, and bile groups at a dose of 120 mg kg^{-1} , and 0.5% CMC-Na aqueous solution was orally administered to 15 rats from the blank blood, urine and feces, and bile groups. All experiments were conducted in accordance with the guides of Animal Care and Use Committee at Hebei Medical University. This study was also performed in strict accordance with the NIH guidelines for the care and use of laboratory animals (NIH Publication No. 85-23 Rev. 1985) and was approved by the Institutional Animal Care and Use Committee of National Tissue Engineering Center (Shanghai, China).

Plasma sample (five SD rats) collection was performed as follows: blood was taken from the canthi of the rats 0.17, 0.50, 0.75, 1, 2, 4, 6, 9, 12 and 24 h after administration. After centrifugation at $1400 \times g$ for 5 min (Hunan Xiangyi Laboratory Instrument Development Co. Ltd., Hunan, China), the supernatant was collected, and all plasma samples were combined. Blank plasma was collected in the same manner from rats (five) administered 0.5% CMC-Na aqueous solution.

Urine and feces (five SD rats) collection was performed as follows: urine and feces were collected during the 0-4 h, 4-8 h, 8-12 h, 12-24 h, 24-36 h, 36-48 h, 48-60 h and 60-72 h periods after administration, and all the urine and feces samples were combined. Rats (five) administered 0.5% CMC-Na aqueous solution were subjected to the same process to collect blank urine and feces samples.

Bile (five SD rats) collection was performed as follows: rats were administered urethane-containing physiological saline solution (1.5–2 g $\rm kg^{-1}$) after gavage, and then, bile duct cannulation. Then, bile samples were collected during 0–1 h, 1–3 h, 3–5 h, 5–8 h, 8–12 h, 12–20 h and 20–24 h periods after

Paper

administration. Finally, all bile samples were consolidated. Rats (five) administered 0.5% CMC-Na aqueous solution were subjected to the same process to collect a blank bile sample.

Three milliliters of blood, urine and bile samples were taken, and the protein in the samples was precipitated by methanol. Then, the supernatant was concentrated to dryness under reduced pressure at 25 °C using a Heidolph Laborota 4001 rotatory evaporator (Heidolph Instruments, GmbH & Co., Schwabach, Germany). The dried samples were dissolved in 300 μ L of methanol in an ultrasonic bath for 5 min, and then, the samples were centrifuged for 10 min at 10 000 \times g. Then, the supernatant was injected into the UHPLC-Q-TOF-MS/MS system for further analysis.

Methanol (20 mL) was added to the feces sample (2.0 g), and then, the sample was ultrasonicated for 45 min (Kun Shan Ultrasonic Instruments Co., Kunshan, China). After the mixture was centrifuged at 10 000 \times g for 10 min, the supernatant was collected and blow-dried in a nitrogen atmosphere. The residue was dissolved in 400 μ L of methanol and centrifuged at 10 000 \times g for 10 min. The supernatant (3 μ L) was injected into the chromatographic instrument.

All the bio-samples were placed in the $-80~^{\circ}\text{C}$ freezer for storage.

2.4. Microsomal incubation

- **2.4.1. Preparation of standard solutions.** The appropriate amount of standard LQ was accurately weighed and dissolved in 50% methanol to make the standard solution (2.15 mg mL $^{-1}$). The solution was stored at 4 °C in a refrigerator.
- 2.4.2. Preparation of liver microsomes. Liver microsomes were prepared by differential centrifugation.³⁹ All surgical instruments and experimental reagents were stored at 4 °C in refrigerator. Five male SD rats (220-250 g) were fasted for 24 h and decapitated. The liver was quickly retrieved, and the blood was blotted with filter paper. Then, the liver was weighed and was repeatedly washed with sugar solution till a khaki color. Then, the liver was added to an ice-cold sugar solution that was 4 times the weight of the liver and was then cut and homogenized. After centrifugation at 20 000 \times g for 20 min at 4 $^{\circ}$ C, the precipitate was discarded. After additional centrifugation at 100 000 \times g for 60 min at 4 °C, the supernatant was discarded. The precipitate was washed with 4 times as much cold Tris-HCl solution. After centrifugation at 100 000 \times g for 60 min at 4 $^{\circ}$ C, the precipitation obtained was resuspended using 4 times as much Tris-HCl solution to obtain liver microsomes. Finally, the liver microsomes were placed at -80 °C in the freezer for storage until further use. In addition, the protein concentration of the liver microsome suspension was determined by the Lowry method.40
- **2.4.3. Phase I metabolism.** The typical incubation mixture (200 μL final volume) consisted of a 0.1 mol L^{-1} K_2HPO_4 buffer containing rat microsomes (1.0 mg m L^{-1}), 3.3 mmol L^{-1} MgCl₂, 1.3 mmol L^{-1} β-NADPH, 3.3 mmol L^{-1} glucose-6-phosphate, 1.0 U m L^{-1} glucose-6-phosphate dehydrogenase and 100 μmol L^{-1} LQ methanol (final amount in the reaction medium: 1%). Preincubation was carried out at 37 °C for 5 min, and then,

NADPH was added to initiate the reaction. After incubation for 30 min at 37 $^{\circ}\text{C}$ in a metabolic shaker (1200 \times g, Hangzhou Miu Instruments Co., Ltd., Hangzhou, China), 1 mL of ice-cold ethyl acetate was added to stop the reaction, and the mixture was vortexed for 5 min. After centrifugation at 10 000 \times g for 10 min, the organic phase was collected and evaporated under nitrogen gas. Residues were redissolved in 100 μL of methanol, and an aliquot (3 $\mu\text{L})$ was injected into the chromatographic system for analysis. The blank sample was incubated without LQ, while the control sample was incubated without the NADPH-generating system by following the method described above.

2.4.4. Phase II metabolism. The reaction mixture (total volume 200 μ L) containing 1.0 mg mL⁻¹ rat liver microsomes, 2 mmol L⁻¹ UDPGA, 8 mmol L⁻¹ MgCl₂, 25 μ g mL⁻¹ alamethicin in PBS (pH 7.4) and 100 μ mol L⁻¹ LQ methanol (final amount in the reaction medium: 1%) solution was preincubated for 20 min at 37 °C, and then, UDPGA was added to start the reaction. After incubation for 60 min at 37 °C in a metabolic shaker (1200 \times *g*), 200 μ L of ice-cold methanol was added to stop the reaction. After vortexing for 5 min, the organic phase was collected and evaporated under nitrogen gas. Methanol (100 μ L) was added to the residues, and an aliquot (3 μ L) was injected into the chromatographic system for analysis. The blank sample was incubated without LQ, while the control sample was incubated without the UDPGA-generating system following the method described above.

3. Results and discussion

A total of 56 *in vivo* metabolites and 15 *in vitro* metabolites were detected in the experimental conditions used. The metabolites detected *in vivo* and *in vitro* are listed in Tables 1 and 2, respectively, and the structures of the metabolites are shown in Fig. 2. The XIC and MS/MS data of the metabolites detected *in vivo* and *in vitro* are presented in Fig. 3 and S1,† respectively.

3.1. Analytical strategy and metabolite analysis

In this study, a method of metabolite identification was developed that was based on a Triple TOFTM instrument with a multiple mass defect filter (MMDF) combined with dynamic background subtraction (DBS)-dependent on-line data acquisition and multiple post-acquisition data mining technologies. First, on-line data and accurate MS/MS data were acquired using a full-scan, unique and effective MMDF and a DBSdependent data acquisition method.41 Then, post-acquisition data mining was performed using various data-mining tools such as XIC, MDF, PIF and NLF.42 Furthermore, structures of the metabolites of LQ were elucidated based on accurate mass measurements, knowledge of relevant drug bio-transformation, previously investigated fragmentation patterns of LQ, and MS/ MS spectra of metabolites. Peakview 1.2 software was used to identify possible metabolites by comparing the extracted ion chromatograms and base peak chromatograms of the sample group with those of the blank group. 43 Finally, the useful clog P parameter, which was calculated using the ChemDraw Ultra 12.0 program, was introduced to distinguish between structural

Open Access Article. Published on 27 March 2018. Downloaded on 11/5/2025 10:00:54 PM.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

Table 1 Summary of phase I and phase II metabolites of LQ in rat blood, urine, bile and feces samples a

												I
Metabolites ID	Composition shift	Formula	z/m	Error (ppm)	$t_{ m R}$ (min)	Score (%)	MS/MS fragments	$\operatorname{clog} P$	Blood	Urine B	Bile Fe	Feces
M1	Hydrolysis + tera-oxidation	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_{8}$	319.0477	1.6	7.95	88.5	273.1616, 255.1246, 239.0917, 221.0782,	0.509837		+	+	
M2	Hydrolysis + tera-oxidation	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_{8}$	319.0477	2.6	9.54	88.5	273.1699, 255.1614, 239.0926, 221.0815,	0.569225		+	+	
M3	Hydrolysis + tera-oxidation	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_{8}$	319.0482	1.0	13.20	92.8	1//.0911, 151.04/6, 255.1595, 239.0923,	0.629837		+	+	
M4	Hydrolysis + tri-oxidation	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_7$	303.0533	1.4	86.8	84.6		1.16623	ı	+	+	
M5	Hydrolysis + tri-oxidation	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_7$	303.0536	4.6	11.42	83.4	153.0922, 137.0967 285.1340, 259.1548, 255.1233, 241.1436, 223.0967, 217.1073, 205.0858, 163.0755,	1.22684	'	+	+	
M6	Hydrolysis + Tri-oxidation	$C_{15}H_{12}O_7$	303.0532	3.2	11.94	90.1	153.0916, 137.0963 285.1342, 259.1540, 255.1299, 241.1430, 223.0975, 217.1075, 205.0865, 163.0750, 175.0020, 137.0070	1.28623		+	+	
M7	Oxidation	$C_{21}H_{22}O_{10}$	433.1115	6.0-	9.01	83.1	153.0920, 137.0972 271.0596, 243.0645, 227.0692, 164.0107, 136.0151 109.0290	0.300687	+	+	+	
M8	Oxidation	$C_{21}H_{22}O_{10}$	433.1111	-2.7	12.61	85.3	271.0593, 243.0646, 227.0695, 164.0103, 136.0152, 109.0289	0.500687	+	+	+	
М9	Oxidation	$C_{21}H_{22}O_{10}$	433.1115	-1.9	15.65	80.4	271.0598, 243.0650, 227.0705, 164.0098, 136.0143, 100.0304	0.558399	+	+	+	
M10	Oxidation	$C_{21}H_{22}O_{10}$	433.1114	-2.1	16.45	83.4	130.0142, $109.0294271.0594$, 243.0652 , 227.0689 , 164.0099 , 136.0154, 109.0285	0.948399	+	+	+	
M11	Oxidation + ketone formation	$\mathrm{C}_{21}\mathrm{H}_{20}\mathrm{O}_{11}$	447.0909	-2.4	9.11	75.0	271.0600, 175.0238, 135.0443, 113.0242	-0.178159		+	ı	
M12	Oxidation + ketone formation	$C_{21}H_{20}O_{11}$	447.0891	-4.4	10.84	78.4	271.0607, 175.0235, 135.0445, 113.0248	0.0218411		+	I	
M13	Oxidation + ketone formation	$C_{21}H_{20}O_{11}$	447.0903	-4.8	11.67	74.4	271.0603, 175.0240, 135.0444, 113.0250	0.0795532	·	+	ı	
M14	Oxidation + ketone formation	$C_{21}H_{20}O_{11}$	447.0909	-4.3	13.12	76.7	271.0601, 175.0232, 135.0446, 113.0255	0.469553		+ -	-	
МТЭ	Hydrolysis + oxidadon	$C_{15}H_{12}O_{5}$	2/1.0635	-0.0	11.52	84.1	253.1385, 235.1150, 227.0705, 209.1546, 191.1430, 183.0113, 151.1109, 145.0525,	1.93694		+	+	
							119.0500					
M16	Hydrolysis + oxidation	$\mathrm{C_{15}H_{12}O_{5}}$	271.0610	-2.5	13.66	90.3	253.1437, 235.1322, 227.1286, 209.1168, 191.1421, 183.1393, 151.0022, 145.0499, 119.0497	2.05485	ı	+	+	
M17	Hydrolysis + oxidation	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_5$	271.0605	-1.4	13.88	74.8	253.1437, 235.1332, 227.1643, 209.1540, 191.1436, 183.1380, 151.0033, 145.0292, 119.0505	2.44485	ı	+	+	
M18	Hydrolysis + tri-oxidation + desaturation	$C_{15}H_{10}O_{7}$	301.1284	4.0	12.89	75.1	283.1184, 257.1402, 255.0857, 241.1068, 239.1275, 221.1159, 211.0960, 169.0858, 118.9799	1.67912	+	+		
M19	Hydrolysis + oxidation + desaturation	$\mathrm{C}_{15}\mathrm{H}_{10}\mathrm{O}_{5}$	269.0464	0.1	14.64	75.1	241.0512, 225.0547, 135.0085, 133.0286	2.90529				
M20a M20b M20c	Hydrolysis + oxidation + methylation	$\mathrm{C_{16}H_{14}O_{5}}$	285.0791	-2.8	16.21	82.8	262.0798, 255.0729, 254.0721, 183.0107, 119.0495, 96.9603, 79.9575	2.50762 2.96762 3.03085	+	+	+	
M21	Oxidation + methylation	$C_{22}H_{24}O_{10}$	447.1278	-4.2	17.00	83.9	385.1283, 285.1180, 271.0971, 255.0731, 175.0243, 165.0551, 113.0248	1.01117	+	+	+	

Open Access Article. Published on 27 March 2018. Downloaded on 11/5/2025 10:00:54 PM.

Pr-No This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

_		Ī
7	7	٠
-	1	
-	0220	
-	>	:
1	Ĺ	3
(١
2	-	
_	_	
	a	1
_	-	
4	c	1
f	0	
L	٠,	

Metabolites												
ID	Composition shift	Formula	m/z	Error (ppm)	$t_{\rm R} $	Score (%)	MS/MS fragments	$\operatorname{clog} P$	Blood	Urine	Bile	Feces
M22	Oxidation + methylation	$\mathrm{C}_{22}\mathrm{H}_{24}\mathrm{O}_{10}$	447.1273	-3.2	17.65	84.7	385.1281, 285.1150, 271.0967, 255.0739,	1.47117	+	+	+	+
M23	Desaturation	$C_{21}H_{20}O_9$	415.1780	-4.1	17.03	82.6	253.0132, 252.0214, 223.0244, 142.9952, 112.9901	1.06205	ı	+	+	+
M24	Hydrolysis + desaturation	$\mathrm{C}_{15}\mathrm{H}_{10}\mathrm{O}_4$	253.0509	0.1	17.58	91.4	135.0082, 133.0291, 117.0346	2.5753	ı	+	+	+
M25	Hydrolysis + methylation	$\mathrm{C_{16}H_{14}O_4}$	269.0780	-4.4	12.40	82.6	253.0637, 191.0387, 96.9607, 79.9585	2.83559	ı	+ -	+ -	+ -
M26 M27	Hydrolysis + metnylation Hydrolysis	$C_{16}H_{14}O_4$ $C_{47}H_{45}O_4$	255.0661	-3.9 -1.2	13.8/	92.3	253.0639, 191.0392, 96.9600, 79.9579 135.0090, 119.0510, 91.0204	3.11994 2.53394	l +	+ +	+ +	+ +
M28	Hydrolysis + bis-glucuronide	$C_{27}H_{28}O_{16}$	607.1310	-4.5	7.33	80.8	431.0975, 255.0655, 175.0240, 113.0248	-1.70101	.	+	+	+
M29	conjugation Hydrolysis + oxidation + sulfate +	$C_{21}H_{20}O_{14}S$	527.0471	-4.8	7.55	81.1	447.0917, 351.0165, 271.0599, 193.0343,	-1.91668	I	+	+	ı
	glucuronide conjugation	 					175.0237, 135.0081, 113.0242					
M30	Hydrolysis + oxidation + sulfate + olucuronide conjugation	$C_{21}H_{20}O_{14}S$	527.0471	-3.6	8.32	75.2	447.2034, 351.0510, 271.0601, 193.0340, 175.0233, 135.0068, 113.0240	-1.61368	ı	+	+	I
M31	Hydrolysis + oxidation + sulfate +	$C_{21}H_{20}O_{14}S$	527.0480	-4.1	10.20	78.1	447.0939, 351.0181, 271.0610, 193.0339,	-1.15368	ı	+	+	ı
	glucuronide conjugation	:	4	4	1	1	175.0239, 135.0084, 113.0245		-	-	_	
M32	Giucuronide conjugation	$C_{27}H_{30}O_{15}$	593.1479	-1.6	7.63	83./	41/.11/6, 255.064/, 1/5.0238, 135.0082, 119.0503, 113.0242	-1.22216	+	+	+	I
M33	Glucuronide conjugation	$C_{27}H_{30}O_{15}$	593.1476	-2.1	12.01	83.3	417.1143, 255.0658, 175.0254, 135.0101,	-1.01791	+	+	+	ı
M3/1	Hydrolysis + his-sulfate conjugation	S. O. H.	715 1517	α ς	20	8 08	335 0230 255 0658 135 0080 119 0494	_0 870/11	ı	+	+	
M35	Sulfate conjugation	C ₁₅ H ₁₂ O ₁₀ S	497 0759	13.0	8 70	86.6	417 1185 283 0806 255 0655 206 9858	-0.806862	ı	- +	- +	+
		021-22-120		2			167.0074, 135.0074, 113.0241					
M36	Hydrolysis + sulfate + glucuronide	$C_{21}H_{20}O_{13}S$	511.0541	-1.3	8.92	74.2	431.0970, 335.0220, 255.0654, 238.9310,	-1.28571	+	+	+	+
	conjugation						175.0244, 135.0084, 119.0506, 113.0242					
M37	Tri-oxidation + glucuronide conjugation	$\mathrm{C}_{27}\mathrm{H}_{30}\mathrm{O}_{18}$	641.1338	-0.7	9.25	79.1	337.0366, 303.0870, 281.1053, 255.0655, 135.0080, 119.0504	-2.78222	ı	I	ı	+
M38	Tri-oxidation + glucuronide conjugation	$\mathrm{C}_{27}\mathrm{H}_{30}\mathrm{O}_{18}$	641.1397	-0.4	12.14	0.97	337.0390, 303.0881, 281.1069, 255.0665,	-2.64222	ı	I	ı	+
M39	Oxidation + glucuronide conjugation	$C_{27}H_{30}O_{16}$	609.1423	-1.2	9.67	82.6	135.0081, 119.0498 433.1126, 255.0806, 175.0239, 151.0395,	-1.55013	1	+	1	ı
M40	Oxidation + glucuronide conjugation	$C_{27}H_{30}O_{16}$	609.1430	-3.2	12.55	81.2	113.0243 433.1146, 255.0812, 175.0237, 151.0395, 113.0240	-1.09013	I	+	I	I
M41	Hydrolysis + oxidation + sulfate	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_{8}\mathrm{S}$	351.0164	-4.5	10.67	80.8	271.0614, 151.0039, 119.0507	0.361621	I	+	+	+
7473	conjugation Undrolucie + oxidation + culfata		251 0162	0	<u> </u>	100	271 0610 161 0040 110 0600	0 031631		4	4	-1
74.74 TM	conjugation	C151112O83	6010.166	4.0	C#-11	0.67	Z/1:0010, 151:0040, 119:0509	0.021021	I	+	+	+
M43	Hydrolysis + oxidation + sulfate conjugation	$\mathrm{C_{15}H_{12}O_{8}S}$	351.0164	-4.7	12.00	93.3	271.0609, 151.0040, 119.0507	0.884852	I	+	+	+
M44	Hydrolysis + glucuronide conjugation	$C_{21}H_{20}O_{10}$	431.0976	-2.6	11.17	85.6	255.0658, 175.0245, 135.0086, 119.0503, 113.0247	0.27429	+	+	+	+
M45	Hydrolysis + glucuronide conjugation	$C_{21}H_{20}O_{10}$	431.0976	-1.5	11.97	94.0	255.0662, 175.0253, 135.0089, 119.0502, 113.0251	0.55864	+	+	+	+
M46	Hydrolysis + desaturation + glucuronide conjugation	$\mathrm{C}_{21}\mathrm{H}_{18}\mathrm{O}_{10}$	429.0828	0.2	12.13	82.6	253.0505, 113.0235	0.390001	I	+	+	I

Table 1 (Contd.)

Metabolites ID	Composition shift	Formula	z/w	Error $t_{\rm R}$ (ppm) (min)	$t_{ m R}$ (min)	Score (%)	MS/MS fragments	$\operatorname{clog} P$	Blood	Blood Urine	Bile	Feces
M47	$\label{eq:conjugation} Hydrolysis + desaturation + glucuronide \qquad C_{21}H_{18}O_{10}$ conjugation	$C_{21}H_{18}O_{10}$	429.0824	-0.7	12.78	85.7	253.0495, 113.0230	0.583201	I	+	+	I
M48	Oxidation + sulfate conjugation	$C_{21}H_{22}O_{13}S$	513.0667	-2.1	12.54	93.4	433.0514, 431.0986, 337.0368, 255.0641, 175.0226, 151.0390, 113.0244	-1.13483	+	+	+	+
M49	Oxidation + sulfate conjugation	$C_{21}H_{22}O_{13}S$	513.0677	-1.1	13.24	91.6	433.0558, 431.0976, 337.0365, 255.0665, 175.0247, 151.0393, 113.0241	-0.674831	+	+	+	+
M50	Hydrolysis + sulfate conjugation	$\mathrm{C}_{15}\mathrm{H}_{12}\mathrm{O}_7\mathrm{S}$	335.0216	-4.4	12.49	84.3	255.0656, 135.0083, 119.0500	0.689589	+	+	+	+
M51	Hydrolysis + sulfate conjugation	$\mathrm{C_{15}H_{12}O_7S}$	335.0217	-4.1	17.30	92.0	255.0660, 135.0088, 119.0502	0.97394	+	+	+	+
M52	Ketone formation	$\mathrm{C}_{21}\mathrm{H}_{18}\mathrm{O}_{11}$	445.1883	-2.1	12.53	89.4	269.0443, 113.0239, 104.9542	0.924951	+	+	+	1
M53	Loss of O + bis-ketone formation	$\mathrm{C}_{21}\mathrm{H}_{18}\mathrm{O}_{9}$	413.0854	-1.8	13.24	83.4	369.0968, 255.0660, 135.0088, 119.0505	0.29614	ı	+	1	ı
M54	Loss of O + bis-ketone formation	$\mathrm{C}_{21}\mathrm{H}_{18}\mathrm{O}_{9}$	413.0853	-2.1	14.22	74.4	369.0983, 255.0666, 135.0087, 119.0501	0.34904	ı	+	ı	ı
M55	Acetylation	$C_{23}H_{24}O_{10}$	459.1297	0.0	16.56	75.3	417.1198, 399.1064, 255.0645, 186.9290, 135.0070, 119.0489	0.769139	+	+	I	1
M56	Methylation	$\mathrm{C}_{22}\mathrm{H}_{24}\mathrm{O}_{9}$	431.0992	2.7	18.33	77.4	369.0977, 255.0666, 175.0244, 135.0086, 113.0248	1.33914	+	+	+	+

 a + Detected, - undetected, a, b and c - possible metabolites

isomers. Generally, compounds with the larger values have longer retention times in reversed-phase liquid chromatography systems. 44,45

All chemical constituents in TMM can be categorized into different families based on structural types. Thus, groups of compounds with identical carbon skeletons may yield similar fragmentation patterns and then generate the same characteristic fragment ions when subjected to collision-induced dissociation (CID) for mass spectrometry. Accordingly, a core supplementary tool in this approach is to use key product ions (KPIs) as markers for compounds detection and identification. $^{46-48}$ In this study, a KPI at m/z 255.0651 could be generated from common substructures as standards and was selected as a diagnostic ion for detecting relevant analogues of these substructures (shown in Fig. 3).

3.2. Mass fragmentation behavior of LQ

The retention time of LQ was 12.01 min, and LQ produced a molecular ion $[M-H]^-$ at m/z 417.1176 under the experimental conditions. Moreover, major secondary fragment ions at m/z 255.0651 $[M-H-C_6H_{10}O_5]^-$, 135.0079 (RDA reaction (retro Diels—Alder reaction)), 119.0496 (RDA reaction) and 91.0190 (RDA reaction) were detected. The proposed MS/MS fragmentation behavior and fragmentation pathways of LQ are shown in Fig. 4. The molecular weight and elemental composition of LQ were used as a baseline for comparison with some metabolites.

3.3. Identification of in vivo phase I metabolites

3.3.1. Oxidation reaction

Metabolites M1, M2 and M3. M1–M3 were eluted at 7.95, 9.54 and 13.20 min, respectively, with deprotonated molecular ions $[M-H]^-$ at m/z 319.0477, 319.0477 and 319.0482, respectively, which were 98 Da less than the value obtained for LQ and corresponded to $C_{15}H_{12}O_8$. The fragment ions at m/z 273.1696, 255.1595, 239.0923, 221.0814 and 151.0475 were produced from M1, M2 and M3 by the loss of CO, H_2O , O and by RDA reaction. Among these fragments, the fragment ion at m/z 255.1595 was hydrolyzed LQ. M1, M2 and M3 were oxidation metabolites of LQ and were assigned based on clog P values of 0.509837, 0.569225 and 0.629837 by ChemDraw 12.0 software.

Metabolites M7, M8, M9 and M10. M7–M10, eluted at 9.01, 12.61, 15.65 and 16.45 min, respectively, were characterized as deprotonated molecular ions $[M-H]^-$ at m/z 433.1115, 433.1111, 433.1115 and 433.1114, respectively, which were 16 Da more than the value obtained for the parent drug LQ. This finding suggested that M7–M10 were oxidation metabolites of LQ and that the chemical formula of M7–M10 was $C_{21}H_{22}O_{10}$. The observed diagnostic fragment ions at m/z 271.0596, 243.0645, 227.0692, 136.0151 and 109.0290 were generated by the loss of $C_6H_{10}O_5$, CO and O and by RDA reaction. The clog *P* values of M7–M10 were 0.300687, 0.550399 and 0.948399, respectively. Therefore, the four compounds were immediately identified on the basis of the retention times and clog *P* values.

Metabolites M11, M12, M13 and M14. M11–M14 were eluted at 9.11, 10.84, 11.67 and 13.12 min, respectively, with

Table 2 Summary of phase I and phase II metabolites of LQ in vitro^a

Metabolites ID	Composition shift	Formula	m/z	Error (ppm)	t _R (min)	Score (%)	MS/MS fragments	clog P	Blanks	Controls	Samples
N1	Oxidation	$C_{21}H_{22}O_{10}$	433.1110	-2.0	8.62	79.1	271.0592, 243.0645, 109.0289	0.300687	_	_	+
N2	Oxidation	$C_{21}H_{22}O_{10}$	433.1111	-2.8	9.01	79.0	271.0595, 243.0647, 109.0282	0.500687	_	_	+
N3	Oxidation	$C_{21}H_{22}O_{10}$	433.1112	-2.6	9.51	82.5	271.0596, 243.0645, 109.0292	0.558399	_	_	+
N4	Oxidation	$C_{21}H_{22}O_{10}$	433.1109	-2.2	14.03	84.4	271.0589, 243.0676, 109.0290	0.948399	_	_	+
N5	Hydrolysis +	$C_{15}H_{12}O_5$	271.0596	-1.1	11.12	89.7	253.0474, 151.0016, 135.0439,	1.93694	_	_	+
	oxidation						119.0487, 91.0191				
N6	Hydrolysis +	$C_{15}H_{12}O_5$	271.0595	-1.3	13.60	81.1	253.0486, 151.0024, 135.0444,	2.05485	_	_	+
	oxidation						119.0495, 91.0136				
N7	Hydrolysis +	$C_{15}H_{12}O_5$	271.0592	-1.5	13.83	73.4	253.0480, 151.0021, 135.0435,	2.44485	_	_	+
	oxidation						119.0480, 91.0131				
N8	Desaturation	$C_{21}H_{20}O_9$	415.1011	-1.7	12.63	76.2	253.0518, 252.0421, 223.0455,	1.06205	_	_	+
							142.99500, 112.9840				
N9	Hydrolysis	$C_{15}H_{12}O_4$	255.0650	-3.4	16.03	96.8	135.0083, 119.0500, 91.0193	2.53394	_	_	+
N10	Hydrolysis +	$C_{15}H_{10}O_5$	269.0437	-3.9	16.38	79.2	135.0090, 133.0287	2.90529	_	_	+
	desaturation + oxidation										
N11	Hydrolysis +	$C_{15}H_{10}O_4$	253.0496	-2.4	17.58	85.6	135.0077, 133.0284,	2.5753	_	_	+
	desaturation						117.0341, 91.0187				
N12a	Glucuronide	$C_{27}H_{30}O_{15}$	593.1565	3.7	7.62	74.4	417.1208, 255.0659, 175.0241,	-1.22216	_	_	+
N12b	conjugation						135.0078, 117.0187, 113.0241	-1.01791			
N13	Hydrolysis +	$C_{21}H_{20}O_{10}$	431.0989	1.3	11.19	86.4	255.0662, 226.9648, 175.0239,	0.27429	_	_	+
	glucuronide						135.0080, 119.0495, 113.0242				
	conjugation										
N14	Hydrolysis +	$C_{21}H_{20}O_{10}$	431.0985	0.2	11.93	82.8	255.0666, 226.9703, 175.0233,	0.55864	_	_	+
	glucuronide						135.0079, 119.0496, 113.0229				
	conjugation										
N15	Glucose	$C_{27}H_{32}O_{14}$	579.1752	3.9	12.45	89.3	417.1273, 402.9939, 255.0669,	-0.743309	_	_	+
	conjuagtion						238.9306, 135.0085, 119.0492				

^a + Detected, – undetected, a and b – possible metabolites.

deprotonated molecular ions $[M - H]^-$ at m/z 447.0909, 447.0891, 447.0903 and 447.0909, respectively, which were 30 Da more than the value obtained for LQ. The major fragment ions detected at m/z 271.0600, 135.0443 and 113.0242 were generated by the loss of $C_6H_8O_6$ and by RDA reaction, which implied that M11–M14 were oxidation products of LQ and that the chemical formula was $C_{21}H_{20}O_{11}$. In addition, M11, M12, M13 and M14 were assigned based on the clog P values of M11, M12, M13 and M14, which were -0.178159, 0.0218411, 0.0795532 and 0.469553, respectively.

Metabolite M18. M18 was detected at a retention time of 12.89 min, with a deprotonated molecular ion $[M-H]^-$ at m/z 301.1284, 46 Da higher than that of M0 losing $C_6H_{10}O_5$, which suggested that M18 was oxidation metabolite. The fragment ions at m/z 283.1184, 255.0857, 239.1275, 169.0858 and 118.9799 were gained by loss of H_2O , CO, O and by RDA reaction. According to the fragmentation, the chemical formula of M18 was $C_{15}H_{10}O_7$.

Metabolite M19. M19 was detected at a retention time of 14.64 min with a deprotonated molecular ion $[M-H]^-$ at m/z 269.0464, which was generated by the loss of two oxygens from M18. Typical fragment ions at m/z 241.0512, 225.0547, 135.0085 and 133.0286 were detected due to successive loss of CO and O and due to RDA reaction. Thus, M19 was deduced to be $C_{15}H_{10}O_5$.

Metabolite M20. M20 had a retention time of 16.21 min and was detected at m/z 285.0791 ([M – H]⁻), which was 30 Da more than the value obtained for hydrolyzed LQ, which implied that oxidation and methylation reactions had occurred. Representative fragment ions were observed at m/z 255.0729 and 119.0495 due to loss of CH₂O and due to RDA reaction, which suggested that the formula of M20 was $C_{16}H_{14}O_5$.

Metabolite M23. A peak was eluted at a retention time of 17.03 min. The MS/MS spectrum of M23 showed a deprotonated molecular ion $[M - H]^-$ at m/z 415.1780, which was 2 Da less than the value obtained for M0, which confirmed the molecular formula to be $C_{21}H_{20}O_9$. In addition, the MS/MS spectrum of M23 showed a number of characteristic fragment ions at m/z 253.0132 $[M-C_6H_{10}O_5-H]^-$, 142.9952 (RDA reaction) and 112.9901 (RDA reaction).

3.3.2. Reduction reaction

Metabolite M24. M24 was detected at 17.58 min and showed a deprotonated molecular ion $[M-H]^-$ at m/z 253.0509, which was 164 Da lower than the value obtained for M0, which suggested that the loss of two hydrogens from hydrolyzed M0. Meanwhile, representative fragment ions at m/z 135.0082, 133.0291 and 117.0346 were obtained due to RDA reaction. The formula was identified as $C_{15}H_{10}O_4$.³⁸

3.3.3. Hydrolysis reaction

Metabolites M25, M26. M25 and M26 displayed molecular ions $[M - H]^-$ at m/z 269.0780 and 269.0781, respectively, and

RSC Advances




Fig. 2 Chemical structures of all metabolites of LQ detected in vitro and in vivo (a, b and c-possible chemical structure).

had retention times of 12.40 and 13.87 min, respectively. The masses of the deprotonated M25 and M26 were 14 Da higher than the mass of hydrolyzed LQ, which suggested that a methylation reaction occurred on LQ after the loss of C₆H₁₀O₅. The MS/MS spectra showed a characteristic fragment ion at m/z253.0637 [M-O-H]⁻, which implied that the formula of M25 and M26 was C₁₆H₁₄O₄. Then, M25 and M26 were identified based on their clog P values of 2.83559 and 3.11994, respectively.

Metabolite M27. The metabolite M27 exhibited a deprotonated ion $[M - H]^-$ at m/z 255.0661, which was 162 Da lower than the value obtained for M0, confirming the occurrence of hydrolysis. In addition, typical fragment ions at m/z 135.0090, 119.0510 and 91.0204 were produced by RDA reaction. The formula of M27 was deduced to be C₁₅H₁₂O₄.38

Paper




Fig. 3 Extracted ion chromatograms of all metabolites of LQ detected *in vitro* and *in vivo*. (A1–2 in rat blood sample, B1–7 in rat urine sample, C in rat bile sample, D in rat feces sample, E1–4 in rat liver microsomes). Additional graph: metabolites detected by KPIs in rat blood, urine, bile and feces samples.

RSC Advances Paper




Fig. 3 (Contd.)


3.4. Identification of in vivo phase II metabolites

Metabolite M28. A peak was detected at 7.33 min and exhibited a deprotonated molecular ion $[M - H]^-$ at m/z607.1310, which was 190 Da higher than the value obtained for LQ. Secondary fragment ions were detected at m/z 431.0975 [M- $C_6H_8O_6-H$ and 255.0655 [M-2C₆H₈O₆-H], suggesting that hydrolyzed LQ had undergone bis-glucuronide conjugation. In addition, the chemical formula was deduced to be C₂₇H₂₈O₁₆.

3.4.2. Metabolites M29, M30, M31. M29-M31 were identified as sulfate and glucuronide conjugation metabolites. M29-M31 were eluted at 7.55, 8.32 and 10.20 min and had deprotonated molecular ions $[M - H]^-$ at m/z 527.0471, 527.0471 and 527.0480, which were 110 Da more than the value obtained for

M0. The product ions detected at m/z 447.0917, 271.0599, 175.0237, 135.0081 and 113.0242 were produced by the loss of SO₃ and C₆H₈O₆ and by RDA reaction, which suggested that the formula was C21H20O14S. In addition, the clog P values of M29, M30 and M31 were -1.91668, -1.61368 and -1.15368, respectively, and M29, M30 and M31 were identified based on this information.

3.4.3. Metabolites M32, M33. Two peaks were eluted at 7.63 and 12.01 min under the chromatography conditions used. M32 and M33 were isomers and had identical MS/MS spectra. M32 and M33 had deprotonated molecular ions $[M - H]^-$ at m/z593.1479 and 593.1476, respectively, which were 176 Da higher than the value obtained for M0, suggesting that M32 and M33 were glucuronide conjugation metabolites. Moreover, the MS/



MS/MS spectrum of LQ and its predominant fragmentation pathways.

MS spectra showed fragment ions at m/z 417.1143, 255.0658, 135.0101 and 119.0517 due to the loss of C₆H₈O₆ and C₆H₁₀O₅ and due to RDA reaction. In addition, the formula was deduced to be C₂₇H₃₀O₁₅. Then, M32 and M33 were identified based on their clog P values of -1.22216 and -1.01791, respectively.

3.4.4. Metabolite M34. M34 was eluted at 8.05 min with a deprotonated molecular ion $[M - H]^-$ at m/z 415.1517, which was 2 Da less than the value obtained for the parent drug M0. Secondary fragment ions at m/z 335.0230 [M-SO₃-H]⁻ and 255.0658 [M-2SO₃-H]⁻ were detected in the MS/MS spectrum, suggesting that hydrolyzed LQ had undergone a bis-sulfate conjugation reaction and that the formula was C₁₅H₁₂O₁₀S₂.

3.4.5. Metabolite M35. M35 had a retention time of 8.70 min with a deprotonated molecular ion $[M - H]^-$ at m/z497.0759, which was 80 Da higher than the value obtained for LQ, suggesting that M35 was a metabolite of sulfate conjugation. Meanwhile, characteristic fragment ions at m/z 417.1185 $[M-SO_3-H]^-$, 255.0655 $[M-SO_3-C_6H_{10}O_5-H]^-$ and 135.0074 (RDA reaction) were seen in the MS/MS spectrum, implying that the chemical formula was C21H22O12S.38

3.4.6. Metabolite M36. M36 exhibited a deprotonated molecular ion $[M - H]^-$ at m/z 511.0541, which was 94 Da more than the value obtained for deprotonated LQ. In addition a series of important fragment ions were observed at m/z $431.0970 [M-SO_3-H]^-$, $255.0654 [M-SO_3-C_6H_8O_6-H]^-$, 135.0084(RDA reaction) and 119.0506 (RDA reaction) in the MS/MS scan. Moreover, the retention time of M36 was 8.92 min, and the formula was identified to be C₂₁H₂₀O₁₃S.³⁸

3.4.7. Metabolites M44, M45. The QTOF-MS mass spectra of the M44 and M45 metabolites exhibited deprotonated ions $[M - H]^{-}$ at m/z 431.0976 and 431.0976, respectively, which were 14 Da higher than the value obtained for the parent drug LQ. In addition, both metabolites exhibited the loss of a glucuronide unit (176 Da) in the MS/MS spectra. A number of typical secondary fragment ions at m/z 255.0658, 135.0086 and

119.0503 were observed due to the loss of C₆H₈O₆ and due to RDA reaction. It was evident that hydrolyzed LQ had undergone glucuronide conjugation, and the molecular formula was identified to be C₂₁H₂₀O₁₀.38 M44 and M45 were eluted at 11.17 and 11.97 min, respectively. The clog P values of M44 and M45 were 0.27429 and 0.55864, respectively. Hence, M44 and M45 were identified on the basis of their retention times and clog Pvalues.

Metabolites M46, M47. The metabolites M46 and 3.4.8. M47, eluted at 12.13 and 12.78 min, respectively, were observed in the metabolic profiles with deprotonated molecular ions [M - H]⁻ at m/z 429.0828 and 429.0824, respectively, corresponding to an elemental composition of C₂₁H₁₈O₁₀. The masses of the deprotonated M46 and M47 were 12 Da more than the mass of LQ. A characteristic fragment ion at m/z 253.0505 [M- $C_6H_8O_6-H$ was observed, which confirmed that M46 and M47 were metabolites of glucuronide conjugation. Moreover, M46 and M47 were assigned based on clog P values of 0.390001 and 0.583201, respectively.

3.4.9. Metabolites M50, M51. M50 and M51 showed deprotonated molecular ions $[M - H]^-$ at m/z 335.0216 and 335.0217, respectively, which were 82 Da less than the value obtained for LQ. M50 and M51 were eluted at 12.49 and 17.30 min, respectively. Typical fragment ions at m/z 255.0656 [M-SO₃-H]⁻, 135.0083 (RDA reaction) and 119.0500 (RDA reaction) were observed in the MS/MS spectra. The neutral loss of SO₃ suggested that hydrolyzed LQ had undergone sulfate conjugation, implying that the molecular formula was $C_{15}H_{12}O_7S.^{38}$ By using ChemDraw 12.0, the clog P values of M50 and M51 were 0.689589 and 0.97394, respectively. Hence, these metabolites were identified based on the retention times and clog P values.

3.4.10. Metabolites M53, M54. Two peaks were eluted at 13.24 and 14.22 min with deprotonated molecular ions $[M - H]^{-}$ at m/z 413.0854 and 413.0853, respectively, which were 4 Da less

Fig. 5 Metabolic profile and proposed metabolic pathways of LQ in vitro and in vivo (a, b, c-possible chemical structure).

RSC Advances Paper

Fig. 5 (Contd.)

than the value obtained for the parent compound LQ. These metabolites had identical fragment ions at m/z 255.0666, 135.0087 and 119.0501. The fragmentation ion at m/z 255.0666 suggested that the position of oxygen loss was glucose and not flavone. Therefore, the molecular formula was deduced to be C₂₁H₁₈O₉. In addition, metabolites M53 and M54 were identified based on the clog P values of 0.29614and 0.34904, respectively.

3.4.11. Metabolite M55. M55 was eluted at a retention time of 16.56 min with a deprotonated molecular ion $[M - H]^-$ at m/z459.1297, which was 42 Da more than the value obtained for LQ. Masses for characteristic ions were observed at m/z 417.1198, 255.0645, 135.0070 and 119.0489, which were generated by the loss of CH₂CO and C₆H₁₀O₅ and by RDA reaction, suggesting that M55 was an acetylation metabolite. Moreover, the chemical formula of M55 was C23H24O10.

3.4.12. Metabolite M56. M56 eluted at 18.33 min and had a deprotonated ion $[M - H]^-$ at m/z 431.0992, which was 14 Da higher than the value obtained for M0, implying that methylation had occurred. The secondary fragment ions at m/z 255.0666 $[M-CH_2-C_6H_{10}O_5-H]^-$ and 135.0086 (RDA reaction) demonstrated that the chemical formula was C₂₂H₂₄O₉. Other metabolites were showed in ESI.†

3.5. Identification and characterization of in vitro metabolites

In this incubation system, fifteen metabolites were detected, including eleven phase I and four phase II metabolites. Furthermore, metabolites N1-N14 were identified in the in vivo metabolic study. However, N15 was not found in vivo bio-samples. The structures of all the metabolites are shown in Fig. 2.

3.5.1. Identification of phase I metabolites

Metabolites N1, N2, M3 and N4. N1-N4, with retention times of 8.62, 9.01, 9.51 and 14.03 min, exhibited sharp peaks of deprotonated molecular ions $[M - H]^-$ at m/z 433.1110, 433.1111, 433.1112 and 433.1109, respectively, which were 16 Da more than the value obtained for LQ, suggesting that N1-N4 were oxidation metabolites and corresponded to the molecular formula C21H22O10. In addition, typical fragment ions at m/z 271.0592, 243.0645 and 109.0289 were generated by the loss of C₆H₁₀O₅ and CO and by RDA reaction. Moreover, N1-N4 were identified based on the clog P values of N1-N4 at 0.300687, 0.500687, 0.558399 and 0.948399, respectively.

Metabolite N8. N8 was detected at 12.63 min and exhibited a sharp peak for a deprotonated molecular ion $[M - H]^-$ at m/z415.1011, which was 2 Da less than the value obtained for M0. Fragment ions at m/z 253.0518 [M-C₆H₁₀O₅-H]⁻ suggested that a desaturation reaction had occurred, which was consistent with a molecular formula of C21H20O9.

Metabolite N9. N9 was detected at 16.03 min. N9 exhibited a sharp peak for a deprotonated molecular ion $[M - H]^-$ at m/z255.0650, which was 162 Da lower than the value obtained for LQ, implying that a hydrolysis had occurred. The secondary fragment ions at m/z 135.0083 and 119.0500, caused by RDA reaction, confirmed that the molecular formula of N9 was C₁₅H₁₂O₄.38

Metabolite N10. N10 exhibited a deprotonated molecular ion $[M - H]^-$ at m/z 269.0437, which was 14 Da more than the value obtained for N9, indicating that the molecular formula of N10 was C₁₅H₁₀O₅. In addition, N10 had a retention time of 16.38 min, and the important fragment ions were at m/z135.0090 (RDA reaction) and 133.0287 (RDA reaction).

Metabolite N11. N11 showed a deprotonated molecular ion [M - H]⁻ at m/z 253.0496, which was 2 Da less than the value obtained for N9, suggesting that a desaturation reaction occurred for N9, and the molecular formula of N11 was predicted to be $C_{15}H_{10}O_4$. In addition, the secondary fragment ions at m/z **RSC Advances**

135.0077, 133.0284 and 117.0341, caused by RDA reaction, were consistent with the chemical structure of N11. Moreover, the retention time of N11 was 17.58 min.

3.5.2. Identification of phase II metabolites

Metabolite N12. N12 presented a deprotonated molecular ion $[M - H]^-$ at m/z 593.1565, which was 176 Da higher than the value obtained for LQ, suggesting that glucuronide conjugation had occurred. In addition, the molecular formula was determined to be C₂₇H₃₀O₁₅.38 Moreover, the MS/MS spectrum exhibited a series of typical fragment ions at m/z 417.1208 [M- $C_6H_8O_6-H_1^-$, 255.0659 $[M-C_6H_8O_6-C_6H_{10}O_5-H_1^-]$, 135.0078 (RDA reaction) and 117.0187 (RDA reaction). In addition, N12 was eluted at 7.62 min.

Metabolite N15. The molecular formula was determined to be $C_{27}H_{32}O_{14}$ based on the deprotonated molecular ion $[M - H]^{-}$ at m/z 579.1752, which was 162 Da higher than the value obtained for LQ. N15 was eluted at 12.45 min and had characteristic fragment ions at m/z 417.1273 [M-C₆H₁₀O₅-H]⁻, 255.0669 $[M-2C_6H_{10}O_5-H]^-$, 135.0085 (RDA reaction) and 119.0492 (RDA reaction), which implied that N15 was a glucose metabolite. Other metabolites were showed in ESI.†

3.6. Metabolic pathways of LQ

Based on the elemental compositions of the metabolites, the accurate MS/MS spectra, the chemical structures of the metabolites and the fragment ions of the metabolites, the metabolic pathway of LQ could be tentatively proposed. In total, 56 in vivo metabolites and 15 in vitro metabolites were characterized. Based on these results, LQ mainly underwent oxidation, reduction, hydrolysis, methylation, acetylation, glucuronide and sulfate conjugation. Based on the in vivo metabolic data, sulfate and glucuronide conjugation reactions were the major bio-transformations. However, based on the in vitro metabolic data, the metabolic pathways of LO were concentrated in phase I. The proposed metabolic pathways of LQ in vitro and in vivo are shown in Fig. 5.

3.7. Comparison of metabolic pathways in vitro and in vivo

Drug metabolism plays an important role in different areas of the pharmaceutical industry and in drug development and toxicology. The in vivo approach is quantitative and very effective in studies of drug metabolism. 49,50 The in vitro method is generally suitable for targeted studies and is often predictive of real hazards and risks.⁵¹ In this study, in vitro (rat liver microsomes) and in vivo (blood, urine, feces and bile in rats) metabolic profiles were investigated by UHPLC-Q-TOF-MS/MS. Consequently, a total of 56 in vivo metabolites and 15 in vitro metabolites were screened and characterized. The in vitro and in vivo metabolisms were concentrated in phase I and II, respectively. The metabolites detected in vitro were all detected in vivo as well. However, the isomer of N12 was detected in vivo, and N15 was not found in vivo. It was speculated that oxidation of N15 had occurred in vivo. Therefore, the isomer of N12 was observed in vivo, which suggested that the metabolism in vivo was more complex and elusive than that in vitro.

3.8. Comparison of metabolites identified in this study and in a previous study

A study of the metabolic pathways of LQ has been reported in China, which only detected 7 metabolites (M24, M27, M32 (N12), M35, M36, M44 and M45 (N13 and N14), and M51). However, in this study, 56 metabolites were detected in vivo and 15 metabolites were detected in vitro, including the 7 metabolites identified in the previous study. Moreover, the metabolic sites of two metabolites were not determined in the previous study, and there were some differences between this study and the previous study. In this manuscript, M32, M33 and M50, M51 were detected as two pairs of isomers; however, these metabolites were detected as a single chromatographic peak in the previous study. Moreover, in this study, M32 and M33 were both not detected in vitro (N12), and no isomers of M36 were detected in vivo. Furthermore, in this manuscript, the metabolic sites of M44 and M45 were identified based on the important clog Pvalues, which were the same as those of N13 and N14 in vitro. In addition, in this study, metabolite N15 was not detected in vivo; therefore, it was speculated that N15 had undergone oxidation and that metabolites M32 and M33 were found in vivo.

Conclusion

In conclusion, the UHPLC-Q-TOF-MS/MS method was first established to screen and identify the metabolites of LQ in vitro and in vivo. Based on the preceding data acquisition and mining strategy, twenty, fifty-four, forty-eight, thirty-seven and fifteen metabolites were detected in rat blood, urine, bile, feces and liver microsomes, respectively. The metabolic pathways of LQ were determined to be oxidation, reduction, hydrolysis, methylation, acetylation, and sulfate and glucuronide conjugation; these findings filled in the gaps remaining from the previous incomplete study. Key product ion (KPI) was used to aid the detection of metabolites of LQ. The results lay the foundation for active screening studies. In addition, this study demonstrated a powerful strategy for rapid screening and identifying metabolites and chemical constituents of traditional Chinese medicines.

Conflicts of interest

All the authors have declared no conflict of interest.

Acknowledgements

The work received financial support from the National Natural Science Foundation of China (No. 81473180).

References

- 1 M. N. Asl and H. Hosseinzadeh, Phytother. Res., 2008, 22,
- 2 B. Q. Fu, H. Li, X. R. Wang, F. S. C. Lee and S. F. Cui, J. Agric. Food Chem., 2005, 53, 7408-7414.

Paper

3 J. Cinatl, B. Morgenstern, G. Bauer, P. Chandra, H. Rabenau and H. W. Doerr, *Lancet*, 2003, **361**, 2045–2046.

- 4 T. Yokozawa, Z. W. Liu and C. P. Chen, *Phytomedicine*, 2000, **6**, 439–445.
- 5 J. He, L. Chen, D. Heber, W. Y. Shi and Q. Y. Lu, *J. Nat. Prod.*, 2006, **69**, 121–124.
- 6 G. Khaksa, M. E. Zolfaghari, A. R. Dehpour and T. Samadian, *Planta Med.*, 1996, **62**, 326–328.
- 7 F. Rauchensteiner, Y. Matsumura, Y. Yamamoto, S. Yamaji and T. Tani, *J. Pharm. Biomed. Anal.*, 2005, **38**, 594–600.
- 8 Y. X. Gao, B. F. Cheng, J. J. Lian, D. D. Guo, J. W. Qin, Y. B. Zhang, H. J. Yang, M. Wang, L. Wang and Z. W. Feng, *J. Funct. Foods*, 2017, 33, 142–148.
- 9 H. J. Kwon, H. H. Kim, Y. B. Ryu, J. H. Kim, H. J. Jeong, S. W. Lee, J. S. Chang, K. O. Cho, M. C. Rho, S. J. Park and W. S. Lee, *Bioorg. Med. Chem.*, 2010, 18, 7668–7674.
- 10 P. Montoro, M. Maldini, M. Russo, S. Postorino, S. Piacente and C. Pizza, *J. Pharm. Biomed. Anal.*, 2011, 54, 535–544.
- 11 X. Y. Meng, S. B. Yang, Z. F. Pi, F. R. Song, H. Y. Jiang and Z. Q. Liu, *J. Liq. Chromatogr. Relat. Technol.*, 2012, 35, 1538–1549.
- 12 C. F. Xue, S. Jiang, J. M. Guo, D. W. Qian, J. A. Duan and E. X. Shang, J. Chromatogr. B: Anal. Technol. Biomed. Life Sci., 2011, 879, 3901–3908.
- 13 C. Y. Li, L. W. Qi and P. Li, J. Pharm. Biomed. Anal., 2011, 55, 146–160.
- 14 G. G. Tan, M. Liu, X. Dong, S. Wu, L. Fan, Y. B. Qiao, Y. F. Chai and H. Wu, *J. Pharm. Biomed. Anal.*, 2014, 96, 187–196.
- 15 Y. Z. Zhang, F. Xu, J. Dong, J. Liang, Y. Hashi, M. Y. Shang, D. H. Yang, X. Wang and S. Q. Cai, *J. Pharm. Biomed. Anal.*, 2012, 70, 425–439.
- 16 M. P. Gonthier, C. Remesy, A. Scalbert, V. Cheynier, J. M. Souquet, K. Poutanen and A. M. Aura, *Biomed. Pharmacother.*, 2006, 60, 536–540.
- 17 A. H. Yang, J. X. Chen, Y. T. Ma, L. L. Wang, Y. W. Fan and X. He, *J. Pharm. Biomed. Anal.*, 2017, **141**, 200–209.
- 18 M. Y. Liu, S. H. Zhao, Z. Q. Wang, Y. F. Wang, T. Liu, S. Li, C. C. Wang, H. T. Wang and P. F. Tu, J. Chromatogr. B: Anal. Technol. Biomed. Life Sci., 2014, 949–950, 115–126.
- 19 H. Han, W. L. Zeng, C. Y. He, S. W. A. Bligh, Q. Liu, L. Yang and Z. T. Wang, *J. Mass Spectrom.*, 2014, **49**, 1108–1116.
- 20 K. Wang, L. W. Chai, X. C. Feng, Z. B. Liu, H. X. Liu, L. Q. Ding and F. Qiu, *J. Pharm. Biomed. Anal.*, 2017, 139, 73–86.
- 21 X. D. Cheng and M. G. Wei, *Molecules*, 2014, **19**, 18881–18896.
- 22 D. G. Wang, T. J. Hang, C. Y. Wu and W. Y. Liu, J. Chromatogr. B: Anal. Technol. Biomed. Life Sci., 2005, 829, 97–106.
- 23 H. L. Ma, Y. Liu, X. Mai, Y. J. Liao, K. Zhang, B. Liu, X. Xie and Q. L. Du, *J. Pharm. Biomed. Anal.*, 2016, **125**, 194–204.
- 24 K. Xiong, T. T. Gao, T. Zhang, Z. T. Wang and H. Han, J. Chromatogr. B: Anal. Technol. Biomed. Life Sci., 2017, 1065–1066, 1–7.
- 25 C. P. Yang, M. H. Liu, W. Zou, X. L. Guan, L. Lai and W. W. Su, J. Asian Nat. Prod. Res., 2012, 14, 68-75.

- 26 C. R. Deng, C. Y. Gao, X. H. Tian, B. Chao, F. Wang, Y. Zhang, J. T. Zou and D. C. Liu, *J. Funct. Foods*, 2017, 35, 332–340.
- 27 Y. Y. Zhao, S. P. Wu, S. M. Liu, Y. M. Zhang and R. C. Lin, *Chem.-Biol. Interact.*, 2014, **220**, 181–192.
- 28 Y. Y. Zhao and R. C. Lin, *Chem.-Biol. Interact.*, 2014, 215, 7–16.
- 29 Y. Y. Zhao, L. Zhang, F. Y. Long, X. L. Cheng, X. Bai, F. Wei and R. C. Lin, *Chem.-Biol. Interact.*, 2013, **201**, 31–38.
- 30 Y. Y. Zhao, J. Liu, X. L. Cheng, X. Bai and R. C. Lin, *Clin. Chim. Acta*, 2012, **413**, 642–649.
- 31 X. Wang, S. S. Johansen, M. K. K. Nielsen and K. Linnet, *Drug Test. Anal.*, 2017, **9**, 1137–1151.
- 32 J. M. Wang, Z. Yang and J. Lechago, *Biomed. Chromatogr.*, 2013, 27, 1463–1480.
- 33 S. Hegstad, S. Hermansson, I. Betner, O. Spigset and B. M. H. Falch, *J. Chromatogr. B*, 2014, **947–948**, 83–95.
- 34 J. J. Pitt, Clin. Biochem. Rev., 2009, 30, 19-34.
- 35 A. H. Wu, R. Gerona, P. Armenian, D. French, M. Petrie and K. L. Lynch, *Clin. Toxicol.*, 2012, **50**, 733–742.
- 36 H. H. Maurer, Ther. Drug Monit., 2010, 32, 324-327.
- 37 H. H. Maurer and M. R. Meyer, Arch. Toxicol., 2016, 90, 2161– 2172.
- 38 S. Q. Dong, H. R. Fan, Q. S. Li, G. L. Wei, W. H. Liu and D. Y. Si, *Chin. Tradit. Herbal Drugs*, 2014, 45, 2499–2505.
- 39 W. T. Liang, H. B. Huang and T. Liu, *Pharm. Today*, 2012, **22**, 13–16.
- 40 H. B. Li, Z. R. Li, Z. L. Wang and L. Y. Sun, *Prog. Biochem. Biophys.*, 1993, **20**, 402–403.
- 41 Y. H. Ma, W. W. Xie, T. T. Tian, Y. R. Jin, H. J. Xu, K. R. Zhang and Y. F. Du, *Anal. Biochem.*, 2016, 511, 61–73.
- 42 P. P. Jia, Y. Q. Zhang, Q. Y. Zhang, Y. P. Sun, H. T. Yang, H. Shi, X. X. Zhang and L. T. Zhang, *Biomed. Chromatogr.*, 2016, 30, 1498–1505.
- 43 M. Y. Liu, S. H. Zhao, Z. Q. Wang, Y. F. Wang, T. Liu, S. Li, C. C. Wang, H. T. Wang and P. F. Tu, J. Chromatogr. B: Anal. Technol. Biomed. Life Sci., 2014, 949–950, 115–126.
- 44 T. T. Tian, Y. R. Jin, Y. H. Ma, W. W. Xie, H. J. Xu, K. R. Zhang, L. T. Zhang and Y. F. Du, J. Chromatogr. B: Anal. Technol. Biomed. Life Sci., 2015, 1006, 80–92.
- 45 J. Liang, F. Xu, Y. Z. Zhang, S. Huang, X. Y. Zang, X. Zhao, L. Zhang, M. Y. Shang, D. H. Yang, X. Wang and S. Q. Cai, J. Pharm. Biomed. Anal., 2013, 83, 108–121.
- 46 J. F. Chen, Y. L. Song, X. Y. Guo, P. F. Tu and Y. Jiang, *Analyst*, 2014, **139**, 6474–6485.
- 47 L. L. Wang, M. M. Sang, E. W. Liu, P. O. Banahene, Y. Zhang, T. Wang, L. F. Han and X. M. Gao, *J. Pharm. Biomed. Anal.*, 2017, 140, 45–61.
- 48 K. W. Luo, Q. S. Shi and F. Feng, *J. Chromatogr. B: Anal. Technol. Biomed. Life Sci.*, 2017, **1040**, 260–272.
- 49 S. J. Fenwick and J. P. Scarth, *Drug Test. Anal.*, 2011, 3, 705–716.
- 50 J. P. Scarth, H. A. Spencer, S. E. Timbers, S. C. Hudson and L. L. Hillyer, *Drug Test. Anal.*, 2010, 2, 1–10.
- 51 I. Wilk-Zasadna, C. Bernasconi, O. Pelkonen and S. Coecke, *Toxicology*, 2015, **332**, 8–19.