


 Cite this: *RSC Adv.*, 2018, 8, 23919

 Received 30th April 2018  
Accepted 15th June 2018

DOI: 10.1039/c8ra03696a

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

# Synthesis of trifluoromethyl-containing isoindolinones from tertiary enamides *via* a cascade radical addition and cyclization process†

 Hui Yu, \* Mingdong Jiao, Xiaowei Fang and Pengfei Xuan

A radical trifluoromethylation reaction of tertiary enamides was investigated and trifluoromethyl-containing isoindolinones were prepared under mild conditions. Using  $\text{TMSCF}_3$  as a radical source,  $\text{PhI}(\text{OAc})_2$  as an oxidant and  $\text{KHF}_2$  as an additive, tertiary enamides were converted to isoindolinones *via* a cascade addition and cyclization process in moderate to good yields.

In recent years, trifluoromethyl-containing azaheterocycles have attracted much attention for their potential application in the fields of pharmaceutical and agricultural chemistry.<sup>1</sup> Thus, lots of efforts have been devoted to the synthesis of trifluoromethyl azaheterocycles,<sup>2</sup> and among these developed methods, radical cascade addition and cyclization has emerged as a remarkable strategy due to its unique properties such as economy and high efficiency. Unsaturated amides are commonly used substrates for this type of transformation, which could be attacked by a  $\text{CF}_3$  radical followed by intramolecular C–O, C–N, or C–C bond formation to give different kinds of trifluoromethyl azaheterocycles. Fu reported a metal-free trifluoromethylation of *N*-allylamides with  $\text{CF}_3\text{SO}_2\text{Na}$  for the synthesis of trifluoromethyl-containing oxazolines *via* oxytrifluoromethylation.<sup>3</sup> In the presence of copper salts, *N*-acyl-2-allylaniline could be converted to trifluoromethylated indolines in moderate to good yields *via* aminotrifluoromethylation process.<sup>4</sup> With Togni's reagent,<sup>5</sup>  $\text{TMSCF}_3$ ,<sup>6</sup>  $\text{CF}_3\text{SO}_2\text{Na}$ ,<sup>7</sup>  $\text{CF}_3\text{SO}_2\text{Cl}$ <sup>8</sup> and other reagents<sup>9</sup> as the  $\text{CF}_3$  source,  $\alpha$ ,  $\beta$ -unsaturated amides, tosyl amides, or imides underwent a tandem conversion to give trifluoromethyl-containing oxindoles or isoquinoline-1,3-diones by trifluoromethylation/arylation reaction. On the other hand, as a special type of unsaturated amide containing an active double bond, enamide also exhibited excellent reactivity in radical reactions.<sup>10</sup> In fact, trifluoromethylation of enamides has already been investigated, and in most cases trifluoromethylated alkenes were obtained as the main products.<sup>11</sup> To the best of our knowledge, the radical trifluoromethylation and cyclization of enamide still remains undeveloped.


Isoindolinones are important *N*-heterocyclic compounds necessary in organic and pharmaceutical chemistry, and these compounds are used widely as anticoagulants and tranquilizers such as aristolactam, pagoclone, and zopiclone.<sup>12</sup> To introduce a  $\text{CF}_3$  group into isoindolinones, Wang and co-workers explored a convenient way to the synthesis of trifluoromethyl-containing isoindolinones by radical aminotrifluoromethylation (Scheme 1a),<sup>13</sup> but this transformation only occurred for *N*-methoxybenzamides, and in case of *N*-alkylbenzamides trifluoromethylated alkenes were obtained as the major products. 1,1-disubstituted terminal alkenes were also not suitable substrates because of the competition between O-trapping and N-trapping process. Thus, development a new method for the synthesis of trifluoromethyl-containing isoindolinones is still in demand. Here in, as a continuation of our efforts on the radical modification of amide derivatives,<sup>14</sup> we wish to present our work on the synthesis of trifluoromethyl-containing isoindolinones using enamides as the start materials by radical carbon trifluoromethylation (Scheme 1b).

Initially, *N*-*n*-butyl-*N*-(2-propenyl) benzamide **1a** was chosen as the model substrate to optimize the reaction conditions of this radical carbontrifluoromethylation process. As shown in Table 1, the reaction of **1a** with  $\text{TMSCF}_3$  (4.0 equiv.) was firstly

School of Chemical Science and Engineering, Shanghai Key Lab of Chemical Assessment and Sustainability, Tongji University, 1239 Siping Road, Shanghai, 200092, P. R. China. E-mail: yuhui@tongji.edu.cn; Fax: +86 21 65981097; Tel: +86 21 65981097

† Electronic supplementary information (ESI) available: Detailed experimental procedures and characterization data for the products. See DOI: 10.1039/c8ra03696a

a) Previous work: aminotrifluoromethylation




b) This work: carbontrifluoromethylation




Scheme 1 Synthesis of trifluoromethyl-containing isoindolinones.






Table 3 Synthesis of the peroxide products<sup>a,b</sup>

<sup>a</sup> The reaction was performed with **3** (0.2 mmol),  $\text{KHF}_2$  (0.2 mmol),  $\text{TMSCF}_3$  (0.8 mmol),  $\text{PhI(OAc)}_2$  (0.8 mmol) in EtOAc (2.0 mL) under  $\text{N}_2$  at  $80^\circ\text{C}$  for 12 h in a sealed tube. <sup>b</sup> Isolated yields.

Scheme 2 Results of heterocyclic substrate **5a** and **5b**.

When *N*-*n*-butyl-*N*-(2-vinyl) benzamide **3a** was subjected to the reaction conditions, no isoindolinone was observed, and the main product was trifluoromethylated alkene (Table 3, **4a'**). Changing the *N*-protecting group to *n*-propyl or cyclohexyl also

caused the formation of trifluoromethylated alkenes (Table 3, **4b'**–**4c'**). It seemed that the substituents on the benzoyl group had significant influence on the reaction result. For example, substrate with methoxy group on the *para* position of the benzoyl moiety still gave trifluoromethylated alkene as the main product (Table 3, **4d'**), but substrate with methyl group on the *para* position led to a mixture of trifluoromethylated alkene and isoindolinone (Table 3, **4e/4e'**). However, substrate with methyl group on the *ortho* position or halides on the *para* position of the benzoyl group gave only isoindolinones as the main products (Table 3, **4f–4h**). Substrate with  $\text{NO}_2$  on the *para* position displayed low reactivity and no reaction occurred (Table 3, **4i**).



Scheme 3 Possible mechanism.



Heterocyclic substrate such as **5a** and **5b** was also examined, but no cyclization product could be found and trifluoromethylated alkene **6a** and **6b** was obtained as the only product (Scheme 2).

To gain insights into the reaction mechanism, a control experiment was carried out to elucidate the mechanism. When 1.0 equiv. TEMPO was added to the reaction, the yield of **2a** decreased significantly to 15%, which indicated the possibility of a radical pathway. Based on the control experimental result and the previous investigation on aryltrifluoromethylation of alkenes, plausible mechanism for our methodology is proposed in Scheme 2. In the presence of  $\text{KHF}_2$ ,  $\text{TMSCF}_3$  reacted with  $\text{PhI}(\text{OAc})_2$  to generate  $\text{CF}_3$  radical, then the  $\text{CF}_3$  radical attacked enamide **1** or **3** affording radical intermediate **A**. Depending on the structure of the substrate, intermediate **A** would be converted to trifluoromethyl-containing isoindolinone or trifluoromethylated alkene according to different pathways as followed: (path a) intramolecular cyclization of **A** gave the resulting radical **B** with an aryl ring, which was oxidized to intermediate **C** then underwent deprotonation to give rise to the final product **2a-r** or **4**; (path b) **A** was oxidized to intermediate **D** then underwent elimination to give trifluoromethylated alkene **2s**, **4'** or **6** (Scheme 3).

## Conclusions

In conclusion, we have demonstrated a simple, facile approach to trifluoromethyl-containing isoindolinones by radical addition and cyclization of enamides with moderate to good yields under mild conditions.  $\text{KHF}_2$  was found crucial for this cyclization process and further investigation into the mechanism is currently underway in our laboratory.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

Financial support from Tongji University (20123231) is gratefully acknowledged.

## Notes and references

- For reviews, see: (a) J. Wang, M. Sanchez-Rosello, J. L. Acena, C. del Pozo, A. E. Sorochinsky, S. Fustero, V. A. Soloshonok and H. Liu, *Chem. Rev.*, 2014, **114**, 2432; (b) A. A. Gakh and Y. Shermolovich, *Curr. Top. Med. Chem.*, 2014, **14**, 952.
- (a) Y. Zheng and J. A. Ma, *Adv. Synth. Catal.*, 2010, **352**, 2745; (b) A. Studer, *Angew. Chem., Int. Ed.*, 2012, **51**, 8950; (c) T. Koike and M. Akita, *Top. Catal.*, 2014, **57**, 967; (d) M. A. Honey, R. Pasceri, W. Lewis and C. J. Moody, *J. Org. Chem.*, 2012, **77**, 1396; (e) J. P. Bouillon, C. Ates, Z. Janousek and H. G. Viehe, *Tetrahedron Lett.*, 1993, **34**, 5075.
- J. Yu, H. Yang and H. Fu, *Adv. Synth. Catal.*, 2014, **356**, 3669.
- (a) J. S. Lin, X. G. Liu, X. L. Zhu, B. Tan and X. Y. Liu, *J. Org. Chem.*, 2014, **79**, 7084; (b) J. S. Lin, Y. P. Xiong, C. L. Ma, L. J. Zhao, B. Tan and X. Y. Liu, *Chem.-Eur. J.*, 2014, **20**, 1332; (c) H. Y. Zhang, W. Huo, C. Ge, J. Zhao and Y. Zhang, *Synlett*, 2017, **28**, 962.
- (a) P. Xu, J. Xie, Q. Xue, C. Pan, Y. Cheng and C. Zhu, *Chem.-Eur. J.*, 2013, **19**, 14039; (b) H. Egami, R. Shimizu and M. Sodeoka, *J. Fluorine Chem.*, 2013, **152**, 51; (c) Y. An, Y. Li and J. Wu, *Org. Chem. Front.*, 2016, **3**, 570; (d) W. Kong, M. Casimiro, E. Merino and C. Nevado, *J. Am. Chem. Soc.*, 2013, **135**, 14480; (e) N. Yang, Z. Li, L. Ye, B. Tan and X. Y. Liu, *Chem. Commun.*, 2016, **52**, 9052–9055.
- (a) L. Li, M. Deng, S. C. Zheng, Y. P. Xiong, B. Tan and X. Y. Liu, *Org. Lett.*, 2014, **16**, 504; (b) W. Fu, F. Xu, Y. Fu, C. Xu, S. Li and D. Zou, *Eur. J. Org. Chem.*, 2014, **4**, 709; (c) X. Mu, T. Wu, H. Wang, Y. Guo and G. Liu, *J. Am. Chem. Soc.*, 2012, **134**, 878; (d) Y. Wang, J. Qiu, D. Kong and F. Chen, *Synlett*, 2014, **25**, 1731.
- (a) L. Shi, X. Yang, Y. Wang, H. Yang and H. Fua, *Adv. Synth. Catal.*, 2014, **356**, 1021; (b) L. Zhang, Z. Li and Z. Q. Liu, *Org. Lett.*, 2014, **16**, 3688; (c) W. Wei, J. Wen, D. Yang, X. Liu, M. Guo, R. Dong and H. Wang, *J. Org. Chem.*, 2014, **79**, 4225; (d) R. Sakamoto, H. Kashiwagi, S. Selvakumar, S. A. Moteki and K. Maruoka, *Org. Biomol. Chem.*, 2016, **14**, 6417; (e) J. Liu, S. Zhuang, Q. Gui, X. Chen, Z. Yang and Z. Tan, *Eur. J. Org. Chem.*, 2014, **15**, 3196; (f) Q. Lu, C. Liu, P. Peng, Z. Liu, L. Fu, J. Huang and A. Lei, *Asian J. Org. Chem.*, 2014, **3**, 273; (g) F. Yang, P. Klumphu, Y. Liang and B. H. Lipshutz, *Chem. Commun.*, 2014, **50**, 936.
- (a) C. Liu, W. Zhao, Y. Huang, H. Wang and B. Zhang, *Tetrahedron*, 2015, **71**, 4344; (b) X. J. Tang, C. S. Thomason and W. R. Dolbier, *Org. Lett.*, 2014, **16**, 4594–4597; (c) L. Zheng, C. Yang, Z. Xu, F. Gao and W. Xia, *J. Org. Chem.*, 2015, **80**, 5730.
- (a) J. Y. Guo, R. X. Wu, J. K. Jin and S. K. Tian, *Org. Lett.*, 2016, **18**, 3850; (b) J. W. Beatty, J. J. Douglas, K. P. Cole and C. R. J. Stephenson, *Nat. Commun.*, 2015, **6**, 7919; (c) S. Tang, Z. H. Li, M. W. Wang, Z. P. Li and R. L. Sheng, *Org. Biomol. Chem.*, 2015, **13**, 5285.
- For review and selected examples, see: (a) M. X. Wang, *Chem. Commun.*, 2015, **51**, 6039; (b) P. Song, P. Yu, J. S. Lin, Y. Li, N. Y. Yang and X. Y. Liu, *Org. Lett.*, 2017, **19**, 1330; (c) Y. Tang, Y. Zhang, K. Wang, X. Li, X. Xu and X. Du, *Org. Biomol. Chem.*, 2015, **13**, 7084; (d) X. M. Xu, L. Zhao, J. P. Zhu and M. X. Wang, *Angew. Chem., Int. Ed.*, 2016, **55**, 3799; (e) T. Taniguchi, A. Ishita, M. Uchiyama, O. Tamura, O. Muraoka, G. Tanabe and H. Ishibashi, *J. Org. Chem.*, 2005, **70**, 1922; (f) L. He, L. Zhao, D. X. Wang and M. X. Wang, *Org. Lett.*, 2014, **16**, 5972; (g) G. K. Friestad and Y. Wu, *Org. Lett.*, 2009, **11**, 819; (h) C. H. Lei, D. X. Wang, L. Zhao, J. P. Zhu and M. X. Wang, *Chem.-Eur. J.*, 2013, **19**, 16981; (i) M. N. Zhao, Z. H. Ren, D. S. Yang and Z. H. Guan, *Org. Lett.*, 2018, **20**, 1287; (j) C. H. Lei, D. X. Wang, L. Zhao, J. P. Zhu and M. X. Wang, *J. Am. Chem. Soc.*, 2013, **135**, 4708.
- (a) R. Ding, Q. Zhang, Y. Xu and T. Loh, *Chem. Commun.*, 2014, **50**, 11661; (b) R. Rey-Rodriguez, P. Retailleau,



- P. Bonnet and I. Gillaizeau, *Chem.–Eur. J.*, 2015, **21**, 3572; (c) H. Jiang, C. Huang, J. Guo, C. Zeng, Y. Zhang and S. Yu, *Chem.–Eur. J.*, 2012, **18**, 15158.
- 12 For reviews, see: (a) A. Di Mola, L. Palombi and A. Massa, *Targets Heterocycl. Syst.*, 2014, **18**, 113; (b) A. Di Mola, L. Palombi and A. Massa, *Curr. Org. Chem.*, 2012, **16**, 2302.
- 13 K. Shen and Q. Wang, *Org. Chem. Front.*, 2016, **3**, 222.
- 14 (a) H. Yu and J. Shen, *Org. Lett.*, 2014, **16**, 3204; (b) H. Yu and J. Shen, *RSC Adv.*, 2015, **5**, 9815; (c) H. Yu, Y. Xu, R. Dong and Y. Fang, *Adv. Synth. Catal.*, 2017, **359**, 39.

