## **RSC Advances**



## CORRECTION

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2018, 8, 20275

## Correction: Ascorbic acid/Fe<sup>0</sup> composites as an effective persulfate activator for improving the degradation of rhodamine B

Xiangyu Wang,\*a Yi Du,a Huiling Liub and Jun Mab

DOI: 10.1039/c8ra90046a

www.rsc.org/advances

Correction for 'Ascorbic acid/Fe<sup>0</sup> composites as an effective persulfate activator for improving the degradation of rhodamine B' by Xiangyu Wang *et al.*, *RSC Adv.*, 2018, **8**, 12791–12798.

The authors regret that the unit on the *x*-axis of Fig. 1 was incorrectly written as "% wt" rather than " $\frac{1}{0}$ " wt" in the original article. The correct version of Fig. 1 is presented below.

<sup>&</sup>lt;sup>a</sup>Faculty of Environmental Science and Engineering, Kunming University of Science and Technology, Kunming 650500, P. R. China. E-mail: imusthlee2014@sina.com

<sup>b</sup>School of Municipal and Environmental Engineering, State Key Laboratory of Urban Water Resources and Environment, Harbin Institute of Technology, Harbin 150090, P. R. China






Fig. 1 (a) Comparison of removal efficiency of RhB in different systems ( $C_0 = 50 \text{ mg L}^{-1}$ , PS dosage  $= 1.4 \text{ g L}^{-1}$ , Fe $^0$  dosage  $= 1 \text{ g L}^{-1}$ , H<sub>2</sub>A/Fe $^0$  dosage  $= 1 \text{ g L}^{-1}$ , H<sub>2</sub>A dosage  $= 1.6 \text{ g L}^{-1}$  and T = 298 K); (b) effect of H<sub>2</sub>A concentration on removal efficiency of RhB in the H<sub>2</sub>A/Fe $^0$ -PS system ( $C_0 = 50 \text{ mg L}^{-1}$ , Fe $^0$  dosage  $= 0.8 \text{ g L}^{-1}$ , T = 298 K and the solution volume is 50 mL).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.