


## CORRECTION

[View Article Online](#)  
[View Journal](#) | [View Issue](#)
Cite this: *Chem. Sci.*, 2018, **9**, 531

DOI: 10.1039/c7sc90073b

[www.rsc.org/chemicalscience](http://www.rsc.org/chemicalscience)

The authors regret that Fig. 4 is incorrect in the original manuscript. In Fig. 4c the chemical structure and mass spectrum of the norbornene-modified adenosine was shown instead of the 2'-deoxyadenosine. The correct figure and caption are displayed below.



**Fig. 4** Norbornene modification of pBR322 plasmid DNA using the  $N^6$ -adenine MTase M. TaqI. (A) Scheme for the functionalization of plasmid DNA using norbornene-modified AdoMet analog **1b**. (B) Fluorescence labeling of plasmid DNA via norbornene-modification followed by labeling with TAMRA-tetrazine and linearization of the plasmid using BamHI. Bands were resolved on a 1% agarose gel (100 V, 50 min), the gel was stained using ethidium bromide and scanned on a Typhoon FLA9500 laser scanner. (C) Mass spectrometric analysis of  $N^6$ -norbornene-modified oligonucleotides. A DNA oligonucleotide was subjected to enzymatic norbornene-modification, followed by digestion using nuclease P1 and dephosphorylation using FastAP (ThermoFisher Scientific). Expected mass for  $C_{26}H_{31}N_6O_4^+ = 491.2401$   $[M + H]^+$ , found: 491.2400. M: GeneRuler 1 kb DNA ladder (ThermoFisher).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>University of Münster, Department of Chemistry, Institute of Biochemistry, Wilhelm-Klemm-Str. 2, 48149 Münster, Germany

<sup>b</sup>Cells-in-Motion Cluster of Excellence (EXC1003-CiM), University of Münster, Germany. E-mail: [a.rentmeister@uni-muenster.de](mailto:a.rentmeister@uni-muenster.de)