# **Chemical Science**

# EDGE ARTICLE

Cite this: Chem. Sci., 2018, 9, 2991

# Gold-catalyzed [4+3]- and [4+2]-annulations of 3-en-1-ynamides with isoxazoles via novel  $6\pi$ electrocyclizations of 3-azahepta trienyl cations†

New gold-catalyzed [4+3]-annulations of 3-en-1-ynamides with isoxazoles afford 4H-azepines efficiently; this process involves 6 $\pi$  electrocyclizations of gold-stabilized 3-azaheptatrienyl cations. In the presence of Zn(OTf)2, the resulting 4H-azepines undergo skeletal rearrangement to furnish substituted pyridine derivatives. We subsequently develop new catalytic [4+2]-annulations between the same 3-en-1 ynamides and isoxazoles to deliver substituted pyridine products using Au(I)/Zn(II) catalysts. This work reports the first success of the  $6\pi$  electrocyclizations of heptatrienyl cations that are unprecedented in

Sovan S[u](http://orcid.org/0000-0002-2011-8124)ndar Giri and Rai-Shung Liu D<sup>\*</sup>

Received 15th January 2018 Accepted 18th February 2018

DOI: 10.1039/c8sc00232k

rsc.li/chemical-science

# Introduction

Electrocyclizations of acyclic conjugated  $\pi$ -motifs are powerful tools to access five-, six- and seven-membered carbocycles;<sup>1</sup> prominent examples include Nazarov cyclizations of pentadienyl cations<sup>2</sup> and  $6\pi$  electrocyclizations of trienes,<sup>3</sup> which have found widespread applications in organic synthesis.

literature reports.



Scheme 1 Electrocyclizations of conjugated  $\pi$ -motifs.

In the context of seven-carbon  $\pi$ -motifs, heptatrienyl anions I undergo facile  $8\pi$  electrocyclizations via rapid interconversions among various anion configurations (Scheme 1).<sup>4</sup> In contrast, heptatrienyl cations III<sup>5</sup> exclusively undergo Nazarov reactions because of the difficulties of forming all  $\sigma$ -cis configured cations V that have a high energy state.<sup>5b</sup> 1-Aza- and 1oxaheptatrienyl cations<sup>6</sup> were also reported to follow Nazarov cyclizations. The realization of a  $6\pi$  electrocyclization of conjugated seven-membered cations is formidable but challenging. This work reveals the first success of such sevenmembered cyclizations of gold-stabilized 3-azaheptatrienyl cations  $V'$  to form azacyclic products 3–4 via a new C–C bond formation. **EDGE ARTICLE**<br> **(a)** Check for updates<br> **Contained and the Contained School Catalyzed [4+3]- and [4+2]-annulations of<br>
Check for updates<br>
Contained and 2.2013 Soviet Control Control Control Control Control Control Contro** 

YAL SOCIETY<br>**CHEMISTRY** 



The advent of gold catalysis has inspired new annulations between alkynes and poor nucleophiles.<sup>7</sup> N-O containing nucleophiles serve as useful building blocks to construct valuable azacyclic frameworks.<sup>7</sup> Ye and Hashmi reported interesting [3+2]-annulations of isoxazoles or benzisoxazoles with electron-

Department of Chemistry, National Tsing-Hua University, Hsinchu, Taiwan, Republic of China. E-mail: rsliu@mx.nthu.edu.tw

<sup>†</sup> Electronic supplementary information (ESI) available. CCDC 1589549, 1589562, 1589561, 1589558, 1589559 and 1589560. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8sc00232k

rich ynamides, yielding substituted pyrrole derivatives through aza-Nazarov cyclizations of the key intermediate  $[eqn(1)]^{7,8}$ These [3+2]-annulations were extensively expanded to other N–O heterocycles including benzisoxazoles, 1,2,4-oxadiazoles, 1,4,2-dioxazoles and 4,5-dihydro-1,2,4-oxadiazoles, yielding additional five-membered azacycles as depicted in  $[eqn(1)]$ .<sup>9</sup> Here, we report two distinct  $[4+3]$ - and  $[4+2]$ -annulations between 3-en-1-ynamides and isoxazoles using varied catalysts. An Au(I) catalyst alone delivers 4H-azepines 3-4 through  $6\pi$  electrocyclizations of intermediates V [eqn (2)] whereas a combined action of  $Au(I)/Zn(I)$  on the same reactants furnishes highly functionalized pyridines 5 [eqn (3)]. With our convenient synthesis, the synthetic utility of new 4H-azepines 3– 4 is also reported.<sup>10</sup>

#### Results and discussion

We examined the reactions of 3-methyl-3-en-1-ynamide 1a with 3,5-dimethylisoxazole 2a using various gold catalysts. Heating this mixture  $(1a/2a = 1 : 2 \text{ ratio})$  in hot DCE with 5 mol% LAuCl/ AgNTf<sub>2</sub>  $[L = p(t-Bu)$ <sub>2</sub>(*o*-biphenyl) and IPr] afforded a [4+3]annulation product, 4H-azepine 3a, in 64% and 75% yields respectively (Table 1, entries 1–2). Under these conditions, a low loading (1.2 equiv.) of 3,5-dimethylisoxazole 2a gave 3a in a decreased yield, ca. 62% (entry 3). With a 10 mol% catalyst, IPrAuCl/AgNTf<sub>2</sub> gave a clean reaction, yielding desired 3a up to 91% (entry 4). We tested other phosphine ligands such as  $PPh<sub>3</sub>$ and P(OPh)3, yielding desired 3a in satisfactory yields (78–81%, entries 5–6). Other counter anions such as  $\mathrm{OTf}^-$  and  $\mathrm{SbF_6}^-$  were also effective in producing 3a in 85–88% yields (entries 7–8). AgNTf<sub>2</sub> alone was not active at all (entry 9).



<sup>a</sup> [1a] = 0.15 M. <sup>b</sup> Product yields are reported after separation from a silica column. <sup>c</sup> L =  $p(t-Bu)_2(o-bipheny)$ . <sup>d</sup> IPr = 1,3 $c$  L =  $p(t-Bu)_{2}(o\text{-biphenyl}).$ bis(diisopropylphenyl)-imidazol-2-ylidene. Ms  $=$  methanesulfonyl,  $DCE = 1,2$ -dichloroethane, and Tf = trifluoromethanesulfonyl.

Table 2 [4+3]-Annulations with various 3-en-1-ynamides



 $a^{\dagger}$  [1] = 0.15 M.  $b^{\dagger}$  Product yields are reported after separation from a silica column. EWG  $=$  electron withdrawing group.

Table 3 [4+3]-Annulations with various isoxazoles

1b <sup>a</sup>	$\mathsf{R}^1$ N $R^2$ Ts $2(2)$ equiv)		10 mol % IPrAuCl/AgNTf <sub>2</sub> DCE, 70 °C $R^2$ time	Тs $R^1$ 4 <sup>b</sup> 7a'	N Ts
Entry	$(R^1, R^2)$	2	Time [h]	Yield $[\%]$	$\boldsymbol{4}$
(1)	H, H	2 <sub>b</sub>	4	84	$4a$ (X-ray)
				8	7a'
(2)	H, Me	2d	3	75	4b
(3)	Me, H	2c	3	87	4c
$\left( 4\right)$	Et, Et	2e	6	85	4d
(5)	$n$ -Bu, $n$ -Bu	2f	7	81	4e
(6)	Me, $n$ -Bu	2g	3	82	4f
(7)	$n$ -Bu, $c$ -Pr	2 <sub>h</sub>	$\overline{2}$	77	4g
(8)	Ph, $n$ -Bu	2i	$\overline{4}$	69	4h
(9)	Ph, Ph	2j	6.5	61	4i
				30	$5i$ (X-ray)
(10)	Me, Ph	2k	4	71	4j
	Ph	Ts	$R^1$ = Ph (5i) $R^1$ = Me (5j)	15	5j

 $[\mathbf{1b}] = 0.15$  M.  $^b$  Product yields are reported after separation from a silica column.

Suitable substituents of 3-en-1-ynamides 1 are crucial to achieve  $6\pi$  cyclizations of 3-azaheptatrienyl cations V' [eqn (2)]. We tested the reactions on 3-en-1-ynes **1b-1m** bearing a C(3)substituent to circumvent aza-Nazarov cyclizations as reported in Ye's work.<sup>7</sup> Herein, only entries 9 and 10 showed the presence of 3-azanorcaradienes  $3^{\prime}.$  We examined these [4+3]-annulations on 3-methyl-3-en-1-ynamides 1b–1e bearing various sulfonamides NTsR<sup>4</sup> ( $R^4$  = Me, cyclopropyl, benzyl and N(n-C<sub>4</sub>H<sub>9</sub>)  $(-SO_2Bu)$ ), affording the desired 4H-azepines 3b-3e in high yields (84–90%, Table 2, entries 1–4). Nevertheless, this new annulation becomes less efficient for 3-en-1-ynamide 1f bearing an oxazolidin-2-one to yield product 3f in 64% yield (entry 5).

We altered the  $C(3)$ -substituents as in substrates  $1g-1i$ ; their resulting products 3g–3h ( $R^1$  = isopropyl and cyclopropyl) were obtained in 74–79%, and 3i ( $R^1$  = Ph) with only 58% yield (entries 6–8). Notably, when a long n-butyl group was present as in species 1j and 1k, their corresponding reactions afforded compounds  $3j/3j' = 5/1$  and  $3k/3k' = 11.1$ : 1, respectively, in 55% and 68% yields (entries 9-10). For E-configured trisubstituted 3-en-1-yne 1l ( $R^1$  = Me,  $R^2$  = Ph and  $R^3$  = H), 4H-azepine 3l



and pyrrole 6l were obtained in equal proportions (entry 11). When a cyclohexenyl group was present for alkene as in species 1m, pyrrole product 6m was dominant over azepine 3m (entry 12). Accordingly, preferable 3-en-1-ynes comprise a small  $R^2$  or  $R<sup>3</sup>$  substituent whereas  $R<sup>1</sup>$  must be substituted. Herein, the structures of 4H-azepines 3b and 3l, and pyrrole species 6m were confirmed with X-ray diffraction.<sup>11</sup>

Isoxazoles of a wide scope are compatible with these [4+3] annulations, as depicted in Table 3. The reaction of unsubstituted isoxazole 2b with model 3-en-1-ynamide 1b afforded the desired  $4H$ -azepine  $4a$  in 84% yield, together with pyrrole  $7a'$  in only 8% yield (entry 1). Mono-substituted 3-methyl or 5-methyl isoxazoles 2c and 2d are also suitable for these annulations to afford compounds 4b and 4c in 75% and 87% yields, respectively (entries 2–3). We prepared additional 3,5-disubstituted isoxazoles 2e–2i with  $R^1$  = alkyl and phenyl, and  $R^2$  = alkyl; their annulations proceed smoothly to produce desired 4d–4h in 69– 85% yields (entries 4–8). For di-substituted isoxazoles 2j and 2k bearing  $R^2 = Ph$ , 4H-azepines 4i and 4j were obtained in 61% and 71% yields respectively, together with their rearrangement products 5i and 5j in 15–30% yields (entries 9–10). Compounds 4a and 5i were characterized by X-ray diffraction.<sup>11</sup> Edge Article<br>
Suitable substitutents of 2-cont-published on 19 February 2018. Download Science Systems Article in the System Common Commo

Our convenient synthesis of 4H-azepines provides new synthetic utilities; several new functionalizations are depicted in Scheme 2. NaBH<sub>4</sub>-reduction of species 3b delivered an alcohol derivative 7a in 84% yield. Selective hydrogenation of the same species afforded 2-aza-1,3-dien-5-one 7b in 71% yield. A final treatment of  $4H$ -azepine 3b with NBS in acetone afforded compound 7c, of which the molecular structure was determined by <sup>1</sup>H NOE spectra.

The Lewis-catalyzed rearrangement of 4H-azepines 3–4 to substituted pyridines 5 [eqn (3)] is unprecedented in 4H-azepine Scheme 2 New functionalization of 4H-azepines. <br>
Scheme 2 New functionalization of 4H-azepines.<br>
Scheme 2 New functionalization of 4H-azepines.

Table 4 [4+2]-Annulations between 3-en-1-ynamides and isoxazoles





 $a [1] = 0.15$  M.  $b$  Product yields are reported after separation from a silica column.  $c$  The value in parentheses is reported using a mixture of IPrAuCl/ AgNTf<sub>2</sub> (10 mol%) and Zn(OTf)<sub>2</sub> (20 mol%) in hot DCE (70 °C, 48 h); 3b was also isolated in 28% yield.



Scheme 3 A plausible reaction mechanism

between 3-en-1-ynamides 1 and isoxazoles 2 using  $Au(i)/Zn(n)$  in a relay series, as depicted in Table 4. In the reactions of various 3-substituted 3-en-1-ynamides 1 ( $R^1$  = methyl, *n*-butyl, cyclopropyl and isopropyl) with 3,5-dimethylisoxazole 2a, substituted pyridines 5a–5d were obtained in satisfactory yields (51–73%, entries 1–4). In entry 1, if the reaction was performed with combined  $Au(1)/Zn(n)$  catalysts in a non-relay operation, compounds 5a and 3b were isolated in 35% and 28% yields respectively. For 3-en-1-ynamide 1a bearing a  $NMs(n-butyl)$ , the corresponding product 5e was obtained in 63% yield (entry 5). We tested the reactions on 3,5-disubstituted isoxazoles 2e–2f & 2h bearing all alkyl substituents, producing desired 5f–5h in good yields (69–78%, entries 6–8). For such disubstituted isoxazoles bearing  $R^4$  = Ph, the reactions afforded the desired pyridine derivatives 5i and 5j in 75–80% yields (entries 9–10). The molecular structures of compounds 5a and 5i were characterized by X-ray diffraction.<sup>11</sup>

Scheme 3 rationalizes the crucial roles of substituents of 3 en-1-ynamides in the chemoselectivity that relies on two conformational structures **D** versus **D** $^{\prime}$ . The N-attack of isoxazole at gold- $\pi$ -ynamide **A** is expected to form a gold-carbene  $\mathbf{D}',$ which can be visualized as a gold-stabilized cycloheptatrienyl cation. Conformation **D** is favorable with  $R = H$ , which prefers aza-Nazarov reactions.<sup>12</sup> When a C(3)-substituent is present (R = alkyl and aryl), all  $\sigma$ -cis configured species  $D'$  are the preferable geometry to induce novel  $6\pi$  electrocyclizations. This ring closure is expected to proceed through an attack of enamide at the alkenylgold moiety that is also visualized as a goldstabilized cation. Additional C(4)-substituents render the formation of cations  $D'$  difficult, thus yielding pyrrole 6 as byproducts. A loss of an acidic proton from seven-membered cations E is expected to yield azepines 3–4. 4H-Azepines 3–4 bear an enone conjugated with a triene; this extensive conjugation is very stable to impede a  $6\pi$  electrocyclization of their triene moieties unless a Lewis acid is present.  $\text{Zn}(\text{OTf})_2$  likely coordinates with the carbonyl of 4H-azepine 3 to generate a 2 azapentadienyl cation F bearing a zinc enolate, further enabling an intramolecular cyclization to generate species G. A 1,2-acyl shift<sup>14</sup> of species G delivers the observed product  $5.^{13}$ 

# **Conclusions**

In summary, this work describes new gold-catalyzed [4+3] annulations<sup>15</sup> of 3-substituted 3-en-1-ynamides with isoxazoles to form 4H-azepines. A relay catalysis is also developed with Au( $I$ )/Zn( $II$ ) catalysts to achieve [4+2] annulations from the same reactants. The mechanisms of gold-catalyzed [4+3] annulations involve unprecedented  $6\pi$  electrocyclizations of 3-azacycloheptatrienyl cations to form 4H-azepines 3–4 efficiently. Control experiments confirm that  $4H$ -azepines  $3-4$  are catalyzed by  $Zn(OTf)<sub>2</sub>$  to undergo new rearrangement reactions to form substituted pyridine derivatives.

# Conflicts of interest

The authors declare no conflict of interest.

# Acknowledgements

The authors thank the Ministry of Science and Technology and the Ministry of Education, Taiwan, for supporting this work.

# Notes and references

- 1 (a) S. Sankararaman, in Pericyclic Reactions-A Textbook. Reactions, Applications and Theory, Wiley-VCH, New York, 2005, ch. 6, pp. 393–398; (b) C. M. Beaudry, J. P. Malerich and D. Trauner, Chem. Rev., 2005, 105, 4757; (c) P. V. R. Schleyer, J. I. Wu, F. P. Cossio and I. Fernandez, Chem. Soc. Rev., 2014, 43, 4909; (d) M. Bian, L. Li and H. Ding, Synthesis, 2017, 49, 4383; (e) E. C. Taylor and I. J. Turchi, Chem. Rev., 1979, 79, 181; (f) F. D. Proft, P. K. Chattaraj, P. W. Ayers, M. T. Sucarrat, M. Elango, V. Subramanian, S. Giri and P. Geerlings, J. Chem. Theory Comput., 2008, 4, 595;  $(g)$  N. Jana and G. Driver, Org. Biomol. Chem., 2015, 13, 9720.
- $2(a)$  W. T. Spencer III, T. Vaidya and A. J. Frontier, *Eur. J. Org.* Chem., 2013, 3621; (b) R. L. Davis and D. L. Tantillo, Curr. Org. Chem., 2010, 14, 1561; (c) T. N. Grant, C. J. Rieder and

F. G. West, Chem. Commun., 2009, 5676; (d) M. J. Riveira, L. A. Marsili and M. P. Mischne, Org. Biomol. Chem., 2017, 15, 9255; (e) N. Shimada, C. Stewart and M. A. Tius, Tetrahedron, 2011, 67, 5851.

- 3 (a) V. A. Guner, K. N. Houk and I. W. Davis, J. Org. Chem., 2004, 69, 8024; (b) R. V. Essen, D. Frank, H. W. Sunnemann, D. Vidovic, J. Magull and A. D. Meijere, Chem.–Eur. J., 2005, 11, 6583; (c) N. A. Magomedov, P. L. Ruggiero and Y. Tang, J. Am. Chem. Soc., 2004, 126, 1624; (d) E. N. Marvell, G. Caple, B. Schatz and W. Pippin, Tetrahedron, 1973, 29, 3781; (e) J. R. Otero, J. Org. Chem., 1999, 64, 6842; (f) P. E. Tessier, N. Nguyen, M. D. Clay and A. G. Fallis, Org. Lett., 2005, 7, 767; (g) C. L. Benson and F. G. West, Org. Lett., 2007, 9, 2545; (h) G. A. Barcan, A. Patel, K. N. Houk and O. Kwon, Org. Lett., 2012, 14, 5388. 4 (a) R. B. Bates, W. H. Delnes, D. A. McCombs and D. E. Potter, J. Am. Chem. Soc., 1969, 91, 4608; (b) K. Marx and W. Eberbach, Chem.–Eur. J., 2000, 6, 2063; (c) M. Reisser and G. Mass, J. Org. Chem., 2004, 69, 4913; (d) T. Hübner, *Dissertation*, University of Freiberg, 1987; (e) A. Arany, D. Bendell, P. W. Groudwater, I. Garnett and M. Nyerges, J. Chem. Soc., Perkin Trans. 1, 1999, 2605. Edge Article<br>
F. G. West, Chem, Consume, 2009, 5676; (d) M. J. Riveria, H. R. Ducha, A. E. Ayeh and M. A. Iyach, Chem, Sc. (c) A. Chem,
	- 5 (a) N. C. Deno, C. U. Pittman and J. O. Turner, J. Am. Chem. Soc., 1965, 87, 2153; (b) O. N. Faza, C. S. López, R. Alvarez and A. R. de Lera, Chem.–Eur. J., 2009, 15, 1944.
	- 6 D. Alickmann, R. Fröhlich, A. H. Maulitz and E. U. Wurthwein, Eur. J. Org. Chem., 2002, 1523.
	- 7 (a) A. H. Zhou, Q. He, C. Shu, Y. F. Yu, S. Liu, T. Zhao, W. Zhang, X. Lu and L. W. Ye, Chem. Sci., 2015, 6, 1265; (b) X. Y. Xiao, A. H. Zhou, C. Shu, F. Pan, T. Li and L. W. Ye, Chem.–Asian J., 2015, 10, 1854; (c) W. B. Shen, X. Y. Xiao, Q. Sun, B. Zhou, X. Q. Zhu, J. Z. Yan, X. Lu and L. W. Ye, Angew. Chem., Int. Ed., 2017, 56, 605; (d) L. Li, T. D. Tan, Y. Q. Zhang, X. Liu and L. W. Ye, Org. Biomol. Chem., 2017, 15, 8483.
	- 8 (a) H. Jin, L. Huang, J. Xie, M. Rudolph, F. Rominger and A. S. K. Hashmi, Angew. Chem., Int. Ed., 2016, 55, 794; (b) H. Jin, B. Tian, X. Song, J. Xie, M. Rudolph, F. Rominger and A. S. K. Hashmi, Angew. Chem., Int. Ed., 2016, 55, 12688.
	- 9 (a) Z. Zeng, H. Jin, J. Xie, B. Tian, M. Rudolph, F. Rominger and A. S. K. Hashmi, Org. Lett., 2017, 19, 1020; (b) M. Chen, N. Sun, H. Chen and Y. Liu, Chem. Commun., 2016, 52, 6324; (c) W. Xu, G. Wang, N. Sun and Y. Liu, Org. Lett., 2017, 19, 3307.
	- 10 For synthesis of  $nH$ -azepines  $(n = 2 \text{ or } 4)$ , see selected examples: (a) U. Gockel, U. Hartmannsgruber, A. Steigel and J. Sauer, Tetrahedron Lett., 1980, 21, 599; (b)

I. R. Dunkin, A. E. Ayeb and M. A. Lynch, J. Chem. Soc., Chem. Commun., 1994, 1695; (c) K. Satake, R. Okuda, M. Hashimoto, Y. Fujiwara, H. Okamoto, M. Kimura and S. Morosawa, J. Chem. Soc. Perkin Trans 1, 1994, 1753; (d) Y. Luo and J. Wu, Chem. Commun., 2011, 47, 11137.

- 11 Crystallographic data of compounds 3b, 3l, 4a, 5a, 5i, and 6m were deposited in Cambridge Crystallographic Data Center: 3b: CCDC 1589549, 3l: CCDC 1589562, 4a: CCDC 1589561, 5a: CCDC 1589558, 5i: CCDC 1589559 and 6m CCDC 1589560.†
- 12 For the aza-Nazarov cyclizations, see: (a) D. A. Klumpp, Y. Zhang, M. J. O'Connor, P. M. Esteves and L. S. de Almeida, Org. Lett., 2007, 9, 3085; (b) Z. X. Ma, S. He, W. Song and R. P. Hsung, Org. Lett., 2012, 14, 5736; (c) R. L. Sahani and R. S. Liu, Angew. Chem., Int. Ed., 2017, 56, 12736; (d) S. K. Pawar, R. L. Sahani and R. S. Liu, Chem.– Eur. J., 2015, 21, 10843; (e) C. Shu, Y. H. Wang, C. H. Shen, P. P. Ruan, X. Lu and L. W. Ye, Org. Lett., 2016, 18, 3254.
- 13 As suggested by one reviewer, an alternative mechanism is also possible for the  $Zn(\pi)$ -catalyzed rearrangement; this process involves an isomerization of initial species 3 to an unconjugated iminoyl ketone H, followed by a  $6\pi$ cyclization to generate species I. A subsequent  $Zn(n)$ catalyzed aromatization of species I is expected to yield the final product 5. In this process, species  $H$  is relatively higher than 3 in energy, but its feasibility is not excluded.



- 14 (a) S. M. Wang and L. Zhang, Org. Lett., 2006, 8, 4585; (b) G. Li, G. Zhang and L. Zhang, J. Am. Chem. Soc., 2008, 130, 3704; (c) S. B. Wagh and R. S. Liu, Chem. Commun., 2015, 51, 15462; (d) R. Chaudhuri, A. Das, H. Y. Liao and R. S. Liu, Chem. Commun., 2010, 46, 4601.
- 15 For reviews of gold-catalyzed cycloaddition or annulation reactions of alkynes,  $\sec(a)$  D. J. Gorin, B. D. Sherry and F. D. Toste, Chem. Rev., 2008, 108, 3351; (b) A. S. K. Hashmi, Chem. Rev., 2007, 107, 3180; (c) F. López and J. L. Mascareñas, Beilstein J. Org. Chem., 2011, 7, 1075; (d) C. Aubert, L. Fensterbank, P. Garcia, M. Malacria and A. Simonneau, Chem. Rev., 2011, 111, 1954; (e) D. Garayalde and C. Nevado, ACS Catal., 2012, 2, 1462; (f) M. E. Muratore, A. Homes, C. Obradors and A. M. Echavarren, Chem.–Asian J., 2014, 9, 3066.