# Chemical Science

# **EDGE ARTICLE**

Check for updates

Cite this: Chem. Sci., 2018, 9, 7498

o All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 28th June 2018 Accepted 27th July 2018

DOI: 10.1039/c8sc02851f

rsc li/chemical-science

### Introduction

Hierarchical assembly is involved in many processes.<sup>1,2</sup> These processes use recognition and supramolecular matching of chemical pairs to help with the organization.<sup>3-5</sup> In chemistry the acquired knowledge from biology is used in the design of organic dendrimers, polymers and molecular machines and extended to coordination compounds as well as metal-organic frameworks.6-10 In physics the modelling of nanostructures on surfaces is a major active area of development as is patterning for logic circuits.<sup>11</sup> Although hierarchical assembly processes have been recognized in chemistry, tandem ones are rarely evoked.12,13

In the course of studying the coordination complexes obtained from different copper salts with planar multidentate 5,5'pyridyl-3,3'-bi-1*H*-pyrazole, rich in  $\pi$ -electrons, by solvothermal reactions in different solvents, we have isolated several series of complexes of variable nuclearities and valencies. One series contains rugby-ball hexamers where the valence of the copper ions varies from Cu<sub>6</sub><sup>II</sup> to Cu<sup>I</sup>Cu<sub>5</sub><sup>II</sup> and Cu<sub>2</sub><sup>I</sup>Cu<sub>4</sub><sup>II</sup> depending on the

## Hierarchical tandem assembly of planar $[3 \times 3]$ building units into $\{3 \times [3 \times 3]\}$ oligomers: mixedvalency, electrical conductivity and magnetism<sup>+</sup>

Fei Yu, 🕩 a Mohamedally Kurmoo, ២ Gui-Lin Zhuang 🕩 c and Jing-Lin Zuo 🕩 \*a

A hierarchical tandem assembly - coordination, oligomerisation and supramolecular organization - is identified using MALDI-TOF of the solid products of the solvothermal reaction of 5.5'-pyridyl-3.3'-bi-1Hpyrazole with  $CuSO_4 \cdot 5H_2O$  in  $CH_3OH-CH_3CN$ . The initial coordination leads to a  $[3\times3]$  $[Cu_4^I Cu_5^{II}(L)_6]$  $(SO_4) \cdot 2CH_3OH$  (Cu<sub>9</sub>) grid followed by oligomerisation to a  $3 \times [3 \times 3]$  $[Cu_6^{l}Cu_{21}^{l}(L)_{18}(CN)_6(SO_4)](SO_4)_2 \cdot 8CH_3CN$  (Cu<sub>27</sub>) grid through bridging sulfate and *in situ* generated cyanide with time and subsequent supramolecular organization. The isolated silvery lustrous crystals, Cus and Cu<sub>27</sub>, are good electrical conductors, with conductivities of  $3.2 \times 10^{-2}$  and  $5.4 \times 10^{-3}$  S cm<sup>-1</sup>, respectively, due to the mixed valency. Their magnetic properties are dominated by antiferromagnetic interactions. DFT and band structure calculations confirm their valence distribution and semiconducting behavior. The planarity of the large  $3 \times [3 \times 3]$  Cu<sub>27</sub> and its electro-activity renders it suitable for the development of large surface area molecular devices.

> anions.<sup>14</sup> While the former monovalent compound is an insulator, those with mixed Cu<sup>I</sup>/Cu<sup>II</sup> are semiconductors. Other crystalline variants were obtained by tuning the solvents and the time of the solvothermal treatment. Using MALDI-TOF of the solids at different periods of the reaction, we have identified a hierarchical tandem assembly starting from the coordination of the ingredients to form  $[3\times3]$   $[Cu_4^I Cu_5^{II}(L)_6](SO_4) \cdot 2CH_3$ -OH (Cu<sub>9</sub>) followed by its oligomerisation to a giant  $3 \times [3 \times 3]$  $[Cu_{6}^{I}Cu_{21}^{II}(L)_{18}(\mu_{2}-CN)_{6}(\mu_{3}-SO_{4})](SO_{4})_{2} \cdot 8CH_{3}CN$  (Cu<sub>27</sub>) disk and finally its supramolecular arrangement into a crystalline solid (Fig. 1). The two solids Cu<sub>9</sub> and Cu<sub>27</sub> were isolated from pure methanol and a mixture of methanol and acetonitrile, respectively.

> Herein, we report their crystal structures (Tables 1, S1, and S2<sup>†</sup>) and characterization using powder diffraction (Fig. S1<sup>†</sup>), infrared and photoelectron spectroscopies and

Fig. 1 The three-step tandem hierarchy in obtaining Cu<sub>27</sub> from Cu (green), 5,5'-pyridyl-3,3'-bi-1H-pyrazole (rust and blue) and sulphate (red and yellow) and cyanide (cyan) bridges via Cua

# Coordination Oligomerization Supramolecu



View Article Online

View Journal | View Issue

Cu2

<sup>&</sup>quot;State Key Laboratory of Coordination Chemistry, School of Chemistry and Chemical Engineering, Collaborative Innovation Center of Advanced Microstructures, Nanjing University, Nanjing, 210023, P. R. China. E-mail: zuojl@nju.edu.cn

<sup>&</sup>lt;sup>b</sup>Institut de Chimie de Strasbourg, Université de Strasbourg, CNRS-UMR 7177, 4 rue Blaise Pascal, 67008 Strasbourg, France

College of Chemical Engineering, Zhejiang University of Technology, Hangzhou, 310032, P. R. China

<sup>†</sup> Electronic supplementary information (ESI) available. CCDC 1844367 and 1844368. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8sc02851f

thermogravimetry (Fig. S2<sup>†</sup>) together with an experimental study of their electrical, magnetic and optical properties. DFT and band structure calculations were performed to understand the valence distribution and the unusual electrical conductivity. Considerable differences in the electrical and magnetic properties were observed which depend on the average oxidation state of the copper ions. The results are quite exceptional for this series of materials where the central metal moieties provide variable carriers, where their electronic densities and magnetic moments increase with the number of metals and their valences, while the organic ligands interact with those of neighbouring units creating bands favouring electron-transfer.<sup>13</sup> This particular distribution of charges and spins is quite unique in rendering these solids electrically conducting.

### Results and discussion

#### Synthesis

The planar multidentate pyridine–pyrazole ligand H<sub>2</sub>L was synthesized by slight modification of a previously reported procedure.<sup>14-16</sup> Solvothermal reaction of the ligand, neutralized with triethylamine, with cupric sulphate in methanol leads to silvery hexagonal crystals of the nonanuclear  $[Cu_4^{I}Cu_5^{II}(L)_6]$ - $(SO_4) \cdot 2CH_3OH$  (Cu<sub>9</sub>), whereas that in a mixture of methanol and acetonitrile leads to silvery square blocks of the heptaco-sanuclear  $[Cu_{21}^{I}(L)_{18}(\mu_2$ -CN)<sub>6</sub>( $\mu_3$ -SO<sub>4</sub>)](SO<sub>4</sub>)<sub>2</sub> · 8CH<sub>3</sub>CN (Cu<sub>27</sub>).

#### **Crystal structures**

The structure of  $Cu_9$  has a planar nonanuclear [3×3] grid core of copper atoms enveloped by six organic ligands, three above the metal plane and three below. The ligands are almost perpendicular to the metal plane and three on one face are parallel to each other but the other three are almost orthogonal to those on the opposite face. There are slight differences in ligand and metal coordination depending on the positions of the copper atoms and their valences while the crystal symmetry of the cluster is satisfied. The grid is highly deviated from a square in order to accommodate the chelating distances of the coordinating nitrogen pairs and the flattened tetrahedron of  $Cu^{II}$ .  $Cu_{9}$ crystallizes in the monoclinic C2/c space group and the molecular unit includes nine Cu atoms (Cu<sub>4</sub><sup>I</sup>Cu<sub>5</sub><sup>II</sup>), six L<sup>2-</sup> ligands, one  $SO_4^{2-}$  anion and two free methanol molecules. The nine metal ions can be categorized into three types: the four corner ones (Cu5, Cu5', Cu4 and Cu6) each have four nitrogen atoms from two pyridines and two pyrazoles, while the four in the middle of the edges (Cu1, Cu3, Cu3', and Cu1') have four nitrogen atoms from one pyridine and three pyrazoles, and the central one (Cu2) has four nitrogen atoms from four pyrazoles (Fig. 1). All the copper ions have a [CuN<sub>4</sub>] coordination sphere and are connected by N…N bridges. The average Cu-N bond lengths are [Cu1-N] = [Cu2-N] = [Cu3-N] = 1.96, [Cu4-N] = [Cu6-N] = 1.99and [Cu5-N] = 2.02 Å. They indicate that Cu4, Cu5, Cu5' and Cu6 are Cu(I), while Cu1, Cu2, Cu3, Cu1' and Cu3' are Cu(II).

The ligands on each face stand in an offset parallel arrangement. The intramolecular  $\pi \cdots \pi$  distances are rather short (3.3–3.5 Å) (Fig. S3<sup>†</sup>), indicating the presence of strong

 $\pi \cdots \pi$  interactions. The ligands between adjacent molecules are stacked through offset  $\pi \cdots \pi$  bond-over-ring mode. The molecules form staircase-like one-dimensional (1D) arrays running along the *b*-axis with a face-to-face distance of 3.32 Å, suggesting strong interaction. Intermolecular H-bonds (Table S2†) between the 1D-arrays extend them into 2D layers in the *ac*-plane (Fig. S4 and S5 and Table S3†).

 $Cu_{27}$  is an oligomer of three  $Cu_9$  with a  $3 \times [3 \times 3]$  heptacosanuclear structure with a distorted grid core of copper atoms enveloped by eighteen organic ligands and combined with one sulphate and six cyanide bridges. It has a pseudo-trigonal symmetry of three Cu<sub>9</sub> held together by a SO<sub>4</sub> sitting centrally on the C<sub>3</sub>-axis and six bridging cyanide ions between the edges (Fig. S3<sup>†</sup>). The twenty-seven metal ions can be categorized into four types: the three corner ones (Cu9, Cu18, and Cu27) have four nitrogen atoms from two pyridines and two pyrazoles, and the average Cu-N bond lengths are [Cu9-N] = 2.03, [Cu18-N] =2.01, and [Cu27-N] = 2.04 Å. The angles are  $\angle N18-Cu9-N36 =$  $120.2^{\circ}$  ( $\angle N54$ -Cu18-N72 =  $128.6^{\circ}$ ,  $\angle N90$ -Cu27-N108 = 122.7°) and  $\angle N17$ -Cu9-N35 = 157.0° ( $\angle N53$ -Cu18-N71 = 161.4°,  $\angle$  N89–Cu27–N107 = 157.3°). Their geometries are closer to a tetrahedron; thus, they are assigned to cuprous ions. The six atoms (Cu6, Cu15, Cu24, Cu8, Cu17, and Cu26) in the middle of the edges are four-coordinated ones with one pyridine and three pyrazoles and the average Cu-N bond lengths are [Cu6-N] = [Cu24-N] = [Cu8-N] = [Cu17-N] = [Cu26-N] = 1.97and [Cu15-N] = 1.95 Å. The angles are  $\angle N12-Cu6-N34 =$  $132.37^{\circ}$  ( $\angle N47$ -Cu15-N69 =  $128.1^{\circ}$ ,  $\angle N84$ -Cu24-N106 = 129.5°),  $\angle$  N11-Cu6-N33 = 170.5° ( $\angle$ N48-Cu15-N70 = 170.7°,  $\angle$  N84-Cu24-N106 = 170.3°),  $\angle$  N16-Cu8-N30 = 130.6° ( $\angle$  N52-Cu17-N66 =  $132.7^{\circ}$ ,  $\angle N88$ -Cu26-N102 =  $135.1^{\circ}$ ), and  $\angle N15$ - $Cu8-N29 = 170.5^{\circ} (\angle N51-Cu17-N65 = 170.3^{\circ}, \angle N87-Cu26 N101 = 169.0^{\circ}$ ). Their geometries are more planar; thus, they are assigned to cupric ions. The central ones (Cu5, Cu14, and Cu23) have four nitrogen atoms from four pyrazoles. The average Cu-N bond lengths are [Cu5-N] = 2.03, [Cu14-N] = 2.01, and [Cu23-N] = 2.04 Å. The angles are  $\angle N18$ -Cu9-N36  $= 120.2^{\circ}$  ( $\angle N54$ -Cu18- $N72 = 128.6^{\circ}, \angle N90-Cu27-N108 = 122.7^{\circ}$  and  $\angle N17-Cu9-N35$  $= 157.0^{\circ} (\angle N53$ -Cu18-N71  $= 161.4^{\circ}, \angle N89$ -Cu27-N107 =157.3°). Their geometries are close to a tetrahedron, and thus they are assigned to cuprous ions (Fig. 1).

The intramolecular  $\pi \cdots \pi$  distances remain short in the range 3.3–3.4 Å and C–H $\cdots \pi$  distances of 2.60–2.89 Å are short and lie in the *bc*-plane. The adjacent molecules are stacked through the strong, offset  $\pi \cdots \pi$  interaction with a face-to-face distance of 3.35 Å to furnish staircase-like one-dimensional (1D) arrays running along the *b*-axis (Fig. S4†). Intermolecular C–H $\cdots \pi$  between the 1D-arrays extend them into 2D layers in the *bc*-plane (Fig. S5 and S6 and Table S3†). These 2D layers form 3D supramolecular frameworks by using acetonitrile molecules and sulphate anions.

#### MALDI-TOF

While  $Cu_9$  was isolated as crystals in pure methanol, the presence of acetonitrile was found to be indispensable for the crystallization of  $Cu_{27}$ . It was therefore necessary to understand

#### **Chemical Science**

the relation between the two compounds. We therefore studied the MALDI-TOF MS of the solids formed from a mixture of methanol and acetonitrile as a function of time (Fig. 2).

For the solids, by 30 minutes only peaks assignable to Cu<sub>9</sub> were located for the black product. Their intensities increase up to 2 hours when the peaks for Cu<sub>27</sub> start to appear. The quantity of the Cu<sub>27</sub> then increases at the expense of Cu<sub>9</sub> with time. Even after 24 hours there is still a trace of Cu<sub>9</sub> but the bulk comprises crystals suitable for crystallography. It is therefore clear from a mechanistic point of view that the initial step is the coordination of the ligand to give the  $[3 \times 3]$  Cu<sub>9</sub> grid cluster and with time the Cu<sub>9</sub> units are oligomerized to the  $3 \times [3 \times 3]$  Cu<sub>27</sub> cluster with the help of bridging SO<sub>4</sub> and *in situ* generated cyanide ions. The final step is the crystallization using the supramolecular interactions. It is worth noting that crystals of  $Cu_{27}$  are insoluble in organic solvents compared to those of Cu<sub>9</sub>. Interestingly, Cu<sub>9</sub> is obtained starting from different copper salts and different solvents but  $Cu_{27}$  is obtained only in the presence of SO<sub>4</sub> and acetonitrile. The three step hierarchical process is new in the chemistry of oligomerisation of clusters.

# X-ray photoelectron spectroscopy and spin density calculations

The valence states of the copper ions and their proportions in the compounds were determined from X-ray photoelectron spectroscopy. The  $2p_{1/2}$  and  $2p_{3/2}$  peaks are characterized by doublets for both **Cu**<sub>9</sub> and **Cu**<sub>27</sub>. The energies of the  $2p_{3/2}$  doublet are 934.8 and 931.9 eV (Fig. 3), corresponding to Cu<sup>II</sup> and Cu<sup>I</sup>, respectively.<sup>17</sup> A similar observation was made for the  $2p_{1/2}$  peaks. Analysis of the areas of these two peaks gave Cu<sup>I</sup> : Cu<sup>II</sup> ratios of 43 : 57 and 23 : 77 respectively, consistent with the calculated balance of charge of the formula and crystallographic findings of  $4Cu^{I} : 5Cu^{II}$  and  $2Cu^{I} : 7Cu^{II}$  for **Cu**<sub>9</sub> and **Cu**<sub>27</sub>. It is noted that the complete structural characterization from single crystals of the



Fig. 2 MALDI-TOF mass spectra of the solid products as a function of time of solvothermal treatment.



Fig. 3 Cu- $2p_{3/2}$  valence band XPS spectra of (a) Cu<sub>9</sub> and (b) Cu<sub>27</sub>. Spin density distributions (green is positive and blue is negative) for (c) Cu<sub>9</sub> and (d) Cu<sub>27</sub>. The black labels are for Cu(II) and the pink labels are for Cu(II).

mixed valence of a common molecular brick is quite rare. The presence of different valences is very interesting in view of charge delocalization and thus band-filling. They are consequently reflected in the conductivity, which is also important in magnetic properties because of the similar structure with different numbers of spin carriers. In contrast to crystallographic data, XPS gives a more accurate definition of the valence states of the compounds and confirms that they belong to Class II of the mixed-valence classification.<sup>18</sup>

Spin density calculations were performed for the two compounds built from the experimentally determined geometries with H atom optimization (Fig. 3 and Table S4†).<sup>19</sup> We find that they are centred principally on the Cu atoms with some delocalization on the coordinated nitrogen atoms. Very little spin density is diffused onto the carbon atoms of the organic ligands. The oxidation states were evaluated by the localized orbital bonding analysis (LOBA) method within Mutliwfn and the results are presented in Table S4.<sup>†20</sup> The results indicated that Cu4, Cu5, Cu5' and Cu6 are Cu(1), and the remaining copper ions are Cu(1) in Cu<sub>9</sub> and Cu5, Cu9, Cu14, Cu18, Cu23 and Cu27 are Cu(1), while all the other copper ions are Cu( $\pi$ ) in Cu<sub>27</sub>.

#### **Electrical conductivity**

The high absorption and silvery lustre of the crystals are consistent with the mixed-valence found and therefore, they are

expected to be conducting (Fig. S7<sup>†</sup>). Gold wires were attached on the surface using silver paste for two-probe resistance and I-V measurements at different temperatures. Measurements on single crystals of Cu<sub>9</sub> found a room temperature conductivity of  $3.2 \times 10^{-2} \ {
m S \ cm^{-1}}$  along the *ab*-plane (almost parallel to the plane of the molecules) and  $4.3 \times 10^{-8}$  S cm<sup>-1</sup> perpendicular to it. The temperature dependence of the resistivity in the *ab*-plane follows an Arrhenius behaviour with a fairly low activation energy of 0.02 eV. Measurements for single crystals of Cu<sub>27</sub> found a slightly lower room temperature conductivity of  $10^{-3}$  S cm<sup>-1</sup> along the *bc*-plane of the molecules but a higher one of  $10^{-5}$  S cm<sup>-1</sup> perpendicular to it (Fig. 4). The crystals of  $Cu_{27}$  lose the solvents with time and the temperature dependence measurement was hampered due to the vacuum needed for the cryostat. Monitoring the crystals under an optical microscope during the solvent evaporation reveals a limited stability and appearance of cleavages after dryness, presumably between the layers (Fig. S8<sup>†</sup>). The observed values of conductivity are quite promising for such molecules where charge becomes mobile within the solid through supramolecular  $\pi \cdots \pi$ interactions as is the low activation energy in the case of Cu<sub>9</sub>.

#### Magnetic properties and electron paramagnetic resonance

The temperature dependences of the magnetic moments, represented by the product of susceptibility and temperature, for the two compounds are shown in Fig. 5. We caution that loss of solvent for  $Cu_{27}$  may have taken place but the molecular structure and valence state are not altered. The values of  $\chi_M T$  at 300 K for the polycrystalline samples are 1.88 and 6.94 cm<sup>3</sup> mol<sup>-1</sup> K for Cu<sub>9</sub> and Cu<sub>27</sub>, respectively. Curie–Weiss fits for data above 150 K gave respectively C = 2.26 and 9.53 cm<sup>3</sup> mol<sup>-1</sup> K and  $\theta = -70$  and -110 K. From the Curie constants the average *g*-value is 2.19  $\pm$  0.01, which is reasonable for Cu<sup>2+</sup>. The EPR spectra of polycrystalline samples are characterized by broad resonances ( $\Delta H_{pp} = 172$  G and g = 2.06 (Cu<sub>9</sub>) and 580 G



Fig. 4 Arrhenius plot of In(resistance) *versus* scaled inverse temperature (red circles represent experimental data and the black line is the linear fit). Inset: calculated band structure and Brillouin zone.



Fig. 5 The temperature dependence of  $\chi_M T$  for Cu<sub>9</sub> (red circles) and Cu<sub>27</sub> (blue circles), measured in a field of 1.0 kOe. The solid black lines are the best fits (see text).

and g = 2.10 (Cu<sub>27</sub>)) (Fig. S9<sup>†</sup>). The large negative Weiss constants indicate strong antiferromagnetic coupling between the nearest neighbouring Cu(II) ions. The temperature dependences of the susceptibility were modelled using the Hamiltonian given in equations (ESI<sup>†</sup>) giving, for Cu<sub>9</sub>, g = 2.15,  $J_1 =$  $-28.3 \text{ cm}^{-1}$  and  $J_2 = -27.8 \text{ cm}^{-1}$ .<sup>21</sup> For Cu<sub>27</sub>, due to the large number of Cu(II) and the limitation of computer ability, it is impossible to apply the irreducible tensor operator method used in PHI. By modifying the LOOP code<sup>22</sup> of the Algorithms and Libraries for Physical Simulations (ALPS) project,23 we herein coupled the Genetic Algorithm with the Quantum Monte Carlo (GA-OMC) method to search for the global optimal parameters.<sup>24</sup> Essentially, the entire spin exchange network per cluster (Fig. 5) was simplified to a 3 J model (see the Hamiltonian operator in the ESI<sup>†</sup>). In view of the large number of atoms per cluster and the considerable  $\pi \cdots \pi$  supramolecular interaction, a temperature-independent paramagnetism (TIP) parameter and intercluster interaction (zJ) were respectively considered. After 200 iterations, the simulation yielded one set of optimal parameters:  $J_1 = -4.9 \text{ cm}^{-1}$ ,  $J_2 = -210.7 \text{ cm}^{-1}$ ,  $J_3 =$  $-3.8 \text{ cm}^{-1}$ , g = 2.19, TIP = 0.00 cm<sup>3</sup> mol<sup>-1</sup>,  $zJ = 0.01 \text{ cm}^{-1}$ , and  $R = 4.25 \times 10^{-5} (R = \sum [(\chi_{\rm M} T)_{\rm obs} - (\chi_{\rm M} T)_{\rm calcd}]^2 / \sum [(\chi_{\rm M} T)_{\rm obs}]^2).$ The values of the moments are direct electron counts and give support to the XPS and crystallography results while the excellent fits suggest that a localized moment picture is an adequate representation. Interestingly, the exchange through the  $C \equiv N$ bridge is the strongest compared to those through the N=N bridge. The reduction in the values of the exchange interactions within the Cu<sub>9</sub> segments of Cu<sub>27</sub> may be due to the effect of the strong field of the bridging cyanide groups.

#### **Optical properties**

The optical properties of the compounds in the limiting region of 200–2000 nm were studied by diffuse reflectance

|   | Cu <sub>9</sub> | Cu <sub>27</sub>      |
|---|-----------------|-----------------------|
| Formula                                 | C98H68Cu9N36O6S | C310H204Cu27N122O12S3 |
| Fw                                      | 2449.80         | 7641.69               |
| Space group                             | C2/c            | $P2_1/n$              |
| a/Å                                     | 32.700(18)      | 14.7327(3)            |
| b/Å                                     | 10.890(6)       | 47.3434(8)            |
| c/Å                                     | 30.813(17)      | 51.2702(8)            |
| $\beta/^{\circ}$                        | 114.970(7)      | 97.581(2)             |
| $V/\text{\AA}^3$                        | 9947(9)         | 35 448.2(11)          |
| Ζ                                       | 4               | 4                     |
| $D_{\rm c}/{\rm g~cm^{-3}}$             | 1.630           | 1.432                 |
| T/K                                     | 150(2)          | 150(2)                |
| F(000)                                  | 4900            | 14 292                |
| R <sub>int</sub>                        | 0.059           | 0.065                 |
| $R_1^a \left[ I \ge 2\sigma(I) \right]$ | 0.070           | 0.066                 |
| $w R_2^{b}$ (all data)                  | 0.214           | 0.141                 |
| GoF                                     | 1.00            | 1.02                  |

measurements on the powdered samples in  $BaSO_4$ . The spectra, shown in Fig. S10,<sup>†</sup> present an edge in each case where a linear extrapolation was used to estimate the band gap. Surprisingly, considering the crude approximated extrapolation, there is a general trend in the spectra. The band-gaps were 0.03 and 0.25 eV for Cu<sub>9</sub> and Cu<sub>27</sub>, respectively, which are close to those found by conductivity measurements and band structure calculations.

### Conclusions

A hierarchical tandem assembly, including coordination to nonanuclear  $[3 \times 3]$  Cu<sub>9</sub>, its oligometrisation to a giant  $3 \times [3 \times 3]$ Cu<sub>27</sub> cluster and finally the supramolecular organization into crystals, is achieved. The second step was revealed by MALDI-TOF and confirmed by single crystal diffraction. The oligomerisation step is facilitated by the presence of sulphate and in situ generated cyanide from acetonitrile. The oligomerisation is not observed in their absence. Their mixed-valence states were confirmed by X-ray photoelectron spectroscopy and the metallic lustre of the crystals is compatible with the fairly high electrical conductivities with three orders of magnitude anisotropy. The structure-electrical property relationship suggests that the  $\pi \cdots \pi$  interaction between neighbouring molecules is the ingredient for the electron-transfer resulting in high conductivity and low activation energy. However, the magnetic susceptibility fittings to localized moment models are a clear indication that these two compounds belong to Class II of the mixed-valence classification. Furthermore, the experimental results are in good agreement with those estimated from DFT and band-structure calculations. Such large planar electroactive clusters which are highly absorbing in the visible region should be suitable for large surface area devices for harvesting light for energy purposes.

### Experimental

#### Materials

All chemicals and solvents were obtained from commercial sources and utilized without further purification.

[Cu<sup>I</sup><sub>4</sub>Cu<sup>T</sup><sub>5</sub>(L)<sub>6</sub>](SO<sub>4</sub>) · 2CH<sub>3</sub>OH (Cu<sub>9</sub>). A CH<sub>3</sub>OH (8 mL, AR-grade  $\geq$  99.7%) solution of CuSO<sub>4</sub> · 5H<sub>2</sub>O (28.0 mg, 0.112 mmol) and H<sub>2</sub>L (21.5 mg, 0.075 mmol) was adjusted to pH ≈ 8 with Et<sub>3</sub>N in a 15 mL Pyrex vial which was then tightly capped and heated at 140 °C for 24 h. After the autoclave was cooled to 30 °C at a rate of 5 °C h<sup>-1</sup>, the black needle shaped crystals of Cu<sub>9</sub> were collected within one hour, washed with methanol, and dried in air (yield: 15.2 mg, 46% based on the ligand). ESI-MS (positive mode, CH<sub>3</sub>OH): *m/z*: 1144.97 (C<sub>96</sub>H<sub>60</sub>N<sub>36</sub>Cu<sub>9</sub>)<sup>2+</sup>. Elemental analysis (%) calculated for C<sub>98</sub>H<sub>68</sub>N<sub>36</sub>O<sub>6</sub>SCu<sub>9</sub>: C, 48.05; H, 2.80; N, 20.58; found: C, 48.56; H, 2.98; N, 20.22. Infrared (KBr, cm<sup>-1</sup>): 3412(m), 3098(m), 1616(s), 1588(w), 1532(m), 1456(s), 1324(m), 1166(s), 1128(s), 1052(m), 986(m), 788(s), 612(m).

 $[Cu_{6}^{I}Cu_{21}^{II}(L)_{18}(CN)_{6}(SO_{4})](SO_{4})_{2} \cdot 8CH_{3}CN$  (Cu<sub>27</sub>). A suspension of  $CuSO_4 \cdot 5H_2O$  (28.0 mg, 0.112 mmol) and  $H_2L$  (21.5 mg, 0.075 mmol) in a mixture of 4 mL CH<sub>3</sub>OH and 4 mL CH<sub>3</sub>CN was adjusted to pH  $\approx$  8 with Et<sub>3</sub>N, filtered, and divided into two equal parts. The first part was slowly concentrated in air to give a black powder of a Cu<sub>9</sub> complex which was only identified by ESI-MS. The second part was placed in a 15 mL Pyrex vial which was tightly capped and heated at 140 °C for 24 h. After the autoclave was cooled at a rate of 5  $^{\circ}$ C h<sup>-1</sup> to 30  $^{\circ}$ C, black square block shaped crystals of Cu<sub>27</sub> were collected, washed with ethanol, and dried in air (yield: 32.0 mg, 53% based on the (positive mode): m/z: 7313.25 ligand). MALDI-TOF  $\{ [Cu_6^I Cu_{21}^{II}(L)_{18}(CN)_6(SO_4)] (SO_4)_2 \}^+$ . Elemental analysis (%) calculated for C310H204Cu27N122O12S3: C, 48.72; H, 2.69; N, 22.36; found: C, 48.53; H, 2.51; N, 22.65. Infrared (KBr, cm<sup>-1</sup>): 2174(w), 2104(w), 1604(s), 1562(m), 1524(w), 1450(s), 1436(s), 1348(s), 1242(w), 1154(m), 1108(w), 1048(m), 988(m), 774(s), 714(m).

#### Measurements

The C, H and N microanalyses were carried out with a Vario Micro Cube elemental analyzer. Thermogravimetric analyses (TGA) were performed on a NETZSCH STA 449 F3 Jupiter TGA instrument in flowing N2 at a heating rate of 5 °C per minute in the range of 40 to 800 °C. Samples for infrared (IR) were prepared as KBr pellets, and spectra were recorded in the 400- $4000 \text{ cm}^{-1}$  range using a Vector 22 Bruker spectrophotometer. UV-Vis-NIR spectra were recorded in diffuse reflectance mode on a UV-3600 Shimadzu spectrometer. Powder X-ray diffraction (PXRD) patterns were recorded at 293 K on a D8 ADVANCE (Cu Ka1,  $\lambda = 1.54056$  Å; Ka2,  $\lambda = 1.54439$  Å) powered at 40 kV and 40 mA. The powder samples were prepared by crushing the crystals and the PXRD scanned from 5 to 50° at a rate of  $5^{\circ}$  min<sup>-1</sup>. Calculated PXRD patterns were generated using Mercury 3.9.25 Mass spectra were recorded with an LCO Fleet instrument for ESI-MS and an ultrafleXtreme TOF/TOF for MALDI-TOF/TOF mass spectrometry. Electron paramagnetic

resonance (EPR) spectra were recorded on a Bruker ER-420 spectrometer with a 100 kHz magnetic field in the X band at room temperature (RT). X-ray photoelectron spectroscopy (XPS) studies were performed using a Thermo Fisher ESCALAB 250 X-ray photoelectron spectrometer (powered at 150 W) using Al K\alpha radiation ( $\lambda = 8.357$  Å). Magnetisation measurements for polycrystalline samples were performed on a Quantum Design MPMS-SQUID-VSM magnetometer in the temperature range from 1.8 to 300 K for direct current (dc) applied fields ranging from 0 to 70 kOe. Diamagnetic corrections were calculated using Pascal's constants,26 and an experimental correction for the diamagnetic sample holder was applied. Temperaturedependent electrical conductivities and I-V curves were measured using a Keithley 4200 semiconductor parameter analyser using single crystal samples by the two-probe method using silver paste and gold wire contacts.

#### X-ray structure determination

The single-crystal X-ray diffraction data for Cu<sub>9</sub> and Cu<sub>27</sub> were collected on a Bruker D8 VENTURE diffractometer with a PHOTON 100 CMOS detector equipped with a Mo-target X-ray tube ( $\lambda = 0.71073$  Å) at T = 150 K by using the  $\varphi$ - $\omega$  scan technique. Data were corrected for absorption using the empirical method SADABS.27 The structures were solved using SHELXT28 and refined using the full-matrix least-squares procedures within the SHELXTL software package.29 All non-hydrogen atoms were refined with anisotropic displacement parameters. The hydrogen atoms were generated geometrically (C-H = 0.95 Å) using the riding model. For all structures, the guest molecules were further confirmed by thermogravimetric analyses (TGA). It was not possible to resolve diffuse electron density residuals (enclosed solvent molecules). The co-crystallized solvent molecules were squeezed with the SQUEEZE facility from PLATON:<sup>30</sup> 8CH<sub>3</sub>CN for Cu<sub>27</sub> (per asymmetric unit). Crystallographic information for all the complexes is presented in Table 1. Selected bond distances and angles are given in the ESI.<sup>†</sup> The crystallographic information files have been deposited at The Cambridge Crystallographic Data Centre with references 1844367 (Cu<sub>9</sub>) and 1844368 (Cu<sub>27</sub>); they can be obtained from http//:www.ccdc.cam.ac.uk/data\_request/cif.

#### DFT calculations

All theoretical calculations were carried out at the DFT level of theory using the hybrid B3LYP/6-31g\* exchange–correlation functional<sup>31–33</sup> as implemented in the Gaussian 09 program.<sup>34</sup> A quadratic convergence method was employed in the SCF process.<sup>35</sup> The triple- $\zeta$  quality basis set proposed by Ahlrichs and coworkers has been used for all atoms.<sup>36</sup> Calculations were performed on the complexes built from the experimentally determined geometries with H atom optimization.

#### Band structure calculations

To gain a good insight into the electrical and optical properties, we performed first-principles electronic structure calculations for  $Cu_9$  based on density functional theory with generalized gradient approximation (GGA) in the form of the PerdewBurke–Ernzerhof (PBE) functional for exchange–correlation potential.<sup>37</sup> We used the Cambridge Serial Total Energy Package (CASTEP),<sup>38</sup> which is a self-consistent-field method (tolerance  $5.0 \times 10^{-5}$  eV per atom) in conjunction with plane-wave basis sets with a cut-off energy of 517 eV using OTFG ultrasoft pseudopotentials in reciprocal space.<sup>39</sup> Calculations were performed starting with the crystal structure data of **Cu**<sub>9</sub> at 150(2) K without optimization. The first Brillouin zone was sampled with a grid spacing of 0.005 Å<sup>-1</sup> for wave functions and 0.05 Å<sup>-1</sup> for the density of states (DOS) on the basis of the Monkhorst–Pack special *k*-point scheme.<sup>40</sup>

### Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

This work was supported by the National Basic Research Program of China (2018YFA0306004) and the National Natural Science Foundation of China (No. 21631006). M. K. was funded by the Centre National de la Recherche Scientifique (CNRS), France. We thank the staff of the BL17B beamline of the National Facility for Protein Science Shanghai (NFPS) at the Shanghai Synchrotron Radiation Facility (SSRF), for assistance with X-ray data collection and Dr Zhenyi Zhang of Bruker (Beijing) Scientific Technology Co., Ltd., for help with structure refinement.

### Notes and references

- 1 J. C. Love, L. A. Estroff, J. K. Kriebel, R. G. Nuzzo and G. M. Whitesides, *Chem. Rev.*, 2005, **105**, 1103–1169.
- 2 W. A. Lopes and H. M. Jaeger, Nature, 2001, 414, 735-738.
- 3 Y. He, T. Ye, M. Su, C. Zhang, A. E. Ribbe, W. Jiang and C. Mao, *Nature*, 2008, **452**, 198–201.
- 4 J.-F. Ayme and J.-M. Lehn, Adv. Inorg. Chem., 2018, 71, 3-78.
- 5 M. Barboiu, A.-M. Stadler and J.-M. Lehn, *Angew. Chem., Int. Ed.*, 2016, 55, 4130–4154.
- 6 Y. Kim, M. F. Mayer and S. C. Zimmerman, *Angew. Chem., Int. Ed.*, 2003, **42**, 1121–1126.
- 7 P. Teng, Z. Niu, F. She, M. Zhou, P. Sang, G. M. Gray,
  G. Verma, L. Wojtas, A. van der Vaart, S. Ma and J. Cai, *J. Am. Chem. Soc.*, 2018, 140, 5661–5665.
- 8 A. Coskun, M. Banaszak, R. D. Astumian, J. F. Stoddart and B. A. Grzybowski, *Chem. Soc. Rev.*, 2012, **41**, 19–30.
- 9 L. S. Xie, L. Sun, R. Wan, S. S. Park, J. A. DeGayner, C. H. Hendon and M. Dincă, *J. Am. Chem. Soc.*, 2018, 140, 7411-7414.
- 10 L. Sun, C. H. Hendon, S. S. Park, Y. Tulchinsky, R. Wan, F. Wang, A. Walsh and M. Dincă, *Chem. Sci.*, 2017, 8, 4550–4557.
- 11 J. V. Barth, G. Costantini and K. Kern, *Nature*, 2005, **437**, 671–679.
- 12 Y.-K. Deng, H.-F. Su, J.-H. Xu, W.-G. Wang, M. Kurmoo, S.-C. Lin, Y.-Z. Tan, J. Jia, D. Sun and L.-S. Zheng, *J. Am. Chem. Soc.*, 2016, **138**, 1328–1334.

- 13 L.-Y. Guo, H.-F. Su, M. Kurmoo, C.-H. Tung, D. Sun and L.-S. Zheng, J. Am. Chem. Soc., 2017, 139, 14033–14036.
- 14 F. Yu, J. Li, Z.-H. Cao, M. Kurmoo and J.-L. Zuo, *Inorg. Chem.*, 2018, 57, 3443–3450.
- 15 J. Janculev and B. Podolesov, God. Zb., Filoz. Fak. Univ., Skopje, Prir.-Mat. Oddel, 1958, 11, 47-49.
- 16 S. Terashima, G. N. Newton, T. Shiga and H. Oshio, *Inorg. Chem. Front.*, 2015, 2, 125–128.
- 17 Z.-L. Wu, C.-H. Wang, B. Zhao, J. Dong, F. Lu, W.-H. Wang, W.-C. Wang, G.-J. Wu, J.-Z. Cui and P. Cheng, *Angew. Chem., Int. Ed.*, 2016, 55, 4938–4942.
- 18 M. B. Robin and P. Day, Adv. Inorg. Chem. Radiochem., 1968, 10, 247–422.
- 19 S. H. Vosko, L. Wilk and M. Nusair, *Can. J. Phys.*, 1980, 58, 1200–1211.
- 20 T. Lu and F. Chen, J. Comput. Chem., 2012, 33, 580-592.
- 21 N. F. Chilton, R. P. Anderson, L. D. Turner, A. Soncini and K. S. Murray, *J. Comput. Chem.*, 2013, 34, 1164–1175.
- 22 A. W. Sandvik, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1999, **59**, R14157–R14160.
- 23 A. F. Albuquerque, F. Alet, P. Corboz, P. Dayal, A. Feiguin, S. Fuchs, L. Gamper, E. Gull, S. Gürtler, A. Honecker, R. Igarashi, M. Körner, A. Kozhevnikov, A. Läuchli, S. R. Manmana, M. Matsumoto, I. P. McCulloch, F. Michel, R. M. Noack, G. Pawłowski, L. Pollet, T. Pruschke, U. Schollwöck, S. Todo, S. Trebst, M. Troyer, P. Werner and S. Wessel, *J. Magn. Magn. Mater.*, 2007, **310**, 1187–1193.
- 24 S. Forrest, Science, 1993, 261, 872-878.
- 25 Mercury CSD 3.9 (Build RC1), CCDC, 2016.
- 26 G. A. Bain and J. F. Berry, J. Chem. Educ., 2008, 85, 532–536.
  27 G. M. Sheldrick, SADABS—Bruker AXS Area Detector Scaling and Absorption, 2008.
- 28 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Adv.*, 2015, A71, 3–8.
- 29 G. M. Sheldrick, *Acta Crystallogr., Sect. C: Struct. Chem.*, 2015, C71, 3–8.
- 30 P. van der Sluis and A. L. Spek, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 1990, **46**, 194–201.

- 31 A. D. Becke, *Phys. Rev. A: At., Mol., Opt. Phys.*, 1988, **38**, 3098–3100.
- 32 C. T. Lee, W. T. Yang and R. G. Parr, Phys. Rev. B: Condens. Matter Mater. Phys., 1988, 37, 785–789.
- 33 A. D. Becke, J. Chem. Phys., 1993, 98, 5648-5652.
- 34 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr, E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, I. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, Gaussian 09, rev. B.01, Gaussian, Inc., Wallingford CT, 2009.
- 35 G. B. Bacskay, Chem. Phys., 1981, 61, 385-404.
- 36 A. Schäfer, C. Huber and R. Ahlrichs, J. Chem. Phys., 1994, 100, 5829–5835.
- 37 J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, 77, 3865–3868.
- 38 S. J. Clark, M. D. Segall, C. J. Pickard, P. J. Hasnip, M. I. J. Probert, K. Refson and M. C. Payne, *Z. Kristallogr.*, 2005, **220**, 567–570.
- 39 Z. Fei, Y. Shi, L. Pu, F. Gao, Y. Liu, L. Sheng, B. G. Wang, R. Zhang and Y. D. Zheng, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2008, 78, 201402.
- 40 H. J. Monkhorst and J. D. Pack, *Phys. Rev. B: Solid State*, 1976, 13, 5188–5192.