CORRECTION

View Article Online
View Journal | View Issue

Cite this: *Chem. Sci.*, 2018, 9, 2010

Correction: Correction: Hypoxia inducible factor (HIF) as a model for studying inhibition of protein–protein interactions

George M. Burslem,^{id} ^{ab} Hannah F. Kyle,^{bc} Adam Nelson,^{id} ^{ab}
Thomas A. Edwards^{id} ^{bc} and Andrew J. Wilson^{id} ^{*ab}

DOI: 10.1039/c8sc90024h

www.rsc.org/chemicalscience

Correction for 'Correction: Hypoxia inducible factor (HIF) as a model for studying inhibition of protein–protein interactions' by George M. Burslem *et al.*, *Chem. Sci.*, 2017, 8, 5214–5215.

The authors regret that in their original correction regarding Fig. 3 there remained errors in the structures of EZN-2208, geldanamycin, ganetespib and echinomycin. The correct figure is displayed below.

^aSchool of Chemistry, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK. E-mail: a.j.wilson@leeds.ac.uk

^bAstbury Centre for Structural Molecular Biology, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK

^cSchool of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK

Fig. 3 Modulators of the HIF pathway (a) topotecan and derivative EZN-2208 (b) 2-oxoglutarate (c) DMOG and oxalyl glycine (d) FG-2216 (e) roxadustat (f) geldanamycin (g) ganetespib (h) X-ray crystal structure of a hydroxyproline derived inhibitor (orange) bound to pVHL (cyan), PDB ID 3zrc (i) optimised hydroxyproline derived pVHL inhibitor VH298 (j) echinomycin (k) DNA sequence specific polyamide (l) bortezomib.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.