Chemical Science

CORRECTION

View Article Online
View Journal | View Issue

Cite this: *Chem. Sci.*, 2018, **9**, 3893

Correction: Spectroscopic observation of two-center three-electron bonded (hemi-bonded) structures of $(H_2S)_n^+$ clusters in the gas phase

Dandan Wang and Asuka Fujii *

DOI: 10.1039/c8sc90072h

www.rsc.org/chemicalscience

Correction for 'Spectroscopic observation of two-center three-electron bonded (hemi-bonded) structures of $(H_2S)_n^+$ clusters in the gas phase' by Dandan Wang *et al.*, *Chem. Sci.*, 2017, **8**, 2667–2670.

The authors regret that some important references were omitted from the original article. These references are presented herein.

The experimental observation of the sulfur–sulfur hemi-bond was pioneered by Asmus and coworkers.[1–6] They observed transient absorption due to the σ*–σ electronic transition in solution. Moreover, they observed the transient absorptions of the hemi-bonds of sulfa with a variety of counter atoms as well as those of N–N and I–I hemi-bonds.[7–11] The electronic spectrum of $(H_2S)_2^+$ in aqueous solution was also reported by Asmus, though detailed structural information is difficult to extract from the broadened electronic transition.[3]

The S–S hemi-bond in gas phase molecules was first reported using mass spectrometry.[5] The stable dimer cation formation of bis(isopropyl)sulfide was observed and hemi-bond formation was proposed on the basis of the fragmentation pattern. Gas phase dimerization equilibrium measurement of dimethyl sulfide cations has also suggested formation of the S–S hemi-bond.[12] Very recently, infrared Stark spectroscopy was applied to $Cl–NH_3$ in He droplets,[13] and hemi-bond formation was concluded by the shift of the NH stretch and dipole moment measurements. This result is consistent with the prediction by high level computation of similar systems.[14]

Theoretical calculations of the S–S hemi-bond were first performed by Clark for $(H_2S)_2^+$,[15] and the series of his study has been extended to a variety of hemi-bonded systems.[16–18]

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

References

1 D. Bahnemann and K.-D. Asmus, *J. Chem. Soc., Chem. Commun.*, 1975, 238–239.
2 K.-D. Asmus, *Acc. Chem. Res.*, 1979, **12**, 436–442.
3 S. A. Chaudri and K.-D. Asmus, *Angew. Chem., Int. Ed.*, 1981, **20**, 672–673.
4 M. Göbl, M. Bonifačić and K.-D. Asmus, *J. Am. Chem. Soc.*, 1984, **106**, 5984–5988.
5 T. Drewello, C. B. Lebrilla, H. Schwarz, L. J. de Koning, R. H. Fokkens, N. M. M. Nibbering, E. Anklam and K.-D. Asmus, *J. Chem. Soc., Chem. Commun.*, 1987, 1381–1383.
6 K.-D. Asmus, in *Sulfur-Centered Reactive Intermediates in Chemistry and Biology, NATO-ASI Series, Series A: Life Sciences*, Springer, 1990, pp. 155–172.
7 M. Bonifačić and K.-D. Asmus, *J. Chem. Soc., Perkin Trans. 2*, 1980, 758–762.
8 K.-D. Asmus, M. Göbl, K.-O. Hiller, S. Mahling and J. Mönig, *J. Chem. Soc., Perkin Trans. 2*, 1985, 641–646.
9 H. Hungerbühler, S. N. Guha and K.-D. Asmus, *J. Chem. Soc., Chem. Commun.*, 1991, 999–1001.
10 R. W. Alder, M. Bonifačić and K.-D. Asmus, *J. Chem. Soc., Perkin Trans. 2*, 1986, 277–284.
11 H. Mohan and K.-D. Asmus, *J. Am. Chem. Soc.*, 1987, **109**, 4745–4746.
12 A. J. Illies, P. Livant and M. L. McKee, *J. Am. Chem. Soc.*, 1988, **110**, 7980–7984.
13 C. P. Moradi, C. Xie, M. Kaufmann, H. Guo and G. E. Douberly, *J. Chem. Phys.*, 2016, **144**, 164301.
14 J. L. Li and H. Guo, *J. Chem. Phys.*, 2013, **138**, 141102.

Department of Chemistry, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan. E-mail: asukafujii@m.tohoku.ac.jp

15 T. Clark, *J. Comput. Chem.*, 1981, **2**, 261–265.
16 T. Clark, *J. Comput. Chem.*, 1982, **3**, 112–116.
17 T. Clark, *J. Comput. Chem.*, 1983, **4**, 404–409.
18 T. Clark, *J. Am. Chem. Soc.*, 1988, **110**, 1672–1678.