ChemComm

Chemical Communications rsc.li/chemcomm

ISSN 1359-7345



COMMUNICATION

Evamarie Hey-Hawkins *et al.* Molecular doping: accessing the first carborane-substituted 1,2,3-triphospholanide *via* insertion of P⁻ into a P–P bond

ChemComm

COMMUNICATION

Check for updates

Cite this: Chem. Commun., 2019, 55. 3187

Received 9th January 2019, Accepted 3rd February 2019

DOI: 10.1039/c9cc00205g

rsc.li/chemcomm

Molecular doping: accessing the first carborane-substituted 1,2,3-triphospholanide via insertion of P⁻ into a P-P bond⁺

Peter Coburger, 📴 a Hansjörg Grützmacher 📴 bc and Evamarie Hey-Hawkins 🗐 *a

Insertion of a P⁻ anion into a P-P bond yielding the first carboranesubstituted 1,2,3-triphospholanide 1 was achieved by treating a carborane-substitued 1,2-diphosphetane with sodium phosphaethynolate. The triphospholanide 1 can serve as a versatile nucleophilic building block for unprecedent functionalised triphospholanes and carboranesubstituted polyphosphanes.

Polyphosphanes and polyphosphanides have received much interest due to their structural diversity, reactivity and coordination properties.¹ Alkyl and aryl groups have been predominantly used as substituents at the phosphorus centre. ortho-Carborane has been very rarely employed as substituent despite its interesting properties such as a flexible Ccluster-Ccluster bond, its electronwithdrawing effect, the possibility of the hydridic B-H hydrogen atoms to participate in metal bonding, and the possibility of converting the closo-cluster structure into an anionic nidocarborane.² In fact, the only examples of carborane derivatives containing more than one P-P bond are the 1,2,3-triphospholanes Ia-d reported by us (Fig. 1). The respective triphospholanides are absent in the literature,³ but structurally related compounds with more common backbones have been published. Pietschnig et al. reported a series of triphospha-[3]ferrocenophanes IIa-f;⁴ here, the respective triphosphanide can be accessed by deprotonation of IId⁵ which opens up further possibilities for functionalisation to polyphosphane derivatives. Schmidpeter et al. obtained a phenylenesubstituted triphospholanide III via degradation of P_4 with 1,2-phenylenebis(phenylphosphanide). Reactivity studies showed that **III** is a versatile nucleophilic building block; in particular,



 $R^1 = R^2 = Ph$, Id; $R^1 = Ph$, $R^2 = Cv$

la: R' = Cl, IIb: R' = Br, IIc: R' = I, IId: R' = H, Ile: R' = F, IIf: R' = NEt₂ IIg: R' = tBu

Fig. 1 Triphospholane and triphospholanide derivatives. Counterions not shown

a tetraphosphane could be obtained via reaction with chlorodiphenylphosphane.⁶ Here, we present the triphospholanide 1, which is formed upon treatment of a strained carborane-substituted 1,2-diphosphetane⁷ with sodium phosphaethynolate,⁸ and its conversion to new carborane-containing triphospholane derivatives.

The carborane-substituted diphosphetane 2 (Scheme 1) has proven to be a versatile synthon.^{3,7,9} Due to the electron-withdrawing effect of the ortho-carborane backbone, reactivities include the reductive cleavage of the P-P bond with elemental lithium and $[K(thf)_{0,2}]$ $[Co(\eta^4 - cod)_2]$ (cod = 1,5-cyclooctadiene).^{9a,c,d} Diphosphetane 2 also exhibits a considerable ring strain of 12.3 kcal mol⁻¹.^{9e} We therefore concluded that 2 should react with a suitable anionic phosphorus source to yield unprecedented carboranesubstituted phosphanido species. Indeed, treatment of 2 with sodium phosphaethynolate in ethereal solvents afforded the triphospholanide 1 by formal insertion of P⁻ into the P-P bond (Scheme 1). Reactions in which the phosphaethynolate anion



Scheme 1 Reaction of 2 with sodium phosphaethynolate. Counterion (Na⁺) not shown

CHEMISTRY

View Article Online

^a Leipzig University, Faculty of Chemistry and Mineralogy, Institute of Inorganic

Chemistry, Johannisallee 29, D-04103 Leipzig, Germany. E-mail: hey@uni-leipzig.de

^b ETH Zurich, Lab. für Anorganische Chemie, Vladimir-Prelog-Weg 1-5/10, 8093 Zürich, Switzerland

^c Lehn Institute of Functional Materials (LIFM), School of Chemistry,

Sun Yat-sen University, 135 West Xingang Road, Guangzhou 510275, China

[†] Electronic supplementary information (ESI) available: Experimental details, NMR investigations, computational details. CCDC 1888315-1888317. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9cc00205g

(formally) inserts into an element–element bond are extremely scarce and limited to a few examples reported by Goicoechea *et al.* (insertion into a Si–Si bond),¹⁰ Weigand *et al.* (insertion into a Ga–P bond)¹¹ and Driess *et al.* (insertion into a Si–C bond).¹² To a certain extent, these reactions can be viewed as molecular doping reactions with P^- as dopant.¹³

The orange carborane-substituted 1,2,3-triphospholanide 1 is formed as a diastereomeric mixture of the anions *rac*-1 (δ^{31} P = 91.9 (d), -86.3 (t) ppm, ${}^{1}J_{PP}$ = 380 Hz) and *meso-1* ($\delta^{31}P$ = 81.3 (d), -105.6 (t) ppm, ${}^{1}J_{PP} = 280$ Hz) in THF in a 3 · 1 ratio. In contrast, only traces of meso-1 are detectable when the reaction is carried out in DME. Additionally, an epimerisation barrier of 29.5 kcal mol⁻¹ between rac-1 and meso-1 was calculated, whilst the energy difference between the two isomers was calculated to be only -0.1 kcal mol⁻¹ (TPSS-D3BJ/def2-TZVP,¹⁴⁻¹⁶ see Computational details in the ESI[†]). These findings indicate that kinetic reasons are responsible for the formation of the observed diastereomeric mixture. A possible reaction mechanism was investigated employing DFT and ab initio methods (Scheme 2). Structures of isolated molecules were optimised with the TPSS-D3BJ^{14,15} functional and the def2-TZVP basis,15 and energies were calculated with the DLPNO-CCSD(T)¹⁷ method by Neese et al. and the def2-QZVPP basis¹⁶ (see Computational details in the ESI[†]). In the first step, the P-P bond in diphosphetane 2 is broken upon attack of the phosphaethynolate anion at one phosphorus atom forming the phosphanido-phosphaketene A. This step proceeds *via* a very low energy barrier ($\Delta E^{\#} = 6.8 \text{ kcal mol}^{-1}$) and is slightly exothermic ($\Delta E = -3.3$ kcal mol⁻¹, see the ESI[†]). Notably, the reaction energy is significantly smaller than the ring strain of 2 (12.3 kcal mol^{-1})^{9e} indicating that release of steric strain might strongly contribute to the driving force of the initial step. The phosphanido moiety in A can react intramolecularly with the phosphaketene moiety in two ways. One possibility is the elimination of carbon monoxide via transition state B forming the triphospholanide *meso-1* irreversibly. The other possibility is the nucleophilic attack at the carbon atom of the phosphaketene moiety via transition state C forming the heterocyclic compound D. Due to the rather high activation energy of the backwards reaction

 $(\Delta E^{\#} = 25.5 \text{ kcal mol}^{-1}, \text{ corresponding to a } t_{1/2} \text{ of } 6.5 \text{ days at r.t.})$ this step is irreversible as well.

Subsequently, **D** can undergo epimerisation (**E**, **F**) followed by ring opening to form the phosphanido-phosphaketene **H**. Finally, elimination of CO *via* transition state **I** yields the triphospholanide *rac-1*. All transition states in the calculated mechanism are accessible at room temperature. In particular, the transition states **B** and **C**, which determine the ratio of *meso-1* to *rac-1*, are close in energy in agreement with the formation of a diastereomeric mixture of **1**. When explicit counterion and solvation effects are incorporated into the calculation, their energy is even closer ($\Delta E^{\#}(\mathbf{B}) = 18.3 \text{ kcal mol}^{-1}, \Delta E^{\#}(\mathbf{C}) = 18.7 \text{ kcal mol}^{-1}$, at the same level of theory, see the ESI[†] for further details).

In the next step, the reactivity of triphospholanide 1 towards several electrophiles (Scheme 3) was probed. Protonation of 1 with water in *n*-hexane afforded the secondary triphospholane **1-H** in good yields (72%). Triphospholane **1-H** is obtained as a colourless solid and can be stored and handled under ambient



Scheme 3 Reactions of ${\bf 1}$ with electrophiles. Counterion (Na^+) in $\it rac/meso-1$ not shown.



Scheme 2 Calculated mechanism for the formation of *meso-1* and *rac-1*. Relative energies are given in kcal mol⁻¹. Counterions (Na⁺) not included.

conditions and can be deprotonated again quantitatively with *n*BuLi at 0 °C to yield **1**. Therefore, we used **1-H** as a convenient source of 1 in further reactivity studies. 1 can be converted to the respective 2-chlorotriphospholane 1-Cl by treatment with hexachloroethane as chloronium ion source in n-hexane. 1-Cl was isolated as an air-stable colourless solid in moderate yield (39%). Triphospholanide 1 can also serve as a building block for more complex polyphosphanes. Thus, the 2:1 reaction with dichlorophenylphosphane affords the heptaphosphane 3 as a colourless solid in 63% yield. 3 is the first carborane-substituted polyphosphane with more than three phosphorus atoms. Of the possible stereoisomers (two rac and two meso isomers),⁴ only one meso isomer is formed in the case of 1-Cl and 3 as confirmed by single-crystal X-ray diffraction studies (vide infra). In contrast, triphospholane 1-H is always obtained as a mixture of a rac and two meso isomers, even when triphospholanide 1 was prepared in DME and thus contained only traces of *meso-1*.

Additional NMR spectroscopic investigations proved that the isomers slowly form an equilibrium in solution with the meso isomers being the major compounds (see the ESI⁺). Overall, the observed stereochemistry of the triphospholane derivatives is in good agreement with a detailed theoretical study (ESI⁺). Single crystals of the triphospholane derivatives rac-1-H, 1-Cl and 3 suitable for X-ray structure analysis were obtained.[‡] The molecular structures of rac-1-H, 1-Cl and 3 are depicted in Fig. 2, and selected bond lengths and angles are summarised in Table 1. The C₂P₃ ring in rac-1-H adopts a distorted, non-symmetrical envelope conformation with two different P-P bond lengths [P1-P3: 2.238(2) Å, P2-P3: 2.192(3) Å] within the standard range of P-P single bonds.¹⁸ rac-1-H also displays a non-symmetric conformation in solution as indicated by the presence of two distinct resonances accounting for the different tert-butylphosphanyl groups in the ³¹P{¹H} NMR spectrum (see the ESI[†]). *meso-1-Cl* crystallises in the monoclinic space group C2/c; the C_2P_3 ring adopts an



Fig. 2 Molecular structures of *rac-1-H*, *1-Cl* and **3** with ellipsoids at the 50% probability level. Hydrogen atoms (other than P3-H1 in *rac-1-H*), disorder within the C_2P_3 ring and the *tert*-butyl groups (in case of *1-H*), co-crystallised *n*-hexane (*1-Cl*) and toluene (*3*) are omitted for clarity.

Table 1 Selected bond lengths [Å] and angles [°] for *rac-1-H*, *1-Cl* and **3** (for clarity, values for the second molecule in the asymmetric unit of **3** are given in the ESI). Numbering scheme according to the respective depiction of the solid-state structures

	<i>rac</i> -1-H	1-Cl	3
C1-C2 (C3-C4) P1-P3 (P5-P6) P2-P3 (P5-P7) P3-P4/Cl1 (P4-P5) P1-P3-P2 (P6-P5-P7) P1-C1-C2 (P6-C3-C4) P2-C2-C1 (P7-C4-C3) P3-P1-C1 (P5-P6-C3)	<i>rac</i> -1-H 1.680(4) 2.238(2) 2.192(3) 97.26(9) 112.6(2) 117.9(2) 100.2(1)	1-Cl 1.689(3) 2.2292(8) 2.2227(9) 2.0846(7) 93.62(3) 114.7(1) 114.3(1) 95.68(7)	3 1.699(2) (1.675(3)) 2.2189(6) (2.2311(6)) 2.2108(6) (2.2276(6)) 2.2428(6) (2.2444(6)) 102.86(2) (92.25(2)) 118.2(1) (114.6(1)) 118.1(1) (113.5(1)) 99.51(6) (94.79(5))
P3–P2–C2 (P5–P7–C4) P3–P4–P5	100.9(1) —	95.68(7) —	99.84(6) (95.63(5)) 99.08(2)

envelope conformation. Overall, the structure of the C_2P_3 rings in *rac*-1-H and 1-Cl is rather relaxed with bond lengths and angles similar to the ones observed in triphospholanes (Fig. 1).^{3,18} This is also the case for the triphospholane rings in 3. However, they markedly differ from each other [*e.g.* P1–P3–P2 is 102.86(2)°, whilst P6–P5–P7 is 92.25(2)°, Table 1]. This unsymmetrical conformation might be adopted to minimise steric repulsion between the substituents on the phosphorus atoms. Additionally, weak attractive interactions between P4 and P1, P4 and P2, P4 and P6, P4 and P7, and P3 and P5 are indicated by their rather short distances [3.2070(7)–3.5687(7) Å, Fig. S4, ESI†] which are well within the sum of their van der Waals radii (3.6 Å).¹⁹ The presence of attractive interactions is further supported by an NBO analysis where Wiberg bond indices (WBI) of 0.02 to 0.04 are found between the respective phosphorus atoms (see Computational details in the ESI†).

The ${}^{31}P{}^{1}H$ NMR spectra of the triphospholane derivatives are in accordance with their solid-state structures with ${}^{1}J_{PP}$ coupling constants of 1-H and 1-Cl in the range of those observed for the triphospholanes Ia-d (ESI⁺).³ In solution, meso-3 displays four multiplets in the ³¹P{¹H} NMR spectrum at room temperature which cannot be assigned easily because the NMR spectra of meso-3 are strongly temperature dependent indicating hindered rotations at room temperature (see the ESI,† Fig. S6 and S7). The multiplets corresponding to the nuclei P3, P4 and P5 become remarkably simple at elevated temperatures. Subsequently, the signals could be simulated as the X and YY' part of the expected AA'BB'XYY' spin system at 60 °C, yielding the following chemical shifts and coupling constants: $\delta_A = \delta_{A'} = 62.9$ ppm, $\delta_{\rm B} = \delta_{\rm B'} = 59.2$ ppm, $\delta_{\rm X} = -4.2$ ppm, $\delta_{\rm Y} = \delta_{\rm Y'} = -16.9$ ppm; $J_{\rm AA'} = -16.9$ ppm; J_{\rm $J_{BB'} = J_{AB'} = J_{BA'} = 0$ Hz, $J_{AB} = J_{A'B'} = 5$ Hz, $J_{BY'} = J_{B'Y} = 20$ Hz, $J_{AY'} = J_{AY'} =$ $J_{A'Y} = 34 \text{ Hz}, J_{YY'} = 86 \text{ Hz}, J_{BX} = J_{B'X} = 180 \text{ Hz}, J_{AX} = J_{A'X} = 188 \text{ Hz},$ $J_{XY} = J_{XY'} = -188 \text{ Hz}, J_{BY} = J_{B'Y'} = -225 \text{ Hz}, J_{AY} = J_{A'Y'} = -233 \text{ Hz}$ (see Fig. S8, ESI,† for a comparison of the experimental and the simulated spectrum). The large ${}^{2}J_{PP}$ coupling constants between P4 and P1, P2, P6 and P7 (180 and 188 Hz) may indicate attractive interactions between these phosphorus atoms even in solution. Significant couplings could also be observed in a ³¹P-³¹P COSY experiment (Fig. S9, ESI⁺). Through-space interactions between phosphorus atoms are also commonly observed in carboranesubstituted bisphosphanes and result in coupling constants of similar magnitudes.^{2c}

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

In summary, the first anionic carborane-substituted triphospholanide **1** was obtained by reacting diphosphetane **2** with the phosphaethynolate anion. Possible mechanisms of this reaction in which OCP⁻ acts as source for the molecular dopant P⁻ were revealed by DFT calculations. Triphospholanide **1** is a valuable synthon and can be easily converted to **1-H**, **1-Cl** and **3** upon reaction with the respective electrophiles. Especially **1-H** and **1-Cl** are promising stable starting materials for further reactivity studies due to the presence of a reactive P–H or P–Cl bond which will eventually allow to incorporate these as building blocks in polymers and other materials. These studies are presently underway.

We gratefully acknowledge financial support from the Studienstiftung des deutschen Volkes (doctoral grant for P. C.). Further support came from the Lehn Institute of Functional Materials (LIFM), School of Chemistry, Sun Yat-sen university, the National Natural Science Foundation of China (NSFC) (project 21720102007), and the ETH Zürich, (project 0-20406-18). We thank Gabriele Hierlmeier (University of Regensburg) for help with the synthesis of sodium phosphaethynolate.

Conflicts of interest

There are no conflicts to declare.

Notes and references

‡ *rac*-1-H co-crystallises with *meso*-1-H a as minor component and additionally displays disorder. Consequently, the precision of the bond lengths and angles is lower in comparison to 1-Cl and 3. However, they are in good agreement with the optimised geometry of *rac*2-1-H in the gas phase (see the ESI† for the Cartesian coordinates).

- (a) M. Baudler and K. Glinka, Chem. Rev., 1993, 93, 1623–1667;
 (b) M. Baudler and K. Glinka, Chem. Rev., 1994, 94, 1273–1297;
 (c) S. Gómez-Ruiz and E. Hey-Hawkins, Coord. Chem. Rev., 2011, 255, 1360–1386; (d) K.-O. Feldmann and J. J. Weigand, J. Am. Chem. Soc., 2012, 134, 15443–15456; (e) A. Wiśniewska, R. Grubba, Ł. Ponikiewski, M. Zauliczny and J. Pikies, Dalton Trans., 2018, 47, 10213–10222.
- 2 (a) V. I. Bregadze, Chem. Rev., 1992, 92, 209–223; (b) M. Scholz and E. Hey-Hawkins, Chem. Rev., 2011, 111, 7035–7062; (c) A. R. Popescu, F. Teixidor and C. Viñas, Coord. Chem. Rev., 2014, 269, 54–84.

- 3 A. Kreienbrink, S. Heinicke, T. T. Duong Pham, R. Frank, P. Lönnecke and E. Hey-Hawkins, *Chem. – Eur. J.*, 2014, **20**, 1434–1439.
- 4 S. Borucki, Z. Kelemen, M. Maurer, C. Bruhn, L. Nyulászi and R. Pietschnig, *Chem. – Eur. J.*, 2017, 23, 10438–10450.
- 5 S. Isenberg, L.-M. Frenzel, C. Bruhn, R. Pietschnig, S. Isenberg, L.-M. Frenzel, C. Bruhn and R. Pietschnig, *Inorganics*, 2018, **6**, 67.
- 6 A. Schmidpeter, G. Burget and W. S. Sheldrick, *Chem. Ber.*, 1985, 118, 3849–3855.
- 7 A. Kreienbrink, M. B. Sárosi, E. G. Rys, P. Lönnecke and E. Hey-Hawkins, Angew. Chem., Int. Ed., 2011, 50, 4701–4703 (Angew. Chem., 2011, 123, 4798–4800).
- 8 H. Grützmacher and J. Goicoechea, Angew. Chem., Int. Ed., 2018, 57, 16968–16994 (Angew. Chem., 2018, 130, 17214–17240).
- 9 (a) A. Kreienbrink, P. Lönnecke, M. Findeisen and E. Hey-Hawkins, *Chem. Commun.*, 2012, 48, 9385–9387; (b) P. Coburger, J. Schulz, J. Klose, B. Schwarze, M. B. Sárosi and E. Hey-Hawkins, *Inorg. Chem.*, 2017, 56, 292–304; (c) P. Coburger, S. Demeshko, C. Rödl, E. Hey-Hawkins and R. Wolf, *Angew. Chem., Int. Ed.*, 2017, 56, 15871–15875 (*Angew. Chem.*, 2017, 129, 16087–16091); (d) J. Schulz, A. Kreienbrink, P. Coburger, B. Schwarze, T. Grell, P. Lönnecke and E. Hey-Hawkins, *Chem. – Eur. J.*, 2018, 24, 6208–6216; (e) P. Coburger, R. Aures, P. Schulz and E. Hey-Hawkins, *ChemPlusChem*, 2018, 83, 1057–1064.
- 10 T. P. Robinson, M. J. Cowley, D. Scheschkewitz and J. M. Goicoechea, Angew. Chem., Int. Ed., 2015, 54, 683–686 (Angew. Chem., 2015, 127, 693–696).
- 11 F. Hennersdorf, J. Frötschel and J. J. Weigand, J. Am. Chem. Soc., 2017, 139, 14592–14604.
- 12 M. Driess, A. Hermannsdorfer and D. W. Stephan, *Chem. Commun.*, 2018, **54**, 13523–13526.
- 13 (a) J. Li, G. D'Avino, A. Pershin, D. Jacquemin, I. Duchemin, D. Beljonne and X. Blase, *Phys. Rev. Mater.*, 2017, **1**, 025602; (b) R. Fujimoto, Y. Yamashita, S. Kumagai, J. Tsurumi, A. Hinderhofer, K. Broch, F. Schreiber, S. Watanabe and J. Takeya, *J. Mater. Chem. C*, 2017, **5**, 12023–12030; (c) I. Salzmann and G. Heimel, *J. Electron Spectrosc. Relat. Phenom.*, 2015, **204**, 208–222.
- 14 J. Tao, J. P. Perdew, V. N. Staroverov and G. E. Scuseria, *Phys. Rev. Lett.*, 2003, **91**, 146401.
- 15 (a) S. Grimme, S. Ehrlich and L. Goerigk, J. Comput. Chem., 2011, 32, 1456–1465; (b) S. Grimme, J. Antony, S. Ehrlich and H. Krieg, J. Chem. Phys., 2010, 132, 154104.
- 16 F. Weigend and R. Ahlrichs, Phys. Chem. Chem. Phys., 2005, 7, 3297-3305.
- (a) C. Riplinger, B. Sandhoefer, A. Hansen and F. Neese, *J. Chem. Phys.*, 2013, **139**, 134101; (b) D. G. Liakos, M. Sparta, M. K. Kesharwani, J. M. L. Martin and F. Neese, *J. Chem. Theory Comput.*, 2015, **11**, 1525–1539.
- 18 A. H. Cowley, Chem. Rev., 1965, 65, 617-634.
- 19 A. Bondi, J. Phys. Chem., 1964, 68, 441-451.