

## PAPER

View Article Online  
View Journal | View IssueCite this: *Dalton Trans.*, 2019, **48**,  
2038Received 6th December 2018,  
Accepted 8th January 2019

DOI: 10.1039/c8dt04818e

rsc.li/dalton

Electrophilic boron carboxylate and phosphinate  
complexes†Diya Zhu,<sup>a</sup> James H. W. LaFortune,<sup>a</sup> Rebecca L. Melen <sup>\*b</sup> and Douglas W. Stephan <sup>\*a</sup>


The reactions of a series of carboxylic acids with  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\text{SMe}_2$  are shown to afford species of the form  $[\text{RC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2\text{O}$  ( $\text{R} = \text{Tol}$  **1**,  $\text{Ph}$  **2**,  $\text{C}_6\text{F}_5$  **3**,  $\text{Me}_2\text{BrC}$  **4**,  $\text{Me}$  **5**) in 87–95% yields with the concurrent reduction of the carboxylic acid to the corresponding aldehyde. A mechanism for the formation of **1–5** is proposed to proceed via a cyclic eight-membered ring species. Analogues of these species were prepared via reactions of carboxylic and phosphinic acids with  $\text{HB}(\text{C}_6\text{F}_5)_2$  and  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\text{SMe}_2$ , respectively, to give  $[\text{ToI}(\text{C}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2)_2]_2$  **6**,  $[(\text{C}_6\text{F}_5)\text{C}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2$  **7**, and  $[\text{Ph}_2\text{P}(\text{O})\text{OBH}(\text{C}_6\text{F}_5)_2]_2$  **8**. These products react subsequently to give  $\text{ToI}(\text{C}(\text{O})\text{OBH}(\text{C}_6\text{F}_5)(\text{NC}_5\text{H}_4\text{NMe}_2))$  **9** and  $\text{Ph}_2\text{P}(\text{O})\text{OBH}(\text{C}_6\text{F}_5)(\text{NC}_5\text{H}_4\text{NMe}_2)$  **10**. The acyloxyborate derivatives **1–4** were shown to be inactive in mediating the direct amidation of carboxylic acids, consistent with previous observations that infer the need for a sterically congested environment about the boron centres.

## Introduction

The formulation of acyloxyboranes generated a lot of confusion in the early literature, specifically over the formulation of the product derived from the reaction of orthoboric acid and acetic anhydride. Originally thought to be boron acetate, this notion was revised in 1957 when Hayter *et al.*<sup>1</sup> correctly formulated the product as the acyloxyborate  $[\text{RC}(\text{O})\text{OBR}']_2\text{O}$  (Scheme 1). Subsequently, a number of related studies have provided structural data or established new protocols to such species. These include reports by the groups of Lappert,<sup>2</sup> Perotti,<sup>3</sup> Sporzynski,<sup>4</sup> Köster,<sup>5</sup> and Wrackmeyer.<sup>6</sup> Despite the relatively few studies of these compounds, acyloxyborates have found important applications in fire-proofing polymer compositions, as stabilisers for synthetic rubbers, and in the pharmaceutical industry.<sup>7</sup>

Related boron carboxylate species have also been proposed as intermediates in dehydrative condensations between a carboxylic acid and an amine using catalysts derived from borate esters<sup>8–10</sup> as well as boric,<sup>11</sup> boronic,<sup>12–21</sup> and borinic acids.<sup>22</sup> Recently however, Whiting and coworkers<sup>23</sup> have suggested an alternative mechanism in which an oxo-bridged bis-boron species, *i.e.* an acyloxyborate, acts as the catalytically active

species for direct amidation reactions (Scheme 1). In this catalysis, Whiting<sup>23</sup> showed that acyloxyborates are generated *in situ* by the reaction of boronic and carboxylic acids in the presence of molecular sieves, although these authors were able to prepare related species directly from the reaction of an



**Scheme 1** (a) Selected synthetic routes to acyloxyborate derivatives. (b) Direct amidation catalysis by acyloxyborates.

<sup>a</sup>Department of Chemistry, University of Toronto, M5S3H6, Canada.

E-mail: dstephan@chem.utoronto.ca

<sup>b</sup>School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff, CF10 3AT, Cymru/Wales, UK

† Electronic supplementary information (ESI) available: Synthetic and spectral details. CCDC 1883339–1883344. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8dt04818e

arylboronic acid and phenylacetic acid. The intermediacy of these acyloxyborates was further supported by Ishihara and co-workers,<sup>24</sup> who showed that *ortho*-substituents on the boron-bound aryl group prevented the coordination of amine to boron, thus accelerating the catalysis.

Our interest in boron-based species that incorporate electron withdrawing substituents has prompted us to probe related carboxylate derivatives. As is well established that borohydride reductions of carboxylic acids proceed *via* boron-carboxylate species,<sup>25,26</sup> we were interested in probing the reactions of carboxylic acids with the electrophilic boranes. In the initial effort, we previously reported the preparation of the salt  $[\text{Cp}^*_2\text{Fe}][\text{PhCO}_2\text{B}(\text{C}_6\text{F}_5)_3]$  *via* the one electron reduction of a peroxide in the presence of a borane.<sup>27,28</sup> Herein, we describe the reactions of  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\cdot\text{SMe}_2$ <sup>29</sup> and  $\text{HB}(\text{C}_6\text{F}_5)_2$  with carboxylic and phosphinic acids. Generally, these reactions result in eight-membered cyclic products, while the reactions of  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\cdot\text{SMe}_2$  and carboxylic acids provide a facile route to bicyclic acyloxyborate derivatives with concurrent acid-reduction to aldehyde. Experimental and computational data support a proposed mechanism for the formation of the latter products. The catalytic utility of these acyloxyborate species in the direct amidation reactions is also probed.

## Results and discussion

The reaction of *p*-toluic acid with Lancaster's reagent,  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\cdot\text{SMe}_2$ ,<sup>29</sup> was performed in DCM at room temperature prompting the evolution of  $\text{H}_2$ . Repeated reactions showed that all the reagents were consumed when combined in an acid to borane ratio of 3 : 2. The  $^{11}\text{B}$  NMR spectrum revealed the complete conversion of  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\cdot\text{SMe}_2$  into a new four-coordinate boron species that exhibited a broad resonance at 5.1 ppm. The  $^{19}\text{F}\{^1\text{H}\}$  NMR spectrum showed a gap between the resonances attributed to the *meta*- and *para*-fluorine atoms of the perfluorinated arene rings ( $\Delta\delta = 8.3$  ppm), consistent with the presence of a four-coordinate boron centre. After work-up, a white solid **1** was isolated in 95% yield (Scheme 2). Single crystals for X-ray diffraction analysis were obtained through diffusion of pentane into a benzene solution at ambient temperature (Fig. 1). The solid-state structure showed that **1** was  $[\text{ToIc}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2\text{O}$ , a [3,3,1] bicycle in which two





Fig. 1 POV-ray depictions of (a) **1**, (b) **2**, and (c) **3**. C: black, O: red, B: yellow-green, and F: pink. Hydrogen atoms have been omitted for clarity.

boron centres are linked by a bridging oxygen atom and two carboxylate ligands.

In a similar fashion, benzoic acid, pentafluorobenzoic acid, 2-bromo-2-methylpropionic acid, and acetic acid reacted with  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\cdot\text{SMe}_2$  to give the products formulated as  $[\text{RC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2\text{O}$  ( $\text{R} = \text{Ph}$  **2**,  $\text{C}_6\text{F}_5$  **3**,  $\text{Me}_2\text{BrC}$  **4**,  $\text{Me}$  **5**) in 87–90% yields (Scheme 1). The spectroscopic data for these compounds were similar to those described for **1**. Crystallographic studies also confirmed the formulations of **2** and **3** (Fig. 1).

The structural data for **1**–**3** confirmed a pseudo-tetrahedral geometry about the two boron centres linked by an oxygen atom and bridged by two carboxylate units. The B–O–B fragments exhibit B–O bond lengths of 1.398(2) and 1.401(2) Å in **1**, 1.401(3) and 1.402(3) Å in **2**, and 1.394(4) and 1.400(4) Å in **3**. The corresponding B–O–B angles were found to be 110.6(2)°, 110.5(2)°, and 112.9(2)°, respectively. The B–O bond distances for the carboxylic oxygen atoms in these species were found to fall in the range of 1.522(3) to 1.612(3) Å. The B–C bond lengths in **1** and **2** ranged from 1.602(3) to 1.612(3) Å, while those in **3** were found to be 1.595(4) and 1.582(4) Å. The shorter B–C bonds in **3** are consistent with the presence of the electron withdrawing carboxylates. Nonetheless, these general



Scheme 2 Synthesis of **1**–**5**.




structural features are similar to those previously reported for the diacyloxy B–O–B species  $[(\text{MeC}(\text{O})\text{O})_2\text{B}]_2\text{O}$ ,<sup>3</sup>  $[\text{MeC}(\text{O})\text{OBET}]_2\text{O}$ ,<sup>5</sup> and  $[\text{MeC}(\text{O})\text{OBCy}]_2\text{O}$ <sup>6</sup> as reported by Perotti, Köster, and Wrackmeyer, respectively.

In addition to  $\text{H}_2$ , the second by-product in the formation of 1–5 was identified as the aldehyde derived from the reduction of the starting carboxylic acid. Examination of the reaction mixture by DART-MS confirmed the formation of the corresponding aldehyde (for details see the ESI†). In addition, the  $^1\text{H}$  NMR spectrum of the reaction mixture of compound 1 showed a broad resonance *ca.* 12 ppm, supporting the presence of the corresponding aldehyde.

The mechanism of the formation of 1–5 is thought to be initiated by protonolysis generating boryl-ester. Dimerization of this species is consistent with the electrophilic nature of the boron centre. In the subsequent reaction with a third equivalent of acid, its carbonyl fragment is in close proximity to a boron hydride, which presumably results in the liberation of aldehyde and the formation of the oxo-bridge between the two boron centres (Scheme 3).

In support of the proposed initial dimerization in this mechanism, a related product was derived from the reaction of *p*-toluic acid with Piers' borane,  $\text{HB}(\text{C}_6\text{F}_5)_2$ .<sup>30</sup> This reaction afforded the near quantitative formation of a product formulated as  $[\text{TolC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2$  6 (Scheme 4). The  $^{19}\text{F}$  NMR spectrum showed resonances at  $-135.1$ ,  $-154.0$ , and  $-162.1$  ppm, while the  $^{11}\text{B}$  NMR signal was observed at 5.2 ppm. These data are consistent with four-coordinate boron centres and this was confirmed crystallographically (Fig. 2). The structure demonstrates the eight-membered ring in which the carboxylic units bridge the two boron centres. This species is structurally similar to dialkyl-group 13 carboxylates reported by Justyniak *et al.*<sup>31</sup> as well as the species  $[\text{Ph}_2\text{PO}_2\text{B}(\text{C}_6\text{F}_5)_2]_2$ .<sup>32</sup> Overall, this ring adopts a pseudo-boat conformation in which the  $\text{C}_6\text{F}_5$  rings on each of the boron atoms are oriented so as to permit  $\pi$ -stacking. This observation appears to be a solid-state packing effect as this inequivalence of the fluoroarene rings is not reflected in the  $^{19}\text{F}$  NMR spectral data.

In a similar fashion, the corresponding reaction of pentafluorobenzoic acid with Piers' borane,  $\text{HB}(\text{C}_6\text{F}_5)_2$ , afforded the



Scheme 4 Synthesis of 6–8.





Fig. 2 POV-ray depiction of 6. C: black, O: red, B: yellow-green, and F: pink. All hydrogen atoms have been omitted for clarity.

product formulated as  $[(\text{C}_6\text{F}_5)\text{C}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2$  7 in 83% isolated yield (Scheme 4). While the  $^{19}\text{F}$  and  $^{11}\text{B}$  NMR data were consistent with the formulation of 7, its poor solubility precluded the acquisition of  $^{13}\text{C}$  data.

In an analogous reaction, the combination of stoichiometric amounts of  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\text{SMe}_2$  and  $\text{Ph}_2\text{P}(\text{O})\text{OH}$  was performed. This led to the release of  $\text{H}_2$  and the formation of a clear and colourless solution after 12 h (Scheme 4). A colourless crystalline solid 8 was isolated in 68% yield after workup. Dissolution of 8 in  $\text{CDCl}_3$  gives rise to two  $^{31}\text{P}$  signals at 40.8 and 37.6 ppm in a 1 : 0.44 ratio. Similarly, two sets of pentafluorophenyl resonances in the  $^{19}\text{F}$  NMR spectrum suggest the presence of two four-coordinate boron environments. Nonetheless, only a single, broad  $^{11}\text{B}$  NMR signal was seen at 0.3 ppm while the corresponding  $^1\text{H}\{^{11}\text{B}\}$  NMR spectrum showed two broad resonances at 4.18 and 3.91 ppm attributed to two B–H environments. To probe the possibility of a monomer/dimer equilibrium (Scheme 4), a variable temperature multinuclear NMR study was performed on 8. Upon cooling to  $-30^\circ\text{C}$ , the ratio of the two resonances at 40.8 and




Scheme 3 Proposed mechanism for the formation of 1–5.



37.6 ppm was altered slightly to 1:0.56. These data suggest a dynamic equilibrium in which a dimer formulated as  $[\text{Ph}_2\text{P}(\text{O})\text{OBH}(\text{C}_6\text{F}_5)]_2$  dissociates to a monomeric species. Subsequently, single crystals of **8** suitable for X-ray diffraction analysis were obtained by cooling a saturated dichloromethane solution of **8** at  $-35^\circ\text{C}$ . The solid-state structure revealed a dimeric structure with the expected eight-membered ring, adopting a pseudo-chair conformation (Fig. 3). The B–O and P–O bond distances were found to be on average 1.438(6) and 1.525(3) Å, respectively.

The nature of **8** is analogous to the proposed intermediate **Int**, in the above reactions with carboxylic acids. However, in contrast to the carboxylic acid analogues, efforts to react **8** with an excess amount of diphenylphosphinic acid or with carboxylic acids failed. In addition, **8** did not react with 1,4-pentadiene in toluene even on heating to  $110^\circ\text{C}$  for 12 h. The latter result is consistent with the strong basicity of the phosphinic acid fragment that binds to boron, precluding the generation of a transient three-coordinate boron centre necessary for hydroboration.

While the NMR data for **8** suggests a monomer/dimer equilibrium, the corresponding data for **6** indicate a robust eight-membered ring. Nonetheless, the reaction of **6** and **8** with DMAP resulted in the formation of new species **9** and **10**, respectively. In the case of **9**, the  $^{19}\text{F}$  NMR spectrum showed signals at  $-133.5$ ,  $-157.3$ , and  $-163.6$  ppm while the  $^{11}\text{B}$  resonance was seen at 1.6 ppm, consistent with a four-coordinate boron centre. A similar conclusion was drawn for **10** based on the  $^{19}\text{F}$  NMR signals at  $-134.9$ ,  $-158.1$ , and  $-164.0$  ppm, with the  $^{11}\text{B}$  signal at 0.8 ppm. The  $^{31}\text{P}$  NMR resonance for **10** was seen at 25.3 ppm. These data suggest the coordination of DMAP to the boron centres of **9** and **10**, resulting in products  $\text{TolCO}_2\text{BH}(\text{C}_6\text{F}_5)(\text{NC}_5\text{H}_4\text{NMe}_2)$  and  $\text{Ph}_2\text{PO}_2\text{BH}(\text{C}_6\text{F}_5)(\text{NC}_5\text{H}_4\text{NMe}_2)$ , respectively (Scheme 5). These formulations were confirmed crystallographically (Fig. 4). The B–O distances in **9** and **10** were found to be 1.484(3) and 1.485(3) Å, while the B–N distances were 1.605(3) and 1.595(4) Å, respectively.



Scheme 5 Synthesis of **9**–**10**.




Fig. 4 POV-ray depictions of (a) **9** and (b) **10**. C: black, O: red, B: yellow-green, F: pink, P: orange, N: blue, and H: white. All hydrogen atoms except those on boron have been omitted for clarity.




Fig. 3 POV-ray depiction of **8**. C: black, O: red, B: yellow-green, F: pink, P: orange, and H: white. All hydrogen atoms except those on boron have been omitted for clarity.

To probe the stability of **8** further, DFT computations were performed at the M062X/def2-TZVPP level of theory with GD3 dispersion and PCM modelling of dichloromethane solvation. Given that **8** is the phosphinic acid analogue of the proposed intermediate, **Int1** (Scheme 2), in the formation of **1**–**5**, the optimized structures of **8** and **Int1** were computed. Interestingly, the lowest energy conformation of **8** is one in which the hydride atoms on boron adopt a *transoid* conformation. In contrast, for **Int1** the orientation in which the hydride atoms are *cisoid* is lower in energy. This difference is attributed to the minimization of steric conflict in **8**. The HOMO for **8** is located on the fluoroarene rings, whereas the hydride atoms on boron in **Int1** contributed to the HOMO (Fig. 5).








Fig. 5 Surface contour plots (isovalue, 0.03) of the HOMO of (a) **8** and (b) **Int1**.

Finally, as Whiting has proposed that the active catalysts in their direct amidation reactions are structurally analogous to **1–5**, the utility of the present compounds was probed. Addition of two equivalents of aniline to the species **1–4** generated insoluble products that could not be characterised. However, the corresponding reaction of **5** gave soluble products and showed the generation of a trace amount of amide after 12 h as evidenced by the  $^1\text{H}$  NMR spectrum (see the ESI†). The corresponding  $^{11}\text{B}$  NMR spectrum revealed the consumption of **5** and the appearance of two signals at 3 and 18 ppm. These results were reminiscent of data presented by Whiting<sup>23</sup> and Ishihara<sup>24</sup> for analogous species with sterically unencumbered boron centres. Although these products could not be isolated from the present reaction, the inability of **5** to catalyse amide formation is consistent with the need for steric hindrance about the boron atom to prompt nucleophilic attack at the carbon of a bridging carboxylic group, thereby prompting the amidation pathway as proposed by Whiting and Ishihara. This notion is also supported by the formation of **9** and **10** which illustrates the ready access of a donor to the respective boron sites in **6** and **8**.

## Conclusions

In summary, synthetic methods to new acyloxyborates of the formula  $[\text{RC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)]_2\text{O}$  (**1–5**) are derived from the reactions of  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\text{SMe}_2$  with carboxylic acids. These species are formed *via* reactions involving protonolysis of the B–H bonds and reduction of carboxylic acid to generate the corresponding aldehyde. The mechanism is proposed to proceed *via* a cyclic intermediate that dissociates and allows the B–H bonds to react with an additional equivalent of carboxylic acid. This proposed intermediate is analogous to the cyclic species **6–8**. For the latter species, the lack of steric effects of the *ortho*-substituent on the boron atoms results in dimer cleavage and facile binding of donors to the boron centres affording **9** and **10**. These observations together with the ineffective catalytic activity of **1–5** in direct amidation reactions are consistent with the previous inference of the need for steric congestion to generate a catalytically active acyloxyborate.<sup>23,24</sup> Efforts to use

boron carboxylate derivatives as catalysts in a variety of reactions are the subject of ongoing study.

## Experimental section

### General remarks

All reactions and work-up procedures were performed under an inert atmosphere of dry, oxygen-free  $\text{N}_2$  by means of standard Schlenk techniques or glovebox techniques (MBRAUN glovebox equipped with a  $-35\text{ }^\circ\text{C}$  freezer) unless otherwise specified. All glassware was oven-dried and cooled under vacuum before use. Dichloromethane (DCM) was distilled over  $\text{CaH}_2$ . Pentane was collected from a Grubbs-type column system manufactured by Innovative Technology and degassed. Solvents were stored over activated 4 Å molecular sieves. Molecular sieves, type 4 Å (pellets, 3.2 mm diameter), purchased from Sigma Aldrich were activated prior to usage by iteratively heating under vacuum for 24 hours.  $\text{CDCl}_3$  purchased from Cambridge Isotope Laboratories was vacuum distilled, further degassed, and stored over activated 4 Å molecular sieves in a glovebox for at least 8 hours prior to use. Unless otherwise mentioned, chemicals were purchased from Sigma Aldrich or TCI. Lancaster's reagent  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\text{SMe}_2$ <sup>29</sup> and Piers' borane  $\text{HB}(\text{C}_6\text{F}_5)_2$ <sup>30</sup> were prepared using literature methods. NMR spectra were recorded at room temperature (298 K) unless otherwise mentioned on a Bruker Avance III 400 MHz, an Agilent DD2 400, and an Agilent DD2 500. Spectra were referenced to the residual solvent signals ( $\text{CDCl}_3$ :  $^1\text{H}$  = 7.26;  $^{13}\text{C}$  = 77.2 ppm; toluene- $d_8$ :  $^1\text{H}$  = 7.09, 7.01, 6.97, and 2.08 ppm and  $^{13}\text{C}$  = 137.48, 128.87, 17.96, 125.13, and 20.43 ppm). Chemical shifts ( $\delta$ ) are reported in ppm and coupling constants ( $J$ ) are listed as absolute values in Hz. Multiplicities are reported as singlet (s), doublet (d), triplet (t), multiplet (m), overlapping (ov), and broad (br). High-resolution mass spectra (HRMS) were obtained on a JMS-T100LC JOEL DART mass spectrometer. Elemental analyses for C, H, and N were performed by ANALEST (University of Toronto) employing a PerkinElmer 2400 Series II CHNS Analyser.

### X-ray diffraction studies

Single crystals were coated with paratone oil, mounted on a cryoloop and frozen under a stream of cold nitrogen. Data were collected on a Bruker Apex2 X-ray diffractometer at 150(2) K for all crystals using graphite monochromated Mo-K $\alpha$  radiation (0.71073 Å). Data were collected using Bruker APEX-2 software and processed using SHELX and an absorption correction applied using multi-scan within the APEX-2 program. All structures were solved and refined by direct methods within the SHELXTL package.

**Synthesis of  $[\text{TolC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)]_2\text{O}$  **1**,  $[\text{PhC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)]_2\text{O}$  **2**,  $[(\text{C}_6\text{F}_5)\text{PhC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)]_2\text{O}$  **3**,  $[\text{Me}_2\text{BrCC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)]_2\text{O}$  **4** and  $[\text{MeC}(\text{O})\text{OB}(\text{C}_6\text{F}_5)]_2\text{O}$  **5**.** These species were prepared in a similar fashion and thus only one preparation is detailed. To a solution of *p*-toluic acid (168.7 mg, 1.24 mmol) in dichloromethane was added Lancaster's reagent,  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\text{SMe}_2$



(200 mg, 0.83 mmol). Bubbles were released immediately. After 40 min, the volatiles were removed under reduced pressure to give a white powder which was washed with 2 × 2 mL pentane and dried. Diffraction quality single crystals were obtained through slow diffusion of pentane into benzene at room temperature. **1**: Yield: 251.7 mg (95% isolated yield).  $^1\text{H}\{^{19}\text{F}\}$  NMR (400 MHz,  $\text{CDCl}_3$ , 253 K):  $\delta$  8.14 (d,  $J$  = 8.0 Hz, 4H, Ar), 7.35 (d,  $J$  = 8.0 Hz, 4H, Ar), 2.48 (s, 6H,  $\text{CH}_3$ ).  $^{19}\text{F}\{^1\text{H}\}$  NMR (377 MHz,  $\text{CDCl}_3$ ):  $\delta$  -134.5 (m,  $^3J_{\text{FF}}$  = 23.2 Hz, 4F, *o*- $\text{C}_6\text{F}_5$ ), -155.1 (t,  $^3J_{\text{FF}}$  = 20.3 Hz, 2F, *p*- $\text{C}_6\text{F}_5$ ), -163.4 (m,  $^3J_{\text{FF}}$  = 22.9 Hz, 4F, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz,  $\text{CDCl}_3$ ):  $\delta$  5.1 (br).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  176.3, 149.1 (dm,  $^1J_{\text{FC}}$  = 242 Hz), 148.9, 141.3 (dm,  $^1J_{\text{FC}}$  = 252 Hz), 137.6 (dm,  $^1J_{\text{FC}}$  = 262 Hz), 131.8, 130.1, 124.0, 22.3. The resonance for the *ipso*-B( $\text{C}_6\text{F}_5$ ) carbon was not observed. HRMS:  $[\text{C}_{28}\text{H}_{15}\text{B}_2\text{F}_{10}\text{O}_5]^+$  cal. 643.0946; exp. 643.0928. Elemental analysis: Calc.: C 52.38%, H 2.20%; Exp.: C 52.23%, H 1.76%.

**2**: Yield: 228.3 mg (90% isolated yield).  $^1\text{H}\{^{19}\text{F}\}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.27 (d,  $J$  = 7.9 Hz, 4H), 7.78 (t,  $J$  = 7.5 Hz, 2H), 7.57 (t,  $J$  = 7.7 Hz, 4H).  $^{19}\text{F}\{^1\text{H}\}$  NMR (377 MHz,  $\text{CDCl}_3$ ):  $\delta$  -134.5 (m,  $^3J_{\text{FF}}$  = 22.6 Hz, 4F, *o*- $\text{C}_6\text{F}_5$ ), -154.74 (t,  $^3J_{\text{FF}}$  = 20.3 Hz, 2F, *p*- $\text{C}_6\text{F}_5$ ), -163.2 (m,  $^3J_{\text{FF}}$  = 21.1 Hz, 4F, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz,  $\text{CDCl}_3$ ):  $\delta$  4.8.  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  176.4, 145.4 (dm,  $^1J_{\text{FC}}$  = 247 Hz), 141.3 (dm,  $^1J_{\text{FC}}$  = 258 Hz), 137.3 (dm,  $^1J_{\text{FC}}$  = 250 Hz), 137.0, 131.5, 129.2, 126.5. The resonance for the *ipso*-B( $\text{C}_6\text{F}_5$ ) carbon was not observed. Elemental analysis: Calc.: C 50.86%, H 1.64%; Exp.: C 50.78%, H 1.50%.

**3**: Yield: 295.0 mg (90% isolated yield).  $^{19}\text{F}\{^1\text{H}\}$  NMR (377 MHz,  $\text{CDCl}_3$ ):  $\delta$  -130.6 (br, 4F, *o*- $\text{C}_6\text{F}_5$ ), -134.31 (d,  $^3J_{\text{FF}}$  = 22.9 Hz, 4F, *o*- $\text{C}_6\text{F}_5$ ), -137.5 (br, 2F, *p*- $\text{C}_6\text{F}_5$ ), -153.1 (t,  $^3J_{\text{FF}}$  = 20.2 Hz, 2F, *p*- $\text{C}_6\text{F}_5$ ), -157.7 (br, 4F, *m*- $\text{C}_6\text{F}_5$ ), -162.68 (m,  $^3J_{\text{FF}}$  = 18.9 Hz, 4F, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz,  $\text{CDCl}_3$ ):  $\delta$  4.2. The  $^{13}\text{C}$  NMR spectrum was not observed due to the poor solubility of compound **3**. Elemental analysis: Calc.: C 39.34%; Exp.: C 38.70%.

**4**: Yield: 55.1 mg (88% isolated yield).  $^1\text{H}$  NMR (400 MHz, toluene- $d_8$ ):  $\delta$  1.61 (s, 12 H,  $\text{CH}_3$ ).  $^{19}\text{F}\{^1\text{H}\}$  NMR (377 MHz, toluene- $d_8$ ):  $\delta$  -134.4 (d,  $^3J_{\text{FF}}$  = 23.7 Hz, 4F, *o*- $\text{C}_6\text{F}_5$ ), -153.7 (t,  $^3J_{\text{FF}}$  = 20.2 Hz, 2F, *p*- $\text{C}_6\text{F}_5$ ), -162.7 (m,  $^3J_{\text{FF}}$  = 22.6 Hz, 4F, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz, toluene- $d_8$ ):  $\delta$  5.4.  $^{13}\text{C}$  NMR (126 MHz, toluene- $d_8$ ):  $\delta$  185.1, 149.1 (dm,  $^1J_{\text{FC}}$  = 246 Hz), 141.8 (dm,  $^1J_{\text{FC}}$  = 247 Hz), 137.7 (dm,  $^1J_{\text{FC}}$  = 249 Hz), 53.8, 28.32. The resonance for the *ipso*-B( $\text{C}_6\text{F}_5$ ) carbon was not observed. Elemental analysis: Calc.: C 34.14%, H 1.72%; Exp.: C 33.84%, H 1.55%.

**5**: Yield: 180.3 mg (89% isolated yield).  $^1\text{H}$  NMR (400 MHz, toluene- $d_8$ ):  $\delta$  1.45 (s, 6 H,  $\text{CH}_3$ ).  $^{19}\text{F}$  NMR (377 MHz, toluene- $d_8$ ):  $\delta$  -134.7 (d,  $^3J_{\text{FF}}$  = 16.2, 4F, *o*- $\text{C}_6\text{F}_5$ ), -154.7 (t,  $^3J_{\text{FF}}$  = 20.5 Hz, 2F, *p*- $\text{C}_6\text{F}_5$ ), -163.5 (m, 4F, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz, toluene- $d_8$ ):  $\delta$  4.2. The  $^{13}\text{C}$  NMR spectrum was not observed due to the poor solubility of compound **5**. Elemental analysis: Calc.: C 39.23%, H 1.23%; Exp.: C 39.02%, H 1.25%.

**Synthesis of  $[\text{ToIc}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2$  **6** and  $[(\text{C}_6\text{F}_5)_2\text{C}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2$  **7**.** These species were prepared in a similar fashion and thus only one preparation is detailed. To a solution of *p*-toluic acid (23.4 mg, 0.17 mmol) in dichloromethane, equi-

molar of Piers' borane,  $\text{HB}(\text{C}_6\text{F}_5)_2$  (60 mg, 0.17 mmol) was added. Hydrogen was released immediately. After 10 min, the solvent was removed under reduced pressure to give a white powder. Diffraction quality single crystals were obtained through slow diffusion of pentane into dichloromethane at  $-35^\circ\text{C}$ . **6**: Yield: 83.0 mg (>99%).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.98 (d,  $J$  = 8.4 Hz, 4H, Ar), 7.35 (d,  $J$  = 7.9 Hz, 4H, Ar), 2.49 (s, 6H,  $\text{CH}_3$ ).  $^{19}\text{F}$  NMR (377 MHz,  $\text{CDCl}_3$ ):  $\delta$  -135.1 (d,  $^3J_{\text{FF}}$  = 22.5 Hz, 8F, *o*- $\text{C}_6\text{F}_5$ ), -154.04 (t,  $^3J_{\text{FF}}$  = 19.3 Hz, 4F, *p*- $\text{C}_6\text{F}_5$ ), -162.1 (s, 8F, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz,  $\text{CDCl}_3$ ):  $\delta$  5.0 (br).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  173.9, 150.1, 147.9 (dm,  $^1J_{\text{FC}}$  ~240 Hz), 141.3 (dm,  $^1J_{\text{FC}}$  ~241 Hz), 137.5 (dm,  $^1J_{\text{FC}}$  ~232 Hz), 132.9, 130.5, 124.8, 112.8, 22.4. Elemental analysis: Calc.: C 50.04%, H 1.47%; Exp.: C 49.47%, H 1.40%.

**7**: Yield: 79 mg (83% isolated yield).  $^{19}\text{F}\{^1\text{H}\}$  NMR (470 MHz, toluene- $d_8$ ):  $\delta$  133.6 (4F, B-*o*- $\text{C}_6\text{F}_5$ ), -135.6, -150.8, -157.7 (2F, B-*p*- $\text{C}_6\text{F}_5$ ), -161.4 (4F, B-*m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (160 MHz, toluene- $d_8$ ):  $\delta$  1.5.  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  was not observed due to poor solubility of compound **7**. Elemental analysis: Calc.: 41.05%; Exp.: C 41.77%.

**Synthesis of  $[\text{Ph}_2\text{PO}_2\text{BH}(\text{C}_6\text{F}_5)]_2$  **8**.** To a suspension of diphenylphosphinic acid (45 mg, 0.21 mmol) in dichloromethane was added Lancaster's reagent,  $\text{H}_2\text{B}(\text{C}_6\text{F}_5)_2\text{SMe}_2$  (50.0 mg, 0.21 mmol). Bubbles were released immediately. After 4 h, the solvent was reduced under vacuum and a white solid was recrystallized from dichloromethane and pentane in a ratio of 1 : 0.5. Diffraction quality single crystals were obtained from a saturated dichloromethane solution at  $-35^\circ\text{C}$ . Yield: 81.6 mg (50% isolated yield).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.81–7.45 (ov, Ar), 4.0 (br, B-H).  $^{19}\text{F}$  NMR (377 MHz,  $\text{CDCl}_3$ ):  $\delta$  -135.4 (d,  $^3J_{\text{FF}}$  = 24.3 Hz, *o*- $\text{C}_6\text{F}_5$ ), -135.7 (d,  $^3J_{\text{FF}}$  = 23.7 Hz, *o*- $\text{C}_6\text{F}_5$ ), -158.0 (t,  $^3J_{\text{FF}}$  = 20.3 Hz, *p*- $\text{C}_6\text{F}_5$ ), -158.1 (t,  $^3J_{\text{FF}}$  = 20.3 Hz, *p*- $\text{C}_6\text{F}_5$ ), -164.1 (m, *m*- $\text{C}_6\text{F}_5$ ), -164.3 (m, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.3 (br).  $^{31}\text{P}$  NMR (162 MHz,  $\text{CDCl}_3$ , 298 K):  $\delta$  40.8, 37.6 (1 : 0.44).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  133.4 (ov), 133.3, 132.3, 132.3, 132.0, 131.9, 131.5, 131.4, 129.1, 129.0, 128.89, 128.86, 128.8, 128.4, 127.2. Elemental analysis: Calc.: C 54.59%, H 2.80%; Exp.: C 54.01%, H 2.64%.

**Synthesis of  $\text{ToIc}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2[\text{NC}_5\text{H}_4\text{NMe}_2]$  **9**.** To a solution of dimer  $[\text{ToIc}(\text{O})\text{OB}(\text{C}_6\text{F}_5)_2]_2$  (40 mg, 0.04 mmol) in dichloromethane, 4-dimethylaminopyridine (10 mg, 0.08 mmol) was added. The solvent was removed and a white powder was obtained; yield: 49.6 mg (99%). Crystals suitable for single crystal X-ray diffraction were obtained from layering of dichloromethane and pentane.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.22 (d,  $J$  = 7.1 Hz, 2H, Ar), 7.95 (d,  $J$  = 8.2 Hz, 2H, Ar), 7.21 (d,  $J$  = 7.9 Hz, 2H, Ar), 6.62 (d,  $J$  = 7.8 Hz, 2H, Ar), 3.17 (s, 6H,  $\text{CH}_3$ ), 2.39 (s, 3H,  $\text{CH}_3$ ).  $^{19}\text{F}$  NMR (470 MHz,  $\text{CDCl}_3$ ):  $\delta$  -133.5 (d,  $^3J_{\text{FF}}$  = 21.1 Hz, 4F, *o*- $\text{C}_6\text{F}_5$ ), -157.3 (t,  $^3J_{\text{FF}}$  = 20.4 Hz, 4F, *p*- $\text{C}_6\text{F}_5$ ), -163.6 (m, 4F, *m*- $\text{C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (160 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.6.  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  168.0, 156.1, 148.1 (dm,  $^1J_{\text{FC}}$  ~242 Hz), 144.1, 142.8, 140.3 (dm,  $^1J_{\text{FC}}$  ~250 Hz), 137.3 (dm,  $^1J_{\text{FC}}$  ~249 Hz), 130.5, 130.0, 129.0, 106.5, 39.8, 21.7. The resonance for the *ipso*-B( $\text{C}_6\text{F}_5$ ) carbon was not observed. Elemental analysis: Calc.: C 53.85%, H 2.85%; Exp.: C 53.67%, H 2.78%.



**Synthesis of  $\text{Ph}_2\text{PO}_2\text{BH}(\text{C}_6\text{F}_5)(\text{NC}_5\text{H}_4\text{NMe}_2)$  10.** To a suspension of  $[\text{Ph}_2\text{PO}_2\text{BH}(\text{C}_6\text{F}_5)]_2$  (45 mg, 0.06 mmol) in dichloromethane 4-dimethylaminopyridine (14 mg, 0.12 mmol) was added. After 10 min, the solvent was reduced under vacuum. The product was recrystallised from a solution of dichloromethane at  $-35^\circ\text{C}$ . Yield: 50 mg (85% isolated yield).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ): 8.20 (d,  $J = 7.2$  Hz, 2H), 7.87 (m, 2H), 7.73 (m, 2H), 7.42–7.30 (m, 6H), 6.54 (d,  $J = 7.6$  Hz, 2H), 3.10 (s, 6H). The B–H was not observed.  $^{19}\text{F}$  NMR (377 MHz,  $\text{CDCl}_3$ ):  $\delta$   $-134.9$  (m,  $o\text{-C}_6\text{F}_5$ ),  $-158.1$  (m,  $p\text{-C}_6\text{F}_5$ ),  $-164.0$  (m,  $m\text{-C}_6\text{F}_5$ ).  $^{11}\text{B}$  NMR (128 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.8.  $^{31}\text{P}$  NMR (162 MHz,  $\text{CDCl}_3$ ):  $\delta$  25.3.  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  156.1, 148.3 (dm,  $^1J_{\text{FC}} = 234$  Hz), 144.3, 140.0 (dm,  $^1J_{\text{FC}} = 249$  Hz), 137.0 (dm,  $^1J_{\text{FC}} = 266$  Hz), 135.7 (d,  $^1J_{\text{PC}} = 106.1$  Hz), 134.57 (d,  $^1J_{\text{PC}} = 96.5$  Hz), 131.78 (d,  $^3J_{\text{PC}} = 10.0$  Hz), 131.55 (d,  $^3J_{\text{PC}} = 9.8$  Hz), 131.04 (d,  $^4J_{\text{PC}} = 2.8$  Hz), 131.01 (d,  $^4J_{\text{PC}} = 2.8$  Hz), 128.22 (d,  $^2J_{\text{PC}} = 13.0$  Hz), 128.05 (d,  $^2J_{\text{PC}} = 12.8$  Hz), 106.7, 39.7. The resonance for the *ipso*-B( $\text{C}_6\text{F}_5$ ) carbon was likely not observed. Elemental analysis: Cal.: C 57.94%, H 4.08%; Exp. C 56.99%, H 3.89%.

## Computational details

Electronic structure calculations were performed using Gaussian 16.<sup>33</sup> Geometry optimisations, frequency calculations, and energy determinations were performed using the M062X<sup>34</sup> functional and the def2-TZVPP<sup>35</sup> basis set with the D3 version of Grimme's dispersion (GD3)<sup>36,37</sup> and the dichloromethane solvation effect calculated using the polarisable continuum model (PCM). The absence of any imaginary frequency with an absolute magnitude greater than  $10\text{ cm}^{-1}$  confirmed that each optimised structure was indeed located at a minimum on its potential energy hypersurface. The Gibbs energy corrections from frequency calculations were added to the single-point energies to obtain the Gibbs free energies in solution. Natural bond orbital and natural population analyses were performed on the optimized structures using the M062X functional and def2-TZVP<sup>35</sup> basis set using NBO 6.0.<sup>38</sup> This work was made possible by the facilities of the Shared Hierarchical Academic Research Computing Network (SHARCNET: <http://www.sharcnet.ca>) and Compute Canada.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

D. W. S. gratefully acknowledges the financial support from the NSERC, Canada, and the award of Canada Research Chair. D. W. S. is also grateful for the award of an Einstein Fellowship at TU Berlin. RLM is grateful for the award of an EPSRC fellowship (CEP R026912/1).

## Notes and references

- R. G. Hayter, A. W. Laubengayer and P. G. Thompson, *J. Chem. Soc.*, 1957, 4243–4244.
- W. Gerrard, M. F. Lappert and R. Shafferman, *J. Chem. Soc.*, 1958, 3648–3652.
- A. D. Negro, L. Ungaretti and A. Perotti, *J. Chem. Soc., Dalton Trans.*, 1972, 1639–1643.
- A. Sporzyński, *J. Organomet. Chem.*, 1993, **448**, C1–C2.
- R. Köster, A. Sporzyński, W. Schuessler, D. Blaeser and R. Boese, *Chem. Ber.*, 1994, **127**, 1191–1199.
- B. Wrackmeyer, E. Khan and R. Kempe, *Z. Naturforsch., B: Chem. Sci.*, 2008, **63**, 275–279.
- M. Periasamy, M. N. Reddy and N. S. Kumar, *Sci. Synth.*, 2005, **6**, 321–335.
- P. Starkov and T. D. Sheppard, *Org. Biomol. Chem.*, 2011, **9**, 1320–1323.
- R. M. Lanigan, P. Starkov and T. D. Sheppard, *J. Org. Chem.*, 2013, **78**, 4512–4523.
- V. Karaluka, R. M. Lanigan, P. M. Murray, M. Badland and T. D. Sheppard, *Org. Biomol. Chem.*, 2015, **13**, 10888–10894.
- P. W. Tang, *Org. Synth.*, 2005, **81**, 262–272.
- K. Ishihara, S. Ohara and H. Yamamoto, *J. Org. Chem.*, 1996, **61**, 4196–4197.
- K. Ishihara, S. Ohara and H. Yamamoto, *Macromolecules*, 2000, **33**, 3511–3513.
- T. Maki, K. Ishihara and H. Yamamoto, *Synlett*, 2004, 1355–1358.
- A. Sakakura, T. Ohkubo, R. Yamashita, M. Akakura and K. Ishihara, *Org. Lett.*, 2011, **13**, 892–895.
- R. Yamashita, A. Sakakura and K. Ishihara, *Org. Lett.*, 2013, **15**, 3654–3657.
- K. Ishihara and Y. H. Lu, *Chem. Sci.*, 2016, **7**, 1276–1280.
- K. Arnold, B. Davies, D. Herault and A. Whiting, *Angew. Chem., Int. Ed.*, 2008, **47**, 2673–2676.
- K. Arnold, A. S. Batsanov, B. Davies and A. Whiting, *Green Chem.*, 2008, **10**, 124–134.
- R. M. Al-Zoubi, O. Marion and D. G. Hall, *Angew. Chem., Int. Ed.*, 2008, **47**, 2876–2879.
- N. Gernigon, R. M. Al-Zoubi and D. G. Hall, *J. Org. Chem.*, 2012, **77**, 8386–8400.
- T. M. El Dine, J. Rouden and J. Blanchet, *Chem. Commun.*, 2015, **51**, 16084–16087.
- S. Arkhipenko, M. T. Sabatini, A. S. Batsanov, V. Karaluka, T. D. Sheppard, H. S. Rzepa and A. Whiting, *Chem. Sci.*, 2018, **9**, 1058–1072.
- K. Wang, Y. Lu and K. Ishihara, *Chem. Commun.*, 2018, **54**, 5410–5413.
- N. M. Yoon, C. S. Pak, H. C. Brown, S. Krishnam and T. P. Stocky, *J. Org. Chem.*, 1973, **38**, 2786–2792.
- H. C. Brown and T. P. Stocky, *J. Am. Chem. Soc.*, 1977, **99**, 8218–8226.
- L. L. Liu, L. L. Cao, Y. Shao and D. W. Stephan, *J. Am. Chem. Soc.*, 2017, **139**, 10062–10071.
- L. L. Cao and D. W. Stephan, *Organometallics*, 2017, **36**, 3163–3170.



- 29 A. M. Fuller, D. L. Hughes, S. J. Lancaster and C. M. White, *Organometallics*, 2010, **29**, 2194–2197.
- 30 D. J. Parks, R. E. v. H. Spence and W. E. Piers, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 809–811.
- 31 I. Justyniak, D. Prochowicz, A. Tulewicz, W. Bury, P. Gos and J. Lewinski, *Dalton Trans.*, 2017, **46**, 669–677.
- 32 R. Kather, E. Rychagova, P. Sanz Camacho, S. E. Ashbrook, J. D. Woollins, L. Robben, E. Lork, S. Ketkov and J. Beckmann, *Chem. Commun.*, 2016, **52**, 10992–10995.
- 33 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, M. C. X. Li, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. J. A. Montgomery, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 09, Revision E.01*, Wallingford CT, 2016.
- 34 Y. Zhao and D. Truhlar, *Theor. Chem. Acc.*, 2008, **120**, 215–241.
- 35 F. Weigend, *Phys. Chem. Chem. Phys.*, 2006, **8**, 1057–1065.
- 36 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 37 S. Grimme, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 38 E. D. Glendening, J. K. Badenhoop, A. E. Reed, J. E. Carpenter, J. A. Bohmann, C. M. Morales, C. R. Landis and F. Weinhold, *NBO 6.0*, Madison, WI, 2013.

