

Featuring work from Dr. Peng Zhang and Prof. Zhiyong Zhang at Key Laboratory for Biological Effects of Nanomaterials and Nanosafety, Institute of High Energy Physics, Chinese Academy of Sciences, China.

#### Plant species-dependent transformation and translocation of ceria nanoparticles

Uptake of nanoparticles by plants is species-dependent, the underlying mechanisms of which are explored here. Root exudate composition and xylem structure are found to be both important factors in determining the transformation and translocation of ceria nanoparticles in plants.

#### As featured in:



See Peng Zhang, Zhiyong Zhang et al., *Environ. Sci.: Nano*, 2019, 6, 60.



ROYAL SOCIETY  
OF CHEMISTRY

Celebrating  
IYPT 2019

[rsc.li/es-nano](http://rsc.li/es-nano)

Registered charity number: 207890



Cite this: *Environ. Sci.: Nano*, 2019, 6, 60

Received 27th September 2018,  
Accepted 9th November 2018

DOI: 10.1039/c8en01089g

rsc.li/es-nano

## Plant species-dependent transformation and translocation of ceria nanoparticles†

Peng Zhang, <sup>\*ab</sup> Yuhui Ma, <sup>a</sup> Changjian Xie, <sup>a</sup> Zhiling Guo, <sup>b</sup> Xiao He, <sup>a</sup> Eugenia Valsami-Jones, <sup>b</sup> Iseult Lynch, <sup>b</sup> Wenhe Luo, <sup>a</sup> Lirong Zheng <sup>c</sup> and Zhiyong Zhang <sup>\*a</sup>

Uptake of nanoparticles (NPs) by plants is species-dependent; however, its underlying mechanisms are rarely investigated. This study compared the transformation, uptake and translocation of CeO<sub>2</sub> NPs (nCeO<sub>2</sub>) in four different plant species. Corn showed the lowest translocation factor (TF) of Ce, while the differences among the plant species reduced or diminished when the phosphates were removed from the nutrient solution (–P treatment). Using transmission electron microscopy (TEM) and synchrotron-based X-ray absorption near edge spectroscopy (XANES), we found that the removal of phosphate from the nutrient solutions reduces or eliminates the immobilization of Ce<sup>3+</sup> on the root surface thus promoting the upward translocation of Ce from roots to shoots, which accounts for the enhanced total Ce detected in all the four plants. Compared with the +P treatment, the Ce(III) contents in the shoots of corn and wheat were enhanced by 27.4 and 4.0 times in the –P treatment, which may account for the reduced biomass of shoots in these two plant species. Our results indicate that different plant species have different abilities of transforming and translocating CeO<sub>2</sub> NPs and that the mechanisms differ depending on the chemistry of the surrounding solution (+P/–P). Plant xylems and root exudate compositions are both important factors in determining the transformation of nCeO<sub>2</sub> and subsequent translocation of Ce species in plants. Further studies are required to determine the composition of root exudates and to identify the components driving the nCeO<sub>2</sub> transformation in different plant species under different culturing conditions.

### Environmental significance

Transformation of nanomaterials (NMs) in plants is a critical process that may determine the final fate of NMs in the environment. We previously found that root surface is the major site for nCeO<sub>2</sub> transformation. In this study, we compared the uptake and phytotoxicity of nCeO<sub>2</sub> to two dicots and two monocots. Different plant species show different capacities to transform nCeO<sub>2</sub> into Ce(III) species, which results in different translocation of Ce in plants. Removal of phosphate from the nutrient solution reduces or eliminates the immobilization of Ce<sup>3+</sup> on the root surface thus promoting the upward translocation of Ce, which leads to the phytotoxicity found in the monocots. Plant xylems and root exudate compositions are both important factors in determining the transformation of nCeO<sub>2</sub> and the subsequent translocation of Ce species in plants. This study reveals the underlying mechanism behind the plant species-dependent uptake and phytotoxicity of NMs in plants. Further studies are required to determine the composition of root exudates in different plant species and identify the determining components for nCeO<sub>2</sub> transformation under different culturing conditions.

## 1. Introduction

CeO<sub>2</sub> minerals were previously recognized as being highly stable in the environment; however, an increasing body of studies now suggest that CeO<sub>2</sub> in its nano form is prone to transform in various environmental surroundings including aquatic mesocosms,<sup>1</sup> wastewater treatment plants,<sup>2</sup> activated sludge,<sup>3</sup> synthetic river water,<sup>4</sup> bacteria,<sup>5</sup> plants,<sup>6</sup> etc., and various transformation products were reported depending on the ambient environment chemistry and biochemistry. For example, in an activated sludge where the sulphur concentration is high, Ce<sup>3+</sup> ions might be released from nCeO<sub>2</sub> upon bacterial reduction and complexation, forming Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> in oxidizing environments or forming Ce<sub>2</sub>S<sub>3</sub> in reducing environments.<sup>3</sup> In plants, low molecular organic acids and reducing substances excreted by plant roots can facilitate the release of Ce<sup>3+</sup> ions from nCeO<sub>2</sub> in the rhizosphere vicinity, which then can be translocated from roots to shoots and form either CePO<sub>4</sub> or Ce carboxylates.<sup>6–8</sup> Retransformation of Ce<sup>3+</sup> ions released from nCeO<sub>2</sub> back to CeO<sub>2</sub> is also

<sup>a</sup> Key Laboratory for Biological Effects of Nanomaterials and Nanosafety, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China. E-mail: pengzhang@ihep.ac.cn, zhangzhy@ihep.ac.cn; Fax: +86 10 88235294; Tel: +86 10 88233215

<sup>b</sup> School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, UK. E-mail: p.zhang.1@bham.ac.uk

<sup>c</sup> Beijing Synchrotron Radiation Facility, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

† Electronic supplementary information (ESI) available: Characterization results of nCeO<sub>2</sub> (Fig. S1 and Table S1), biomass of plants (Fig. S2), TEM figures of root sections (Fig. S3–S6), fraction of Ce species obtained by LCF analysis (Tables S2 and S3), Ce species contents (Table S4), translocation factors of Ce species in plants (Fig. S7), and DLS results of nCeO<sub>2</sub> in root exudates (Fig. S8 and S9 and Table S5). See DOI: 10.1039/c8en01089g



possible.<sup>3,9</sup> Transformation processes such as carboxylation/complexation facilitate the release and mobility of Ce(III) thus increasing the toxicity, while sulfidation or phosphorylation reduce the mobility and toxicity of NPs.<sup>10</sup> However, redissolution of CePO<sub>4</sub> is also possible in the presence of strong Ce<sup>3+</sup> ion-binding ligands in the root exudates.<sup>11</sup>

nCeO<sub>2</sub> transformation is highly associated with the physicochemical characteristics of nCeO<sub>2</sub> such as surface coating,<sup>3</sup> size,<sup>12</sup> and shape.<sup>13</sup> For example, surface coating with citric acid may act as a shield reducing the direct contact of the particle surface with the ambient environment and subsequent dissolution.<sup>3</sup> Compared with larger size nCeO<sub>2</sub>, smaller size nCeO<sub>2</sub> released more Ce(III) ions and thus induced higher toxicity to *Lactuca* plants.<sup>12</sup> Rod-shaped nCeO<sub>2</sub> with highly reactive crystal facets transformed to a greater extent in cucumber plants than other shapes of nCeO<sub>2</sub> such as octahedra and truncated octahedra.<sup>13</sup>

Another important factor, *i.e.* the plant species, has been largely ignored in the study of the uptake and transformation of nCeO<sub>2</sub> in plants. Previous studies have suggested that the uptake and translocation of NPs are highly dependent on plant species.<sup>9,14,15</sup> The discrepancy might not be attributed only to the structural differences but also to the composition difference of the root exudates, considering that the root surface is the major site for the transformation of nCeO<sub>2</sub> in plants.<sup>16</sup>

In this study, we exposed two monocots (corn and wheat) and two dicots (cabbage and soybean) to nCeO<sub>2</sub> (2000 mg L<sup>-1</sup>) under two different hydroponic conditions (with and without phosphates in the nutrient solution, called “+P” and “-P” treatment, respectively), with the aim to understand how plant species and culturing conditions impact the transformation, uptake and translocation of nCeO<sub>2</sub> by plants. To supply P for plant growth in the “-P” treatment, 1 mM KH<sub>2</sub>PO<sub>4</sub> was sprayed onto the foliage of the plants every two days. Translocation and transformation of nCeO<sub>2</sub> in plants were then examined by ICP-MS, TEM and X-ray absorption near edge spectroscopy (XANES).

## 2. Materials and methods

### 2.1. Materials, plant cultivation and exposure

nCeO<sub>2</sub> with an average size of 25 nm (Fig. S1†) were synthesized by a precipitation method<sup>13</sup> and the physicochemical properties of nCeO<sub>2</sub> are described in Table S1.† Seeds of corn, wheat, cabbage and soybean purchased from the Chinese Academy of Agricultural Sciences were sterilized in a 5% NaClO solution and rinsed with deionized water. Then the seeds were placed on filter paper which was soaked with deionized water in a plastic Petri dish and allowed for germination in the dark for 5 days. The seedlings were then transferred into 200 mL beakers containing 100 mL  $\frac{1}{4}$  strength Hoagland solution. In the -P treatment, nutrient solution without phosphate was used. To supply P for normal growth of plant, KH<sub>2</sub>PO<sub>4</sub> (1 mM) solution was sprayed on the foliage of the plants every two days. The seedlings were allowed to

grow for 10 days in an artificial climate chamber, and then exposed to 2000 mg L<sup>-1</sup> nCeO<sub>2</sub> suspension in  $\frac{1}{4}$  strength Hoagland solution; 2000 mg L<sup>-1</sup> was the maximum exposure concentration used in the phytotoxicity test established by the U.S. Environmental Protection Agency guideline.<sup>17</sup> After 2 weeks of exposure, the seedlings were harvested, rinsed with deionized water and lyophilized for further analysis.

### 2.2. Phytotoxicity, Ce localization, quantification and chemical species analysis

The dry weights of the shoots and roots of the plants were measured for assessing the phytotoxicity. The root apices were collected, fixed, dehydrated, embedded, and sectioned following the procedures for TEM observation as described previously.<sup>6</sup> A JEM-1230 (JEOL) transmission electron microscope was used for observing the sections. The shoots and roots were ground into fine powders using a pestle and mortar. The powders were then digested with a mixture of nitric acids and hydrogen peroxide (v/v: 4/1), and the total Ce contents were analysed by ICP-MS (Thermo).

For XANES analyses, the fine powders of the shoots and roots were compressed into thin slices (~2 mm). The Ce L-III edge spectra were collected using a 19-element high purity Ge solid state detector array at beamline 1W1B of the Beijing Synchrotron Radiation Facility. The XANES spectra of nCeO<sub>2</sub> and Ce-containing compounds including CePO<sub>4</sub>, Ce acetate and Ce oxalate were collected for reference. Linear combination fitting (LCF) of the XANES data was performed on Athena software. LCF is justified because the X-ray adsorption of different species in the sample is additive.<sup>18</sup> The total absorption coefficient calculated by LCF can be written as

$$M = \sum f_i(\text{STD}_i),$$

where  $M$  is the least squares fit to the sample spectrum, (STD<sub>*i*</sub>) represents the absorption coefficient of the standard reference spectra, and  $f_i$  is the fraction of each reference spectrum in the sample spectrum, summed over the number of references,  $i$ . Thus, with appropriate standard references, LCF can identify and quantify the fraction of chemical species in an unknown sample.

## 3. Results and discussion

### 3.1. Plant species-dependent Ce uptake – influence of phosphate

Fig. S2† shows that the four plant species respond differently to nCeO<sub>2</sub> exposure. In the +P treatment, nCeO<sub>2</sub> increased the root biomass of soybean by 36%, but had no effect on the other plant species tested. In the -P treatment, the positive effects of nCeO<sub>2</sub> on the soybean roots were remarkably enhanced, with the root biomass increased by 60% compared with that of the control, while nCeO<sub>2</sub> still had no effect on the other plant species tested. nCeO<sub>2</sub> had no significant effects on the shoot biomass of soybean and cabbage in the -P treatment; however, nCeO<sub>2</sub> reduced the shoot biomass of





corn and wheat by 24% and 63%, respectively, in the -P treatment. These results suggest that the phytotoxicity of nCeO<sub>2</sub> is plant species-dependent and is influenced by the composition of the nutrient solution used in the hydroponic growth system.

Table 1 shows that the Ce uptake and translocation in the plants were also highly species-dependent. In the +P treatment, the Ce content in the plants and the TFs of the monocots (corn and wheat) were lower than those of the dicots (cabbage and soybean) except for wheat where the Ce content in the root is higher than those in the other plant species. This might be due to the fact that cabbage and soybean have larger leaf surface areas resulting in higher transpiration rates, because the driving force for long distance translocation of nutrients mainly depends on the transpiration rate.<sup>19</sup>

The removal of phosphate (P) from the nutrient solution remarkably promoted the uptake of Ce into the roots of corn and cabbage and the upward translocation of Ce in all the four species. The Ce contents in the roots of corn and cabbage exposed under the -P treatment condition were 5.5 and 1.9 times higher, respectively, than those under the +P treatment. The Ce contents in the shoots of all the plant species were higher under the -P treatment than those under the +P treatment. In particular, the Ce contents in the shoots of corn, wheat, cabbage and soybean under the -P treatment were 28.4, 5.0, 2.3, and 2.4 times higher, respectively, than those under the +P treatment. Obviously, P has greater effects on the translocation of Ce in the monocots (corn and wheat) than in the dicots (cabbage and soybean). This indicates that the accumulations of Ce in the shoots of all the plant species under the -P treatment are equivalent.

### 3.2. Ce uptake and transformation by TEM

To further understand the above results, we observed the root sections by TEM (Fig. S3–S6†). In all the plant species, we found needle or rod-like particles on the root surfaces in the +P treatment, which are completely different from the morphology of the pristine octahedral nCeO<sub>2</sub>; however, in the -P treatment, we found large amounts of flocculated particles in the vacuole of root cells while no needle or rod-like particles were found on the root surface. The needle, rod, or the flocculated particles were CePO<sub>4</sub> which have been identified in previous studies.<sup>6,20</sup> Given that the transformation of CeO<sub>2</sub> mainly occurs on the root surface rather than inside of the

plant,<sup>16</sup> the Ce<sup>3+</sup> ions released from CeO<sub>2</sub> can be easily immobilized on the root surface in the presence of P in the nutrient solution, and the upward translocation of Ce will be then inhibited. The removal of P from the medium reduces or eliminates the immobilization of Ce on the root surface, and thus the free translocation of Ce<sup>3+</sup> from the roots to shoots can occur. The formation of CePO<sub>4</sub> in the vacuoles in the -P treatment was because foliar applied P was adsorbed by plants and stored in the vacuole as a P “pool” which then reacts with the free Ce<sup>3+</sup> ions as they translocate into the plant.<sup>21</sup> The above results suggest that the P-containing ions have a significant impact on the uptake of nCeO<sub>2</sub> in all the plant species tested.

It should be noted that the particulate CeO<sub>2</sub> NPs were also observed inside the root of soybean but were not observed in the other plants (Fig. 1 and S5B†). As shown in Fig. 1A and C, nCeO<sub>2</sub> were found in the intercellular regions. Interestingly, the morphology of some particles in the large aggregates (Fig. 1B) was different from that of the original nCeO<sub>2</sub>, suggesting the transformation of nCeO<sub>2</sub>. We previously found that nCeO<sub>2</sub>, which have been translocated into the cucumber plant, cannot transform;<sup>16</sup> however, a recent study shows that nCeO<sub>2</sub> may also be dissolved inside the leaves of sunflower resulting from pH variation and interaction with antioxidant substances in apoplast.<sup>22</sup> The nCeO<sub>2</sub> that we found here may be transformed either before or after their entering into the plants.


### 3.3. Transformation of nCeO<sub>2</sub> in plants analyzed by XANES

To further understand the translocation of Ce in the four plant species, we quantified the chemical species of Ce in the plant tissues by XANES and linear combination fitting (LCF) analyses (complete LCF results are given in the ESI† Tables S2 and S3). The XANES spectra show how X-rays are absorbed by a specific atom, and how the absorption is modified by the chemical and physical environment of the atom in different compounds. With appropriate reference compounds, quantification of different chemical species can be achieved by doing the linear combination fitting (LCF) analysis on the XANES spectra. The results show that Ce was mainly presented as CeO<sub>2</sub> (>74.1%) in the roots of the four plant species in both +P and -P treatments (Fig. 2A and 3A and B); the Ce(III) species are mainly CePO<sub>4</sub> in the +P treatment but are mainly Ce carboxylates in the -P treatment. This is in

**Table 1** Ce content (mg kg<sup>-1</sup> dry weight) in the roots and shoots of the plants treated with nCeO<sub>2</sub>, and translocation factors (TFs, defined by dividing the Ce contents in the shoots by that in the roots) of Ce in the different plant species. Data are expressed as mean ± SD (*n* = 8). \* indicates a significant difference between +P and -P treatment for each plant species (*p* < 0.05). Significant differences between different species in each treatment are compared separately and expressed as lowercase letters. Different superscript letters indicate a significant difference (*p* < 0.05)

	Root		Shoot		TFs (×10 <sup>-3</sup> )	
	-P	+P	-P	+P	-P	+P
Corn	68 093 ± 20 530 <sup>a*</sup>	12 499 ± 2600 <sup>b</sup>	158 ± 56 <sup>a*</sup>	5.6 ± 1.6 <sup>c</sup>	2.3 ± 0.8 <sup>b*</sup>	0.5 ± 0.1 <sup>c</sup>
Wheat	39 269 ± 9278 <sup>b</sup>	34 322 ± 10 207 <sup>a</sup>	140 ± 45 <sup>a*</sup>	28 ± 11 <sup>b</sup>	3.6 ± 1.0 <sup>b*</sup>	0.8 ± 0.3 <sup>b</sup>
Cabbage	58 315 ± 22 799 <sup>ab</sup>	31 418 ± 14 821 <sup>a</sup>	175 ± 44 <sup>a*</sup>	76 ± 15 <sup>a</sup>	3.0 ± 0.7 <sup>b</sup>	2.4 ± 0.4 <sup>a</sup>
Soybean	35 003 ± 6615 <sup>b</sup>	31 527 ± 5669 <sup>a</sup>	181 ± 37 <sup>a*</sup>	77 ± 9 <sup>a</sup>	5.2 ± 0.5 <sup>a*</sup>	2.4 ± 0.2 <sup>a</sup>






**Fig. 1** TEM images of the root sections of soybean treated with  $n\text{CeO}_2$  under the  $-P$  condition. (A) and (C) show the intercellular regions. (B) and (D) are the magnified images of the highlighted rectangle areas in A and C. Red arrows in B indicate the  $n\text{CeO}_2$  that were morphologically “transformed”.

accordance with the TEM results that the  $\text{Ce}^{3+}$  ions were mainly immobilized on the root surface with a minimum amount being translocated into the roots and shoots in the  $+P$  treatment; however, in the  $-P$  treatment, the  $\text{Ce}^{3+}$  ions moved easily into the roots, bound to  $\text{PO}_4^{3-}$  in the vacuoles

of root cells, and bound to carboxyl groups in the xylem during the upward translocation.

The Ce species in the shoots were different from that in the roots (Fig. 2B and 3C and D). In the  $+P$  treatment,  $\text{CeO}_2$  was the predominant species in the shoots, with the highest proportion of 94.7% in corn and the lowest of 78.2% in wheat; only a small amount of Ce presented as Ce carboxylates. However, in the  $-P$  treatment, the proportion of Ce carboxylates remarkably increased, with a proportion of 93.3% in corn and the lowest proportion of 43.3% in the wheat shoots. Phosphate deficiency can induce a high abundance of metabolites with chelating ability and increase the active exudation of organic acids out from the roots.<sup>23</sup> This process will enhance the dissolution of  $n\text{CeO}_2$ . We further calculated the absolute contents of  $\text{Ce(IV)}$  and  $\text{Ce(III)}$  in the roots and shoots by multiplying their proportion by the total Ce contents (Table S4†). The Ce (IV) and Ce(III) contents in the roots and shoots in the  $-P$  treatment were both higher than those in the  $+P$  treatment (consistent with the data in Table 1). Notably, the Ce(III) contents in the shoots of corn in the  $-P$  treatment were 500 times higher than those in the  $+P$  treatment. Ce(III) is more toxic to plants than Ce in its  $n\text{CeO}_2$  form,<sup>24</sup> therefore, the enhanced Ce(III) accumulation in the shoots might be the reason leading to the decrease of shoot biomasses of the corn and wheat plants under the  $-P$  treatment.

The mechanism of the transformation and translocation of  $n\text{CeO}_2$  in the cucumber plant has been well discussed in previous studies.<sup>6,16</sup> However, discrepancies of the transformation and translocation of  $n\text{CeO}_2$  in different plant species are not fully explored. We deduced that two factors may account for the species-dependent translocation of the  $\text{CeO}_2$  NPs found in this study (Fig. 4). Firstly, structural differences between monocots and dicots result in different translocation abilities of substances. The xylem walls of dicots usually have higher affinity for nutrient ions than those of monocots because there are more negatively charged groups (*e.g.* carboxyl



**Fig. 2** Ce L-III edge XANES spectra of the root (A) and shoot (B) of the plants with treatment of  $2000 \text{ mg L}^{-1} n\text{CeO}_2$  for 2 weeks under the  $+P$  (solid lines) and  $-P$  (dotted lines) conditions. Vertically dotted line indicates the peak of the Ce(III) species centered at 5726.7 eV.






Fig. 3 Fraction of the Ce species in the four types of plants treated with 2000 mg L<sup>-1</sup> nCeO<sub>2</sub> for 2 weeks. (A) Root, +P, (B) root, -P, (C) shoot, +P, and (D) shoot, -P.




Fig. 4 A schematic illustration of the transformation of nCeO<sub>2</sub> on the root surface and the translocation of nCeO<sub>2</sub> and Ce<sup>3+</sup> ions in the root xylem of the dicot and monocot using soybean and corn as examples.



groups) on the xylem walls of dicots than those of monocots.<sup>25</sup> Therefore, the translocation of cations in dicots is usually more difficult than in monocots, as the cations become immobilised on the xylem walls. However, this is not consistent with the higher Ce contents found in the dicots in the +P treatment. This can be explained by the different transformation under the two culturing conditions. In the +P treatment, Ce is mainly translocated in its particulate form (Table S4†), which is mainly driven by transpiration and is not affected by xylem walls/ion charges. However, in the -P treatment, more free  $\text{Ce}^{3+}$  ions entered into the roots (Table S4†) and can easily bind to the negatively charged groups (carboxyl) on the xylem wall; structural differences will then have significant effects on the translocation of Ce. The xylems of dicots are more negatively charged than the xylems of monocots, thus can bind more  $\text{Ce}^{3+}$  ions and restricted the upward translocation of the  $\text{Ce}^{3+}$  ions. Different translocation patterns of  $\text{nCeO}_2$  and  $\text{Ce}^{3+}$  in plant tissues are also reported in a previous study which compared the two-dimensional distribution of Ce in cucumber leaves with  $\text{nCeO}_2$  and  $\text{Ce}^{3+}$  ion treatments.<sup>26</sup> This suggests that the particulate  $\text{CeO}_2$  mostly move to the leaf margins along with the water flow within the vascular bundle which is driven by transpiration; however, the  $\text{Ce}^{3+}$  ions mostly accumulate along the leaf veins, suggesting the binding of  $\text{Ce}^{3+}$  to the negatively charged groups on the xylem wall. The xylem structure-dependent translocation is further evidenced by looking at the translocation factor of the  $\text{Ce(IV)}$  and  $\text{Ce(III)}$  species in the plants (Fig. S7†). The TFs of  $\text{Ce(IV)}$  in the dicots are higher than those in the monocots, and remain largely the same between the +P and -P treatments for each plant species, suggesting a transpiration-driven translocation process of particulate Ce. However, the TFs of  $\text{Ce(III)}$  change drastically between the +P and -P treatments especially for the dicots. In the corn plants, the TFs of  $\text{Ce(III)}$  in the -P treatment are 221 times higher than that in the +P treatment, which leads to the accumulation of enough  $\text{Ce}^{3+}$  inducing the toxicity. These results indicate that the translocation of positively charged  $\text{Ce}^{3+}$  depends on the xylem structures.

Secondly, root exudates play a critical role in the transformation of  $\text{nCeO}_2$  in plants. We extracted the root exudates from the plants according to a previously described method,<sup>27</sup> and found that the hydrodynamic sizes of  $\text{nCeO}_2$  in root exudate solutions varied among the four plant species (Fig. S8 and S9†). The largest size (199 nm) was found for corn, which might be one possible reason for the low uptake of Ce in corn. The zeta potential of  $\text{nCeO}_2$  in the exudates under the +P and -P conditions and among the different species are not significantly different (Table S5†), suggesting that surface charges are not associated with the Ce translocation. Since the transformation of  $\text{nCeO}_2$  mainly occurs on the root surface with the assistance of the root exudates,<sup>16</sup> the compositional difference in the root exudates may lead to different extent of transformation of  $\text{nCeO}_2$  on the root surface and subsequently different translocation of Ce. Root exudates are composed of low molecular compounds such as saccharides,

organic acids, amino acids, phenols, high molecular polysaccharides and proteins.<sup>27</sup> The low molecular organic acids (e.g., succinic acid and malic acid)<sup>8</sup> and reducing substances such as phenols,<sup>28</sup> which are critical for the transformation of  $\text{nCeO}_2$ , varied among different species. Moreover, different acids and reducing substances have different capacities to dissolve the  $\text{nCeO}_2$ .<sup>8,29</sup> For example, mugineic acids, which have been found to promote the transformation of  $\text{CuO}$  NPs,<sup>30</sup> are specifically excreted by gramineous plants (e.g., corn and wheat).<sup>31</sup> Turnip cabbage excretes two more kinds of organic acids (i.e. malic and glycolic acids) than cucumber.<sup>32</sup> Except for the compositional differences, amounts of exudates vary among different species and at different growth stages for the same plant species.<sup>33</sup> Plant roots at the seedling stage generally excrete a greater amount of reducing substances than that at the germination stage, which may explain why no transformation of  $\text{CeO}_2$  in the roots of the four plant species was found in a previous study.<sup>14</sup> We also noticed that in the +P treatment, the TFs of  $\text{Ce(III)}$  in the dicots are actually higher than those in the monocots (Table S4 and Fig. S7†), which may not be explained by either the xylem structures or the exudate compositions. This indicates an additional mechanism influencing the translocation of Ce on top of the xylem structures and the composition of exudates, e.g., dissolution of  $\text{nCeO}_2$  inside the leaves.

## 4. Conclusions

In this study, the plant species-dependent transformation and translocation of  $\text{nCeO}_2$  were comprehensively studied for the first time. Although the exposure concentration used in this study is much higher than the predicted environmental concentration of  $\text{nCeO}_2$  (e.g., 0.25–1  $\text{mg kg}^{-1}$ )<sup>34</sup> and natural background concentration of Ce (48  $\text{mg kg}^{-1}$  dry soil),<sup>35</sup> the mechanistic understanding of  $\text{nCeO}_2$  behaviour in plants in this study is of vital importance. Our results suggest that different plant species have different abilities of transforming and translocating the  $\text{CeO}_2$  NPs and that the mechanisms differ depending on the chemistry of the surrounding solution (+P/-P). Plant xylems and root exudate compositions are both important factors in determining the transformation of  $\text{nCeO}_2$  and subsequent translocation of Ce species in plants. Further studies are required to identify the root exudate components and the determining components for  $\text{nCeO}_2$  transformation. In addition, the components of root exudates may vary (e.g., increase of organic acid levels)<sup>36</sup> due to ambient stress (e.g., NP exposure) for the same plant species, and therefore, extracting and determining the exudate composition after NP exposure are important. Furthermore, different plant species respond differently to  $\text{nCeO}_2$  under different culturing conditions. For some plant species (e.g., corn and wheat in this study), deficiency of P can increase the potential phytotoxicity of  $\text{nCeO}_2$ ,<sup>37</sup> and enhance the accumulation of Ce (mainly in the form of  $\text{Ce}^{3+}$ ) in plants. Besides, deficiency of P is only one of the conditions that might occur for plants, and more factors (e.g., pH values, redox state, and the





presence of other ions) that may affect the conditions around the root should be considered in order to comprehensively understand the transformation of NPs in plants.

## Conflicts of interest

There is no conflict of interest to declare.

## Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (Grants 11405183, 11575208, 11675190, 11875267, and 21507153) and the Ministry of Science and Technology of China (Grant 2016YFA0201600). Financial support from a Marie Skłodowska-Curie Individual Fellowship (NanoLabels 750455) is appreciated.

## References

- 1 T. Marie, A. Mélanie, B. Lenka, I. Julien, K. Isabelle, P. Christine, M. Elise, S. Catherine, A. Bernard and A. Ester, Transfer, transformation, and impacts of ceria nanomaterials in aquatic mesocosms simulating a pond ecosystem, *Environ. Sci. Technol.*, 2014, **48**, 9004–9013.
- 2 L. E. Barton, M. Auffan, L. Olivi, J.-Y. Bottero and M. R. Wiesner, Heteroaggregation, transformation and fate of CeO<sub>2</sub> nanoparticles in wastewater treatment, *Environ. Pollut.*, 2015, **203**, 122–129.
- 3 L. E. Barton, M. Auffan, M. Bertrand, M. Barakat, C. Santaella, A. Masion, D. Borschneck, L. Olivi, N. Roche and M. R. Wiesner, Transformation of pristine and citrate-functionalized CeO<sub>2</sub> nanoparticles in a laboratory-scale activated sludge reactor, *Environ. Sci. Technol.*, 2014, **48**, 7289–7296.
- 4 R. C. Merrifield, K. P. Arkill, R. E. Palmer and J. R. Lead, A high resolution study of dynamic changes of Ce<sub>2</sub>O<sub>3</sub> and CeO<sub>2</sub> nanoparticles in complex environmental media, *Environ. Sci. Technol.*, 2017, **51**, 8010–8016.
- 5 X. He, Y. Pan, J. Zhang, Y. Li, Y. Ma, P. Zhang, Y. Ding, J. Zhang, Z. Wu and Y. Zhao, Quantifying the total ionic release from nanoparticles after particle-cell contact, *Environ. Pollut.*, 2015, **196**, 194–200.
- 6 P. Zhang, Y. Ma, Z. Zhang, X. He, J. Zhang, Z. Guo, R. Tai, Y. Zhao and Z. Chai, Biotransformation of ceria nanoparticles in cucumber plants, *ACS Nano*, 2012, **6**, 9943–9950.
- 7 Y. Ma, P. Zhang, Z. Zhang, X. He, Y. Li, J. Zhang, L. Zheng, S. Chu, K. Yang and Y. Zhao, Origin of the different phytotoxicity and biotransformation of cerium and lanthanum oxide nanoparticles in cucumber, *Nanotoxicology*, 2015, **9**, 262–270.
- 8 W. Zhang, Y. Dan, H. Shi and X. Ma, Elucidating the mechanisms for plant uptake and in-plant speciation of cerium in radish (*Raphanus sativus* L.) treated with cerium oxide nanoparticles, *J. Environ. Chem. Eng.*, 2017, **5**, 572–577.
- 9 F. Schwabe, S. Tanner, R. Schulín, A. Rotzetter, W. Stark, A. Von Quadt and B. Nowack, Dissolved cerium contributes to uptake of Ce in the presence of differently sized CeO<sub>2</sub> nanoparticles by three crop plants, *Metallomics*, 2015, **7**, 466–477.
- 10 C. Levard, E. M. Hotze, B. P. Colman, A. L. Dale, L. Truong, X. Yang, A. J. Bone, G. E. Brown Jr, R. L. Tanguay and R. T. Di Giulio, Sulfidation of silver nanoparticles: natural antidote to their toxicity, *Environ. Sci. Technol.*, 2013, **47**, 13440–13448.
- 11 J. Cervini-Silva, D. A. Fowle and J. Banfield, Biogenic dissolution of a soil cerium-phosphate mineral, *Am. J. Sci.*, 2005, **305**, 711–726.
- 12 P. Zhang, Y. Ma, Z. Zhang, X. He, Y. Li, J. Zhang, L. Zheng and Y. Zhao, Species-specific toxicity of ceria nanoparticles to Lactuca plants, *Nanotoxicology*, 2015, **9**, 1–8.
- 13 P. Zhang, C. Xie, Y. Ma, X. He, Z. Zhang, Y. Ding, L. Zheng and J. Zhang, Shape-Dependent Transformation and Translocation of Ceria Nanoparticles in Cucumber Plants, *Environ. Sci. Technol. Lett.*, 2017, **4**, 380–385.
- 14 M. L. López-Moreno, G. de la Rosa, J. A. Hernández-Viezas, J. R. Peralta-Videa and J. L. Gardea-Torresdey, XAS corroboration of the uptake and storage of CeO<sub>2</sub> nanoparticles and assessment of their differential toxicity in four edible plant species, *J. Agric. Food Chem.*, 2010, **58**, 3689.
- 15 Z. Zhu, H. Wang, B. Yan, H. Zheng, Y. Jiang, O. R. Miranda, V. M. Rotello, B. Xing and R. W. Vachet, Effect of surface charge on the uptake and distribution of gold nanoparticles in four plant species, *Environ. Sci. Technol.*, 2012, **46**, 12391–12398.
- 16 Y. Ma, P. Zhang, Z. Zhang, X. He, J. Zhang, Y. Ding, J. Zhang, L. Zheng, Z. Guo and L. Zhang, Where does the transformation of precipitated ceria nanoparticles in hydroponic plants take place?, *Environ. Sci. Technol.*, 2015, **49**, 10667–10674.
- 17 EPA Ecological effects test guidelines, OPPTS 850.4150 Terrestrial Plant Toxicity, Tier I (vegetative Vigor), EPA 712-C-96-163 (Public Draft); Office of Prevention, Pesticides and Toxic Substances, U.S. Environmental Protection Agency, Washington, DC, 1996.
- 18 A. Gaur, B. Shrivastava and S. Joshi, Copper K-edge XANES of Cu(I) and Cu(II) oxide mixtures, *J. Phys.: Conf. Ser.*, 2009, **190**, 012084.
- 19 R. Russell and V. Shorrocks, The relationship between transpiration and the absorption of inorganic ions by intact plants, *J. Exp. Bot.*, 1959, **10**, 301–316.
- 20 P. Zhang, Y. Ma, S. Liu, G. Wang, J. Zhang, X. He, J. Zhang, Y. Rui and Z. Zhang, Phytotoxicity, uptake and transformation of nano-CeO<sub>2</sub> in sand cultured romaine lettuce, *Environ. Pollut.*, 2017, **220**, 1400–1408.
- 21 R. Bielecki, Phosphate pools, phosphate transport, and phosphate availability, *Annu. Rev. Plant Biol.*, 1973, **24**, 225–252.
- 22 S. Schymura, T. Fricke, H. Hildebrand and K. Franke, Elucidating the Role of Dissolution in CeO<sub>2</sub> Nanoparticle Plant Uptake by Smart Radiolabeling, *Angew. Chem., Int. Ed.*, 2017, **56**, 1–5.
- 23 N. Edayilam, D. A. Montgomery, B. O. Ferguson, A. S. Maroli, N. Martinez, B. A. Powell and N. Tharayil,





- Phosphorus stress-induced changes in plant root exudation could potentially facilitate uranium mobilization from stable mineral forms, *Environ. Sci. Technol.*, 2018, 52, 7652–7662.
- 24 W. Zhang, S. D. Ebbs, C. Musante, J. C. White, C. Gao and X. Ma, Uptake and accumulation of bulk and nanosized cerium oxide particles and ionic cerium by radish (*Raphanus sativus* L.), *J. Agric. Food Chem.*, 2015, 63, 382–390.
  - 25 P. J. White, in *Marschner's Mineral Nutrition of Higher Plants*, Elsevier, 3rd edn, 2012, pp. 49–70.
  - 26 Z. Zhang, X. He, H. Zhang, Y. Ma, P. Zhang, Y. Ding and Y. Zhao, Uptake and distribution of ceria nanoparticles in cucumber plants, *Metallomics*, 2011, 3, 816–822.
  - 27 K. Tawarayama, R. Horie, T. Shinano, T. Wagatsuma, K. Saito and A. Oikawa, Metabolite profiling of soybean root exudates under phosphorus deficiency, *Soil Sci. Plant Nutr.*, 2014, 60, 679–694.
  - 28 V. Vančura, Root exudates of plants, *Plant Soil*, 1964, 21, 231–248.
  - 29 Y. Rui, P. Zhang, Y. Zhang, Y. Ma, X. He, X. Gui, Y. Li, J. Zhang, L. Zheng and S. Chu, Transformation of ceria nanoparticles in cucumber plants is influenced by phosphate, *Environ. Pollut.*, 2015, 198, 8–14.
  - 30 P. McManus, *Rhizosphere interactions between copper oxide nanoparticles and wheat root exudates in a sand matrix; influences on bioavailability and uptake*, Utah State University, 2016.
  - 31 T. W.-M. Fan, A. N. Lane, M. Shenker, J. P. Bartley, D. Crowley and R. M. Higashi, Comprehensive chemical profiling of gramineous plant root exudates using high-resolution NMR and MS, *Phytochemistry*, 2001, 57, 209–221.
  - 32 V. Vančura and A. Hovadik, Root exudates of plants: II. Composition of root exudates of some vegetables, *Plant Soil*, 1965, 21–32.
  - 33 V. Vančura and A. Hanzlíková, Root exudates of plants: IV. Differences in chemical composition of seed and seedlings exudates, *Plant Soil*, 1972, 271–282.
  - 34 F. Gottschalk, T. Sun and Bernd Nowack, Environmental concentrations of engineered nanomaterials: Review of modeling and analytical studies, *Environ. Pollut.*, 2013, 181, 287–300.
  - 35 G. Tyler, Rare earth elements in soil and plant systems - A review, *Plant Soil*, 2004, 267, 191–206.
  - 36 P. Wang, S. Bi, S. Wang and Q. Ding, Variation of wheat root exudates under aluminum stress, *J. Agric. Food Chem.*, 2006, 54, 10040–10046.
  - 37 G. Wang, Y. Ma, P. Zhang, X. He, Z. Zhang, M. Qu, Y. Ding, J. Zhang, C. Xie and W. Luo, Influence of phosphate on phytotoxicity of ceria nanoparticles in an agar medium, *Environ. Pollut.*, 2017, 224, 392–399.

