



Cite this: *Food Funct.*, 2019, **10**, 4452

## Correction: Different antitumor effects of quercetin, quercetin-3'-sulfate and quercetin-3-glucuronide in human breast cancer MCF-7 cells

Qiu Wu,<sup>a</sup> Paul W. Needs,<sup>\*b</sup> Yalong Lu,<sup>a</sup> Paul A. Kroon,<sup>b</sup> Daoyuan Ren<sup>a</sup> and Xingbin Yang<sup>\*a</sup>

DOI: 10.1039/c9fo90027f  
rsc.li/food-function

Correction for 'Different antitumor effects of quercetin, quercetin-3'-sulfate and quercetin-3-glucuronide in human breast cancer MCF-7 cells' by Qiu Wu *et al.*, *Food Funct.*, 2018, **9**, 1736–1746.

The authors regret that the panel for Q3G-100  $\mu\text{M}$  in Fig. 3A was shown incorrectly in the original article. The correct version of Fig. 3 is as shown below.

Consequently, sections of the text in the manuscript should be adjusted according to this change, and these are detailed below.


The sentence in the abstract beginning “Moreover, it was found that 70.8%, 58.2%, and 48.0% of MCF-7 cancer cells...” should be correctly given as “Moreover, it was found that 70.8%, 58.2%, and 49.8% of MCF-7 cancer cells entered the early phase of apoptosis when treated with 100  $\mu\text{M}$  Que, Q3'S, and Q3G for 48 h, respectively”.

The sentence on page 1739 beginning “Furthermore, Que, Q3'S, and Q3G at the concentrations of 100  $\mu\text{M}$ ...” should be correctly given as “Furthermore, Que, Q3'S, and Q3G at the concentrations of 100  $\mu\text{M}$  dose-dependently resulted in 70.8%, 58.2%, and 49.8% of early apoptotic cells, which were similar to that of the 5-Fu group (75.2%,  $p > 0.05$ )”.

<sup>a</sup>Shaanxi Engineering Laboratory for Food Green Processing and Safety Control, and Key Laboratory of Ministry of Education for Medicinal Resource and Natural Pharmaceutical Chemistry, College of Food Engineering and Nutritional Science, Shaanxi Normal University, Xi'an 710062, China. E-mail: xbyang@snnu.edu.cn

<sup>b</sup>Quadram Institute Bioscience, Norwich Research Park, Colney, Norwich, NR4 7UA, UK. E-mail: needsp@quadram.ac.uk





(B)




Fig. 3 Quantitative analysis of apoptotic cells induced by Que, Q3'S, and Q3G using the annexin V/PI double staining assay. (A) Representative dot plots of Annexin V/PI staining. (B) Column bar graph of apoptotic cells. Cells were treated with Que, Q3'S, and Q3G at 25 and 100  $\mu\text{M}$  for 48 h, respectively. 3000 cells were analyzed by flow cytometry. The results are expressed as mean  $\pm$  SD of three independent experiments.  $p < 0.01$ (\*\*), as compared to the control.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

