## Food & Function



## CORRECTION

View Article Online
View Journal | View Issue



Cite this: Food Funct., 2019, 10, 5240

## Correction: Total saponins from *Rosa laevigata*Michx fruit attenuates hepatic steatosis induced by high-fat diet in rats

Deshi Dong, a,b,c Yan Qi, a Lina Xu, a Lianhong Yin, a Youwei Xu, a Xu Han, a Yanyan Zhao a and Jinyong Peng\*a,b

DOI: 10.1039/c9fo90039j

Correction for 'Total saponins from *Rosa laevigata* Michx fruit attenuates hepatic steatosis induced by high-fat diet in rats' by Deshi Dong, et al., Food Funct., 2014, **5**, 3065–3075.

The authors regret that there were errors in Fig. 4 and Fig. 6 in the original manuscript. The GAPDH bands in these figures for the western blot assay were incorrect due to copy and paste errors. The corrected versions of Fig. 4 and Fig. 6 are presented below.




Fig. 4 Effects of RLTS on the protein expressions of LXRα, PPARα, MCAD, ACADS, CYP2E1, CYP4A11, and the gene expressions of PPAR $_{\gamma}$ , SREBP-1c, FAS, ACC, SCD1, CPT1 from rat livers in normal control (II), model (III), RLTS (70 mg kg $^{-1}$ ) + HFD (IV), RLTS (140 mg kg $^{-1}$ ) + HFD (V) and RLTS (210 mg kg $^{-1}$ ) + HFD (VI) groups. Values were expressed as mean  $\pm$  SD in each group (n = 5). \*p < 0.05, \*\*p < 0.01 vs. model group.

<sup>&</sup>lt;sup>a</sup>College of Pharmacy, Dalian Medical University, 9 Western Lyshun South Road, Dalian 116044, China

<sup>&</sup>lt;sup>b</sup>Research Institute of Integrated Traditional and Western Medicine of Dalian Medical University, Dalian 116011, China. E-mail: jinyongpeng2005@163.com; Fax: +86411 8611 0411; Tel: +86 411 8611 0411

<sup>&</sup>lt;sup>c</sup>The First Affiliated Hospital of Dalian Medical University, Dalian 116011, China

Food & Function Correction




Fig. 6 Effects of RLTS on the protein expressions of HO-1, Nrf2, kEAP1, SOD2, ALDH7A1, NF- $\kappa$ B, COX-2, HMGB-1, AP-1, and the gene expressions of TNF- $\alpha$ , IL-6 from rat livers in normal control (II), model (III), RLTS (70 mg kg<sup>-1</sup>) + HFD (IV), RLTS (140 mg kg<sup>-1</sup>) + HFD (V) and RLTS (210 mg kg<sup>-1</sup>) + HFD (VI) groups. Values were expressed as mean  $\pm$  SD in each group (n = 5). \*p < 0.05, \*\*p < 0.01 vs. model group.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.