Cite this: *Mol. Syst. Des. Eng.*, 2019, 4, 984

DOI: 10.1039/c9me90025j

rsc.li/molecular-engineering

Correction: Star-shaped π -gelators based on oxadiazole and thiadiazoles: a structure–property correlation

Subrata Nath,^a Suraj Kumar Pathak,^a Joydip De,^b
Santanu Kumar Pal^b and Ammathnadu S. Achalkumar^{*a}

Correction for ‘Star-shaped π -gelators based on oxadiazole and thiadiazoles: a structure–property correlation’ by Subrata Nath *et al.*, *Mol. Syst. Des. Eng.*, 2017, 2, 478–489.

The authors regret that Fig. 9 and 12(d) of the published article were incorrect. The corrected versions of both affected figures are given below. Please note that the results and conclusions reported in the article are not affected by these errors.

Fig. 9 AFM image obtained for the xerogel of compound **4a** (a); XRD profile depicting the intensity against the 2θ obtained for the crystalline phase of compound **4a** in xerogel state (inset shows the XRD image pattern obtained) (b); SEM images obtained for compound **4a** (c); an expanded region of the image in c (scale bar is 1 μm) (d).

^a Department of Chemistry, Indian Institute of Technology Guwahati, Guwahati, 781039, Assam, India. E-mail: achalkumar78@gmail.com, achalkumar@iitg.ernet.in

^b Indian Institute of Science Education and Research Mohali, Sector-81, Knowledge City, Manauli 140306, Punjab, India

Fig. 12 Angular frequency (rad s^{-1}) dependence of G' and G'' obtained with small stress amplitudes: (a) 0.9 wt% of compound **4a** in dodecane at 25 °C using 0.2 kPa stress; (b) 0.6 wt% of compound **7a** in dodecane at 25 °C using 0.2 kPa stress; (c) 0.5 wt% of compound **7b** in dodecane at 25 °C using 0.2 kPa stress; stress dependence of G' and G'' of the same gels measured at constant frequency (at 25 °C, and frequency at 0.1 Hz) for compound **4a** (d); compound **7a** (e) and compound **7b** (f).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.