



Cite this: *Org. Biomol. Chem.*, 2019, **17**, 2287

## Correction: Glycosylated naphthalimides and naphthalimide Tröger's bases as fluorescent aggregation probes for Con A


Elena Calatrava-Pérez,<sup>a</sup> Jason M. Delente,<sup>a</sup> Sankarasekaran Shanmugaraju,<sup>a</sup> Chris S. Hawes,<sup>b</sup> Clive D. Williams,<sup>c</sup> Thorfinnur Gunnlaugsson\*<sup>a</sup> and Eoin M. Scanlan\*<sup>a</sup>

DOI: 10.1039/c9ob90017a

rsc.li/obc

Correction for 'Glycosylated naphthalimides and naphthalimide Tröger's bases as fluorescent aggregation probes for Con A' by Elena Calatrava-Pérez *et al.*, *Org. Biomol. Chem.*, 2019, DOI: 10.1039/c8ob02980f.

The authors regret that there were errors in the structure of compounds 2/3 in Scheme 1. The correct scheme is shown below.



**Scheme 1** The structure of Nap 1, and the synthetic strategy for the preparation of TBNap 3 from the TB anhydride 4 and the precursor 2.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>School of Chemistry and Trinity Biomedical Sciences Institute (TBSI), Trinity College Dublin, The University of Dublin, Dublin 2, Ireland. E-mail: gunnlaut@tcd.ie, eoin.scanlan@tcd.ie

<sup>b</sup>School of Chemical and Physical Sciences, Keele University, Keele ST5 5BG, UK

<sup>c</sup>School of Biochemistry and Immunology and Trinity Biomedical Sciences Institute (TBSI), Trinity College Dublin, The University of Dublin, Dublin 2, Ireland

