



Cite this: RSC Adv., 2019, 9, 172

# Secondary metabolites (SMs) of *Isaria cicadae* and *Isaria tenuipes*

Xiaofeng Zhang,  Qiongbao Hu \* and Qunfang Weng \*

Both *Isaria cicadae* and *Isaria tenuipes* are important entomopathogenic fungi used in health foods and traditional herbal medicines in East Asia. However, the safety concerns for both fungal species have been attracting significant attention. Thus, surveying their secondary metabolites (SMs) will be beneficial to improving the safety of their fungal products. In the case of *I. cicadae*, its SMs mainly include nucleosides, amino acids, beauvericins, myriocin, and oosporein. In contrast, trichothecene derivatives, isariotins, cyclopenta benzopyrans and PKs, are found in the case of *I. tenuipes*. Among them, beauvericins, myriocin, oosporein and many trichothecene derivatives are toxic compounds. The toxicity and side effects of the fungal products may be related to these SMs. Thus, to ensure the safety of fungal products, the residues standards of SMs need to be reported. Furthermore, methods for the detection of their SMs and biological identification of their strains must be considered. This review gives new insight into the secondary metabolites of medical and edible fungi.

Received 1st November 2018  
Accepted 29th November 2018

DOI: 10.1039/c8ra09039d

rsc.li/rsc-advances

## 1 Introduction

*Isaria* is an entomopathogenic fungal genus with more than 100 species (<http://www.mycobank.org/>), which plays important roles in biodiversity conservation and is utilized in medicines and agriculture.<sup>1,2</sup> Both *I. cicadae* and *I. tenuipes* (Fig. 1) belonging to the anamorph of *Cordyceps* genus, are popular mushrooms mostly infecting insects (in nature, often the underground nymphs of cicadas), leading to their death, and then produces white fruit body from the dead insect, which releases white powdery spores. The fruit body is used in expensive traditional medicines and health products in East Asia.<sup>2–4</sup>

*I. cicadae*, the asexual type of *C. cicadae*, is a famous mushroom. In nature, the fruiting bodies form uniquely on pupated cicadae (*Platydomia* spp.) and are referred to as “cicada flower” or the Chinese name “Jingchanhua” in China. In the laboratory, this fungus can be inoculated in the pupae of Chinese tussah silkworm (*Antheraea pernyi*). Interestingly, in contrast to other *Cordyceps* species that produce sexual fruiting bodies in nature, *C. cicadae* forms synnema-like asexual structures.<sup>5</sup> *I. sinclairii* was once known as *C. cicadae*, *C. kobayashii*, *Paecilomyces cicadae*, *P. sinclairii* and *C. sinclairii*.<sup>6–8</sup> However, it is considered as the synonym of *I. cicadae*.<sup>9</sup> This species is found in Asia (particularly China, Japan, and Korea) and New Zealand. It infects cicadas of the genera *Amphipsalta* and *Melampsalta* in nature, and in the lab it can be cultured on the bodies of silkworms,<sup>10</sup> which also shows potential as a biological control agent against the

agricultural pest *Plutella xylostella*.<sup>11</sup> Its fruit body is used to treatment cancers and improve the eyesight and renal function and immunity of patients.<sup>12</sup> It also has anti-hypertensive effect in spontaneously hypertensive rats.<sup>13</sup> However, it was reported that it induced kidney toxicity in rats exposed to its fungus fruit bodies.<sup>14,15</sup>

*I. tenuipes* is a common entomopathogenic fungus that usually infects Lepidopteran insects.<sup>16</sup> It has the synonymies of *C. tenuipes* and *P. tenuipes*, and it was also called *Spicaria heliothis*, *C. concurrens*, *C. polyarthra*, and *C. subpolyarthra* (<http://www.mycobank.org/>). Moreover, it is considered as the anamorph of *C. takaomontana*<sup>17</sup> and synonymy *I. japonica*.<sup>18</sup> Its fruiting body is used in the production of health foods and traditional medicine with the functions of lowering blood-glucose, antitumor, antibacterial, anti-depression, anti-oxidation, and anti-aging activity, lowering blood fat and immunoregulation.<sup>7,18,19</sup>

However, safety concerns of consuming these entomopathogenic fungi have been frequently raised due to the uncertainty of the fungal production of human-toxic mycotoxins.<sup>20</sup> In fact, *Isaria* fungi have rigorous secondary metabolisms through different pathways, including non-ribosomal peptide synthetase (NRPS), polyketide synthase (PKS) and terpenoid synthetase (TS), which produce various secondary metabolites (SMs), such as ribosomal peptides (NRPs), polyketides (PKs), terpenoid and miscellaneous types.<sup>5,21,22</sup> According to the research (patents not included) published in the past forty years, it is estimated that more than 200 SMs have been isolated and identified from both *Isaria* fungi. Many SMs are the functional components with anti-virus, anti-bacteria, and anti-tumor activity and immunity regulation.<sup>23,24</sup> Interestingly, SMs are

College of Agriculture, South China Agricultural University, Guangzhou 510642, China. E-mail: hqbscau@scau.edu.cn; wengweng@scau.edu.cn






Fig. 1 General characteristics and morphology of *Isaria cicadae* and *Isaria tenuipes*. (A–D) *Isaria cicadae* (<http://www.thai2bio.net/>; <http://www.biophar.cn/>) and (E–F) *Isaria tenuipes* (<http://www.naro.affrc.go.jp/>). (A) and (E) wild complex of fruiting body and host, FB: fruiting bodies, HI: host insects, (B) and (F) microscopic photos, CS: conidial structures, (C) cultured fruiting bodies and (D) commercial product.

considered as important drug resources. For example, fingolimod, a novel type of immunosuppressive agent, is a new medicine derived from *I. sinclairii*, which is used for treating multiple sclerosis, renal cancer, and asthma.<sup>25,26</sup> However, many SMs belong to mycotoxins, which are toxic to humans and animals. Nevertheless, there are few documents reviewing the risks of *Isaria* fungi mycotoxins. Thus, the current review focuses on the mycotoxins of both *Isaria* fungi, including their structures, bioactivities and toxicities, as well as their risk evaluation.

## 2 SMs from *I. cicadae*

*I. cicadae* produces numerous SMs (Table 1 and Fig. 2). Using metabolism omics analysis, Lu *et al.* reported that more than 110 compounds were detected in the mycelia, primordia and stroma stages of the fungus.<sup>5</sup> While eight compounds were found in *C. cicadae* sporoderm-broken spore powders (CCBSP) using UPLC-ESI-Q-TOF-MS, and forty-nine volatile molecules representing 99.56% of CCBSP were clearly identified by GC-MS.<sup>27</sup> Moreover, another experiment reported that 60 metabolites including amino acids and their derivatives, saccharides, organic acids and fatty acids were identified in the methanol extract of the complex of the cicadae body and *I. cicadae* stroma, while inositol, gamma-aminobutyric acid, ornithine and threonine were the most abundant compounds.<sup>28</sup>

Based on their molecular structures, *I. cicadae* SMs are mainly categorized as organic acids, amino acids, lipids and phospholipids, nucleosides, carbohydrates and their derivatives. Among them, some compounds are functional or bioactive components, for example, cordycepin (1) and cordycepic acid (3) are considered as the bioactive components of *I. cicadae*, similar with *Cordyceps* spp.<sup>29</sup> Furthermore, many other compounds such as adenosine, guanosine, uridine, inosine and

thymidine and other nucleosides, ergosterol and its peroxide, and D-mannitol beneficial for pharmacological functions.<sup>30,31</sup> However, some of the SMs are toxic to humans and animals, while probably most SMs are unknown, and their molecular structures and biological functions have not been elucidated.<sup>5</sup> The main nucleoside of this fungus is cordycepin (1), 3'-deoxyadenosine, which was isolated from the sporoderm-broken spore powders and fruit bodies of *C. cicadae*.<sup>27,32</sup> As an adenosine derivative, it is an important parameter for quality control in medicines, referring to Dongchongxiacao CORDYCEPS, *Cordyceps sinensis* (BerK.) Sacc. in the Chinese Pharmacopoeia, 2015. Since cordycepin (1) is similar to adenosine, it can participate in certain biochemical reactions (for example, be incorporated into an RNA molecule, thus causing the premature termination of its synthesis)<sup>33</sup> and has multiple bioactivities such as anti-cancer, antimetastatic, antioxidant, antidepressant and immunoregulation properties.<sup>34–37</sup> It also enhances male reproduction by stimulating *in vitro* and *in vivo* steroidogenesis in mouse Leydig cells by activating the PKA pathway.<sup>38</sup> In 2007, the US Food and Drug Administration (FDA) granted cordycepin orphan drug designation for the treatment of TdT-positive acute lymphocytic leukemia (<https://www.fda.gov/>). Cordycepin plus pentostatin was subjected to a clinic trial of phase 1/phase 2 in patients with refractory TdT-positive leukemia in 2008–2010 (<https://www.clinicaltrials.gov/>). In China, it was reported that cordycepin was used for clinic adjuvant treatment of non-small cell lung cancer<sup>39</sup> and nephropathy of type 2 diabetes mellitus.<sup>40</sup> However, the main fungal species used to produce cordycepin in industry is *C. militaris*.<sup>41</sup>

Another adenosine derivative, *N*-(2-hydroxyethyl) adenosine (2) was isolated and proven to exhibit antioxidant activity by measuring its radical scavenging effect on 1,1-diphenyl-2-picrylhydrazyl (DPPH) radicals.<sup>42</sup>



Table 1 SMSs isolated from *I. cicadae* and their biological activities

Metabolite	CAS no.	Material source	Biological activity
Cordycepin (1)	73-03-0	Sporoderm-broken spore powders of wild <i>C. cicadae</i>	Immunomodulatory, antibacterial, antiviral, and anti-tumor effects on cells
<i>N</i> -(2-Hydroxyethyl) adenosine (2)	4338-48-1	Cultivated complex of fruit body of <i>I. sinclairii</i> and silkworm	Bacteriostatic activity
Cordycepic acid (3)	69-65-8	Sporoderm-broken spore powders of wild <i>C. cicadae</i>	
Betaine (4)	107-43-7	Mycelia of <i>C. cicadae</i> CCAD02	
Hercynine (5)	534-30-5	Mycelia of <i>C. cicadae</i> CCAD02	Regulate internal osmotic pressure, relieve stress, and promote fat metabolism and protein synthesis
Ergothioneine (6)	497-30-3	Mycelia of <i>C. cicadae</i> CCAD02	Antioxidant
Fumimycin (7)	942472-95-9	Mycelia, cultivated complex of fruit body of <i>C. cicadae</i> CCAD02 and <i>Antheraea pernyi</i>	Antioxidant
Cordysin A (8)	1330197-09-5	Mycelia of <i>C. cicadae</i> CCAD02	Inhibiting the proliferation of human glioma U87 mg and U251 cells. Anti-inflammatory activity in human neutrophils
Beauvericin (9)	26048-05-5	Mycelia, cultivated complex of fruit body of <i>C. cicadae</i> CCAD02 and <i>Antheraea pernyi</i>	Significant inhibitory effect on hepg2 and hepg2/ADM cells with IC <sub>50</sub> values ranging from 2.40–14.48 μM. Cytotoxicity against hepg2/ADM cell line with IC <sub>50</sub> value 25-fold more sensitive to doxorubicin
Beauvericin A (10)	165467-50-5	Wild complex of fruit body of <i>C. cicadae</i> and insect-body (specimen no. 2012081827)	
Beauvericin E (11)	728912-27-4	Wild complex of fruit body of <i>C. cicadae</i> and insect-body (specimen no. 2012081827)	
Beauvericin J (12)	1342821-26-4	Wild complex of fruit body of <i>C. cicadae</i> and insect-body (specimen no. 2012081827)	Significant inhibitory effect on hepg2 and hepg2/ADM cells with IC <sub>50</sub> values ranging from 2.40–14.48 μM
Cordycecin A (13)	1799406-02-2	Wild complex of fruit body of <i>C. cicadae</i> and insect-body (specimen no. 2012081827)	Significant inhibitory effect on hepg2 and hepg2/ADM cells with IC <sub>50</sub> values ranging from 2.40–14.48 μM
Aspergilliamide (14)	1691204-76-8	Mycelia of <i>C. cicadae</i> CCAD02	Medium brine shrimp toxicity
Myriocin (15)	35891-70-4	Sporoderm-broken spore powders of wild <i>C. cicadae</i> . Cultivated mycelia of <i>C. cicadae</i>	
Phytosphingosine (16)	554-62-1	Mycelia, cultivated complex of fruit body of <i>C. cicadae</i> CCAD02 and <i>Antheraea pernyi</i>	Food refusal effect on crop pests <i>Prodenia litura</i> and <i>Spodoptera exigua</i> . Insecticide to <i>Lepidopteran</i> pests
Oosporein (17)	475-54-7	Mycelia, cultivated complex of fruit body of <i>C. cicadae</i> CCAD02 and <i>Antheraea pernyi</i>	
Stipitonic acid (18)	606-39-3	Mycelia, cultivated complex of fruit body of <i>C. cicadae</i> CCAD02 and <i>Antheraea pernyi</i>	Bactericidal and insecticidal activities, as well as growth-inhibitory effects on some pathogenic fungi
Cycloheximide acid A (19)	1610848-67-3	Mycelia, cultivated complex of fruit body of <i>C. cicadae</i> CCAD02 and <i>Antheraea pernyi</i>	
2-(3-Carboxy-2-hydroxypropyl)-3-methyl-2-cyclopentenone (20)	1019196-97-4	Cultivated mycelia of <i>C. cicadae</i>	
5-(2-Hydroxyethyl)-2-furanacetic acid (21)	1018901-08-0	Cultivated mycelia of <i>C. cicadae</i>	



Table 1 (Contd.)

Metabolite	CAS no.	Material source	Biological activity
Lichenicolin A (22)	883977-62-6	Mycelia, cultivated complex of fruit body of <i>C. cicadae</i> CCAD02 and <i>Antheraea pernyi</i>	
2-(5-(3-Oxobutyl)furan-2-yl)acetic acid (23)		A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	Weak inhibitory activity against AChE
Cordycepene (24)		A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	
Cephalosporolide E (25)	97373-15-4	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	
Cephalosporolide J (26)	97344-02-0	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	
Methyl-2-(5-(3-hydroxybutyl)furan-2-yl)acetate (27)	851868-04-7	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	Weak inhibitory activity against AChE
2-(5-(3'-Hydroxybutyl) furan-2-yl)acetic acid (28)	170445-59-7	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	Weak inhibitory activity against AChE
$\alpha$ -furoic acid (29)	88-14-2	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	Weak inhibitory activity against AChE. Moderate inhibitory activity against the nematode <i>Panagrellus redivivus</i>
(22E,24R)-Ergosta-5,7, trien-3 $\beta$ -ol (ergosterol) (30)	57-87-4	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	
3-Benzyl-6-isopropyl-2,5-piperazinedione (31)	14474-71-6	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	Moderate inhibitory activity against the nematode <i>Panagrellus redivivus</i>
3-Isobutyl-6-isobutylpiperazine-2,5-dione (32)	5625-50-3	A strain of cell fusion from <i>C. cicadae</i> and <i>C. militaris</i>	

Cordycepic acid (3) (D-mannitol), an isomer of quinic acid with a polyol structure, was isolated from sporoderm-broken spore powders.<sup>27</sup> It is believed to be another active ingredient and is used as an important index of quality control for *C. sinensis* products. It exhibits diuretic action and prophylaxis against postoperative acute renal failure, relieving cough and asthma, and anti-free radical activities. Specifically, it can be used to provide effective protection or treatment for patients after cerebral ischemia and trauma, such as improving cerebral microcirculation and cerebral blood flow.<sup>43</sup>

The amino derivative betaine (4) (*N,N,N*-trimethylglycine), which was first found in sugar beet in the nineteenth century, was detected in the mycelia, primordia and stroma of *C. cicadae*.<sup>5</sup> Betaine (4) is a methyl donor of increasingly recognized significance in biology, and serves as an organic osmolyte, which is a substance synthesized or taken up from the environment by cells for protection against osmotic stress, drought, high salinity, and high temperature.<sup>44</sup> Hercynine (5), a derivative of histidine, was detected in the mycelia, primordia and stroma of *C. cicadae*.<sup>5</sup> However, ergothioneine (6) was identified in its fruit body and mycelia.<sup>45</sup> Hercynine (5) is considered as the primer for ergothioneine biosynthesis, which is a secreted antioxidant that protects cells from oxidative stress.<sup>46</sup> Fumimycin (7), an unusual metabolite incorporating an unusual alanine unit linked to a phenyl group at the alpha-carbon with both lactone and amide moieties, was first isolated from *Aspergillus fumisynnematus*, and it was detected in the mycelia, primordia and stroma of *C. cicadae*.<sup>5</sup> It showed antibacterial activity against *Staphylococcus aureus* with an IC<sub>50</sub> value of 4.1  $\mu$ M, inhibiting peptide deformylase.<sup>47</sup>

Cordysin A (8), a diketopiperazine first identified from *C. sinensis*,<sup>48,49</sup> was detected in the mycelia of *C. cicadae*.<sup>5</sup> It shows activity in inhibiting the proliferation of human glioma U87-MG and U251 cells.<sup>50</sup> It has anti-inflammatory activity in human neutrophils, which was assessed by its inhibition of FMLP/CB-induced superoxide anion generation (<https://pubchem.ncbi.nlm.nih.gov>).

Some of the SMSs in *I. cicadae* are toxic to animals and humans. Among them, five NRPs are common mycotoxins. Beauvericin (9) and its analogues A, E, J (10–12), and cordycecin A (13), were isolated from its fruit body,<sup>51</sup> and beauvericin was also detected in other stages (mycelia, primordia and stroma).<sup>5</sup> The four beauvericins (9–12) exhibited a significant inhibitory effect on HepG2 and HepG2/ADM cells with IC<sub>50</sub> values ranging from 2.40–14.48  $\mu$ M. However, cordycecin A (13) had no such activity.<sup>51</sup> Beauvericins have insecticidal effects at a microgram level to several insects.<sup>52</sup> The cytotoxicity of beauvericins on human cells and cancer cells was also discovered.<sup>53–55</sup> Thus, the risks of beauvericins to contaminate foods and influence human health are attracting the attention of researchers.<sup>56,57</sup> Aspergilliamide (14) is also an amino derivative, which was detected in the primordia of *I. cicadae*.<sup>5</sup> It was toxic to brine shrimp with an LC<sub>50</sub> value of 71.09 nM.<sup>58</sup>

Myriocin (15) (ISP-1, thermozymocidin), an atypical amino acid, was isolated from the culture broth, mycelia and sporoderm-broken spore powders of *C. cicadae*<sup>27,59–61</sup> and from the culture broth of the *I. sinclairii* ATCC24400 strain.<sup>62</sup> In fact, it was named myriocin because it was first identified from the thermophilic ascomycete *Myriococcum albomyces*. It exhibited good anti-fungal activity against *Candida albicans*. However, the








Fig. 2 Structures of the SMs isolated from *I. cicadae*.

acute toxicity of myriocin is too high for therapeutic purposes. It had intraperitoneal toxicity with an  $LD_{50}$  of 5–10 and 2–5  $mg\ kg^{-1}$  for mice and rats, respectively, and resulted in the death of dogs 48–72 hours after they were treated by subcutaneous injection of 0.25  $mg\ kg^{-1}$ .<sup>63</sup> Therefore, myriocin did not attract more attention until it was found in the culture broth of the

strain CTCC 24400 of *I. sinclairii* and exhibited 10- to 100-fold more potency than cyclosporin A as an immunosuppressive agent of the immune response *in vitro* and *in vivo*.<sup>64</sup> In 1995, a novel immunosuppressant, FTY720 (fingolimod), was first synthesized and screened based on the knowledge that 2-substituted 2-aminoethanol is the minimum essential structure



for the immunosuppression activity of myriocin.<sup>65,66</sup> Different from the action mechanism of myriocin, which inhibits serine palmitoyl transferase, FTY720 is a sphingosine 1-phosphate (S1P) receptor modulator, which is mainly phosphorylated by sphingosine kinase 2 *in vivo*. The phosphorylated drug (-P) acts as a potent agonist of four of the five G protein-coupled receptors for S1P: S1P(1), S1P(3), S1P(4), and S1P(5).<sup>67</sup> FTY720 is now used worldwide for the treatment of multiple sclerosis.<sup>68</sup>

Phytosphingosine (16), a myriocin-like long-chain compound, was also found in the mycelia, primordia and stroma of *C. cicadae*.<sup>5</sup> In the budding yeast *Saccharomyces cerevisiae*, the endoplasmic reticulum stress surveillance pathway is activated by phytosphingosine, which ensures that daughter cells inherit a functional ER.<sup>69</sup>

Oosporein (17), an important dibenzoquinone pigment identified early,<sup>70</sup> was also detected in the mycelia, primordia and stroma of *C. cicadae*.<sup>5</sup> It exhibited median oral toxicity to 1 day-old cockerels.<sup>71</sup> Oosporein inhibits total erythrocyte membrane ATPase activity in a dose-dependent manner due to alterations in erythrocyte morphology and promotes varying degrees of cell lysis.<sup>72</sup> Oosporein also exhibits broad spectrum *in vitro* antimicrobial, antioxidant and cytotoxic activities.<sup>73</sup> Oosporein is a rather strong organic acid and can hardly be adsorbed by organisms; thus, oosporein is unlikely to enter the food chain and influence human health.<sup>74</sup> Also, this fungus produces other pigments and is considered as a resource of natural color materials for the food industry.<sup>75,76</sup> However, the exact components in its pigments are not clear.

Stipitatic acid (18), a tropolone compound isolated in early 1959 from *Penicillium stipitatum*, was also detected in the mycelia, primordia and stroma of *C. cicadae*.<sup>5</sup> Stipitatic acid shows bactericidal and insecticidal activities, as well as growth-inhibitory effects on some pathogenic fungi, especially *Pythium aphanidermatum*.<sup>77</sup>

Many SMs are unknown, and their bioactivity, toxicity, and even molecular structure are unclear. For example, twelve unknown compounds were detected in *C. cicadae* by Lu *et al.*<sup>5</sup> Meanwhile, cycloheximide acid A (19), 2-(3-carboxy-2-hydroxypropyl)-3-methyl-2-cyclopentenone (20), 5-(2-hydroxyethyl)-2-furanacetic acid (21) and lichenicolin A (22) were identified in different stages of *C. cicadae*,<sup>78</sup> but their bioactivities are not yet known.

Moreover, some new compounds were found in fungal hybrids. A strain from the cell fusion of *C. cicadae* and *C. militaris* produces different SMs including two new compounds, 2-(5-(3-oxobutyl) furan-2-yl) acetic acid (23) and cordycepon (24), as well as eleven known compounds, cephalosporolide E (25), cephalosporolide J (26), methyl 2-(5-(3-hydroxybutyl)furan-2-yl) acetate (27), 2-(5-(3-hydroxybutyl) furan-2-yl) acetic acid (28),  $\alpha$ -furoic acid (29), (22*E*,24*R*)-ergosta-5,7,22-trien-3 $\beta$ -ol (ergosterol) (30), 3-benzyl-6-isopropyl-2,5-piperazinedione (31), and 3-isobutyl-6-isobutylpiperazine-2,5-dione (32).<sup>79</sup> The AChE inhibitory activity of all the compounds was tested, and the results showed that compounds 23 and 27–29 at a concentration of 50  $\mu\text{g mL}^{-1}$  showed weak inhibitory activity against AChE with inhibition rates of 15.45%, 17.89%, 15.07%, and 16.41%, respectively. Moreover, the bioassay for the anti-nematode

activity of all the isolates showed that compounds 29 and 31 exhibited moderate inhibitory activity against the nematode *Panagrellus redivivus* with mortality ratios of 71.66% and 72.29% at 2.5  $\text{mg mL}^{-1}$ , respectively.<sup>79</sup>

### 3 SMs from *I. tenuipes*

More than twenty SMs were isolated from *I. tenuipes*, and most of them belong to compounds from the PKS pathway (Table 2 and Fig. 3). Cephalosporolides B (33), F (34) and tenuipyron (35) were isolated in 2012 by Asai *et al.*<sup>80</sup> Cephalosporolide B is thought to be a true biosynthetic precursor of tenuipyron,<sup>80</sup> which can be used as a precursor for the chemical synthesis of cephalosporin C (36), G and (4-OMe-) G.<sup>81</sup> Cephalosporolide F (34) showed moderate inhibitory activity against the nematode *Panagrellus redivivus* with a mortality ratio of 79.0% at 2.5  $\text{mg mL}^{-1}$ .<sup>79</sup>

The ten-membered macrolide, (4*R*,10*R*)-4-hydroxy-10-methyloctane-2,8-dione (37) and a novel diketopiperazine, (3*S*)-6-benzyl-3-isopropyl-1-methylpiperazine-2,5-dione (38), were obtained from the culture broth.<sup>82</sup> They showed moderate cytotoxicity in the MTT assay against prostate cancer cells 22RV1 and DU-145, and their inhibition rates were 37.8% and 38.6% and 32.5% and 40.6%, respectively.

The trichothecene derivative, 4- $\beta$ -acetoxyscirpindiol (4-ASD) (39) was isolated from the fruit body in 2001, which induced apoptosis in human leukemia cells (HL-60).<sup>83</sup> It lowered the blood sugar levels in the circulatory system by inhibiting Na<sup>+</sup>/glucose transporter-1 (SGLT-1)<sup>84</sup> and also showed significant apoptosis-inducing activity in various human cancer cell lines.<sup>85</sup>

Tenuipesine A (40) and paecilomycines A–C (41–43) were isolated from the cultivated fruiting bodies also, but only paecilomycine A (41) exhibited activity in neurotrophic factor biosynthesis in glial cells.<sup>86,87</sup> The spirocyclic trichothecanes, spirotenuipesines A (44) and B (45), were isolated from fruiting body grown in barley grain. The two compounds had the potent activity in neurotrophic factor biosynthesis in glial cells.<sup>88</sup> As is known, trichothecenes are very important mycotoxins with multiple toxicities including carcinogenic, teratogenic and mutagenic effects.<sup>89</sup>

Cyclopenta benzopyran and penostatins A–C (46–48) and J (49) were isolated in 2014 by Chen *et al.* All of them are protein tyrosine phosphatase 1B (PTP1B) inhibitors.<sup>90</sup> In fact, penostatins A–C were identified in a strain of *Penicillium* sp. originally separated from the marine alga *Enteromorpha intestinalis*. They exhibited significant cytotoxicity against cultured P388 cells.<sup>91</sup>

*I. tenuipes* produces isariotins, which are alkaloids probably biosynthesized through the hybrid NRPS-PKS pathway. Isariotins A–D (50–53) were isolated from the culture broth of *I. tenuipes* BCC7831 strain in 2007 by Haritakun *et al.* These compounds showed no bioactivity against malaria parasites and fungi, and no cytotoxicity against three cancer cell lines and Vero cells.<sup>92</sup> Isariotins E (54), F (55) and TK-57-164A (56) were isolated from the strain BCC12625 in 2009 by Bunyapaiboonsri *et al.*<sup>93,94</sup> Isariotin F (55) exhibited activity against the malaria parasite *Plasmodium falciparum* K1 with an IC<sub>50</sub> value of 5.1 nM and cytotoxic activities against cancer cell lines (KB, BC, and



Table 2 SMS isolated from *Isaria tenuipes* and their biological activities

Metabolites	CAS no.	Material source	Biological activity
Cephalosporolide B (33)	97344-03-1	Culture broth	Moderate inhibitory activity against the nematode <i>Panagrellus redivivus</i>
Cephalosporolide F (34)	97344-04-2	Culture broth of <i>I. tenuipes</i>	
Tenuipyrone (35)	1354559-27-5	Culture broth of <i>I. tenuipes</i>	Inhibit prostate cancer cells 22RV1 and DU-145 at the rates of 32.5% and 40.6% at 5 $\mu\text{mol L}^{-1}$ , respectively
Cephalosporolide C (36)	97344-02-0	Culture broth of <i>I. tenuipes</i>	
(4R,10R)-4-Hydroxy-10-methyloxecane-2,8-dione (37)	2098945-28-7	Culture broth of <i>I. tenuipes</i>	
(3S)-6-Benzyl-3-isopropyl-1-methylpiperazine-2,5-dione (38)	2101845-87-6	Culture broth of <i>I. tenuipes</i>	Inhibit prostate cancer cells 22RV1 and DU-145 at the rates of 37.8% and 38.6% at 5 $\mu\text{mol L}^{-1}$ , respectively
4- $\beta$ -Acetoxyscirpendiol (39)	2531-11-5	Cultivated fruiting bodies of <i>I. japonica</i> (voucher no. CHO-6133)	Induces apoptosis of human leukemia cells (HL-60). Lower blood sugar levels in the circulatory system as effective SGLT-1 inhibitors
Tenuipesine A (40)	816448-01-8	Cultivated fruiting bodies of <i>P. tenuipes</i> MH19912	Activity in neurotrophic factor biosynthesis in glial cells
Paecilomycine A (41)	594840-03-6	Cultivated fruiting bodies of <i>P. tenuipes</i> MH19912	
Paecilomycine B (42)	594840-04-7	Cultivated fruiting bodies of <i>P. tenuipes</i> MH19912	
Paecilomycine C (43)	765942-39-0	Cultivated fruiting bodies of <i>P. tenuipes</i> MH19912	Potent activity in neurotrophic factor biosynthesis in glial cells
Spirotenuipesine A (44)	594840-05-8	Cultivated fruiting bodies of <i>P. tenuipes</i> MH19912	
Spirotenuipesine B (45)	594840-06-9	Cultivated fruiting bodies of <i>P. tenuipes</i> MH19912	
Penostatin A (46)	173485-70-6	Mixture of cultivated mycelium and medium of <i>P. tenuipes</i> RCEF 37776	Protein phosphatase 1B inhibitors
Penostatin B (47)	173655-56-6	Mixture of cultivated mycelium and medium of <i>P. tenuipes</i> RCEF 37776	Protein phosphatase 1B inhibitors
Penostatin C (48)	173485-71-7	Mixture of cultivated mycelium and medium of <i>P. tenuipes</i> RCEF 37776	Protein phosphatase 1B inhibitors
Penostatin J (49)	1695557-12-0	Mixture of cultivated mycelium and medium of <i>P. tenuipes</i> RCEF 37776	Protein phosphatase 1B inhibitors
Isariotin A (50)	952703-96-7	Culture broth of <i>I. tenuipes</i> BCC7831	Marginal activity against the <i>Mycobacterium tuberculosis</i> H37Ra with the MIC value of 486 $\mu\text{M}$
Isariotin B (51)	952703-97-8	Culture broth of <i>I. tenuipes</i> BCC7831	Marginal activity against the <i>Mycobacterium tuberculosis</i> H37Ra with the MIC value of 488 $\mu\text{M}$
Isariotin C (52)	952703-98-9	Culture broth of <i>I. tenuipes</i> BCC7831	
Isariotin D (53)	952703-99-0	Culture broth of <i>I. tenuipes</i> BCC7831	
Isariotin E (54)	1137665-60-1	Culture broth of <i>I. tenuipes</i> BCC12625	Antimalarial activity against <i>Plasmodium falciparum</i> K1; Antitubercular activity against <i>Mycobacterium tuberculosis</i> H37Ra; Antifungal activity against <i>Candida albicans</i> ; cytotoxic activities against three cancer cell lines (KB, BC, and NCI-H187)
Isariotin F (55)	1137665-61-2	Culture broth of <i>I. tenuipes</i> BCC12625	
TK-57-164A (60)	745050-50-4	Culture broth of <i>I. tenuipes</i> BCC12625, BCC21283	Antimalarial activity; cytotoxic activities against KB, MCF-7, NCI-H187 and Vero cells



Table 2 (Contd.)

Metabolites	CAS no.	Material source	Biological activity
Isariotin G (56)	1383930-91-3	Culture broth of <i>I. tenuipes</i> BCC21283	Antimalarial activity; cytotoxic activities against KB, MCF-7, NCI-H187 and Vero cells
Isariotin H (57)	1383930-93-5	Culture broth of <i>I. tenuipes</i> BCC15621	Antimalarial activity; cytotoxic activities against KB, MCF-7, NCI-H187 and Vero cells
Isariotin I (58)	1383930-95-7	Culture broth of <i>I. tenuipes</i> BCC15621	Antimalarial activity; cytotoxic activities against KB, MCF-7, NCI-H187 and Vero cells
Isariotin J (59)	1383930-97-9	Culture broth of <i>I. tenuipes</i> BCC15621	Antimalarial activity; cytotoxic activities against KB, MCF-7, NCI-H187 and Vero cells
Hanasanagin (61)	1187336-16-8	Fruiting body of <i>I. japonica</i> cultivated on silkworm pupae	Antioxidant activity
3'-Deoxyhanasanagin(3,4-diguanidinobutanoyl-tyrosine) (62)	1187059-33-1	Fruiting body of <i>I. japonica</i> cultivated on silkworm pupae	

NCI-H187) and nonmalignant (Vero) cells with IC<sub>50</sub> values of 15.8, 2.4, 1.6, and 2.9 nM, respectively. However, TK-57-164A (56) exhibit no such activity.<sup>93</sup> Both isariotin G (57) and TK-57-164A (56) were isolated from the strain BCC21283, while isariotins H–J (58–60) were isolated from the strain BCC15621<sup>95</sup>. Isariotins G (57), I (59) and J (60) showed antimalarial activity with IC<sub>50</sub> values of 2.10–5.51 µg mL<sup>-1</sup>, while isariotin H (58) had IC<sub>50</sub> values of >10 µg mL<sup>-1</sup>. Isariotins G–J (57–60) all exhibited cytotoxic activities against KB, MCF-7, NCI-H187 and Vero cells.<sup>95</sup>

Two pseudo-di-peptides, hanasanagin (61) (3,4-diguanidinobutanoyl-DOPA) and 3,4-diguanidinobutanoyl-tyrosine (62), were isolated from the fruiting bodies of *I. japonica* cultivated on silkworm pupae.<sup>96,97</sup> Hanasanagin (61) exhibited antioxidant activity, but 3,4-diguanidinobutanoyl-tyrosine (62) showed no antioxidant activity.<sup>96,97</sup>

In addition, *I. tenuipes* produces the depsipeptide beauvericin (9). It was confirmed that beauvericin is one of the active principles of three strains of *I. tenuipes*, which strongly inhibited mycelial growth of the two phytopathogens *Phytophthora sojae* and *Aphanomyces cochlioides*.<sup>98</sup>

## 4 Problems and perspectives

In recent years, the concerns about the safety risks of medical and edible *Isaria* fungi have been attracting the attention of researchers and consumers. In fact, several research reports about the biosafety analysis of both fungi were published.

For *I. cicadae*, there were four reports indicating its safety to humans, but one report demonstrated its toxicity to the kidneys (Table 3). The possible toxicity arising from repeated exposure to freeze-dried submerged mycelial culture of *C. cicadae* for 90 days was evaluated. The results indicated that there were no adverse effects to male and female Sprague Dawley rats by gavage >2000 mg kg<sup>-1</sup> *C. cicadae* whole broth.<sup>99</sup> Also, toxicological effects were not recorded in the adult Sprague Dawley rats administered orally with a complex powder suspension of

*P. sinclairii* and larvae of *Bombyx mori* at doses ranging from 0.008 to 5 g kg<sup>-1</sup> body weight for 2 weeks, except for a decline in the weight of the thymus in males.<sup>100</sup> The extract of wild *C. cicadae* fruit bodies was not toxic to mice even at a dosage of 80 g kg<sup>-1</sup>, which was 444 times the clinical daily dosage.<sup>101</sup> Moreover, the genotoxicity of the fruiting bodies of *I. sinclairii* together with its parasitic host larva was evaluated by using short-term genotoxicity tests, namely, the Ames, chromosome aberration (CA), and micronuclei (MN) tests. The results indicated that this complex has no mutagenic potential in the *in vitro* and *in vivo* systems.<sup>102</sup> However, concerns about the nephrotoxicity of the complex of *P. sinclairii* and its host insect (*Bombyx mori*) have been raised. Kidney toxicity was investigated after 13 weeks of administering the complex orally to rats. Dose-dependent kidney cell karyomegaly and tubular hypertrophy were observed. There was a dose-dependent increase in kidney injury molecule 1 (KIM-1) and matrix metalloproteinase 1 (TIMP-1) levels in the kidney and urinary KIM-1, cystatin C, beta-2-microglobulin, and osteopontin levels. KIM-1 and TIMP-1 increased in the male kidneys, and they did not recover 2 weeks after stopping exposure. Cystatin C in the kidney was significantly lowered in all treatment groups at 13 weeks of administration. All the changes were more noticeable in males.<sup>14</sup> In addition, some clinical cases of poisoning by eating wild the complex of *C. cicadae* and the host insect (*Cicadae flammata*) were reported in China.<sup>103</sup>

For *I. tenuipes*, there were two different research reports about its biosafety (Table 3). In the first report, the genotoxicity, acute and subchronic toxicity of the water-macerated extracts of its fungus fruiting bodies were evaluated in Korea.<sup>104</sup> The acute oral LD<sub>50</sub> to rats was >2000 mg kg<sup>-1</sup> of body weight. In the subchronic test, the oral treatment of rats with 500, 1000 or 2000 mg kg<sup>-1</sup> extract daily for 13 weeks did not induce any dose-related changes (body weight, food consumption, clinical observation, urinalysis, hematology, clinical chemistry and organ weight). However, histopathological observation revealed that the *I. tenuipes* extract induced karyomegaly in the outer








Fig. 3 Structures of SMs isolated from *I. tenuipes*.

medulla of the kidneys of all the treated rats. Importantly, the *I. tenuipes* extract exerted mutagenic potential in the Ames assay. Since karyomegalic alterations are known to be associated with carcinogenicity, *I. tenuipes* probably has the risk of carcinogenicity.<sup>104</sup> In the second experiment, the aqueous and ethanol extracts of *I. tenuipes* N45 caused neither mortality nor toxicological signs in mice and rats with the maximum tolerance dose of 15 g kg<sup>-1</sup>. No mortality or adverse effects was observed in the subchronic toxicity tests, in which no significant difference in bodyweight, relative organ weight or hematological parameters,

and no abnormality of internal organs were found between the treatment and control groups. This suggests that the fungus extract is safe.<sup>105</sup>

Obviously, there are contradictions in these research reports, but some safety risks indeed exist in the both fungi. Of course, the toxicity is based on the substances in the fungus or the complex of the fungus and its insect hosts. Thus, from rational deduction, the toxicities of both *Isaria* fungi to humans may be related to the toxic MSs described above; however, this needs to be further validated because other toxic substances including



Table 3 Toxicity tests of the two *Isaria* fungi

Material source	Toxicity descript
Freeze-dried powder of <i>C. cicadae</i> (MU30106, Bioresource Collection and Research Center at the Food Industry Research and Development Institute, Taiwan, China) submerged mycelial culture	A total of eighty 8 week-old Sprague Dawley rats were divided into 4 groups (10 males and 10 females in each group). <i>C. cicadae</i> was administered daily to the animals by gavage at doses of 0, 500, 1000, and 2000 mg kg <sup>-1</sup> body weight for 90 days. No observed adverse effect level to both sexes of Sprague Dawley rats exposed dietary >2000 mg kg <sup>-1</sup> for 90 days
Water extract from the powder of wild <i>C. cicadae</i> fruit bodies and hosts (Hangzhou, China)	No acute toxicity to mice even at a dosage 80 g kg <sup>-1</sup> for 7 days, which was 444 times the clinical daily dosage
Water extract from the powder of <i>P. sinclairii</i> from the National Institute of Agricultural Science and Technology (Suwon, Korea)	No toxicological effects to the adult Sprague Dawley rats orally treated at doses of 0.008–5 g kg <sup>-1</sup> body weight for 2 weeks, except for a decline in the weight of the thymus in the males
Extract of phosphate buffered saline (PBS) from the complex of the fruiting bodies of <i>I. sinclairii</i> and its parasitic host larva (Rural Development Administration, Korea)	In the Ames test, no mutagenic response in the absence or presence of 59 mix with TA98, TA100, TA1535, and TA1537. In the chromosome aberration (CA) test, no significant effect on Chinese hamster ovary (CHO) cells. In the micronuclei (MN) test, no significant change in the occurrence of micronucleated polychromatic erythrocytes in male ICR mice intraperitoneally administered at doses of 15, 150, or 1500 mg kg <sup>-1</sup>
Powder of the cultured fruiting bodies of <i>P. sinclairii</i> (isolated from parasitizing dead or living <i>Cicadae</i> subspecies) and its silk worm larvae hosts (Rural Development Administration, Korea)	Kidney toxicity was investigated after 13 weeks of administering the complex orally to Sprague Dawley rats at doses of 0, 5000, 10 000, and 50 000 ppm. Dose-dependent kidney cell karyomegaly and tubular hypertrophy were observed, with higher severity in males. There was a dose-dependent increase in kidney injury molecule 1 (KIM-1) and tissue inhibitor of matrix metalloproteinase 1 (TIMP-1) levels in the kidney and urinary KIM-1, cystatin C, b2m, and osteopontin levels. KIM-1 and TIMP-1 increased in the male kidneys, and they did not recover 2 weeks after stopping exposure. Cystatin C in the kidney was significantly lowered in all treatment groups at 13 weeks of administration. All the changes were more noticeable in males. These data indicate that the complex damages renal tubule cells with histopathological lesions and changes in biomarker levels. Kidney and urinary KIM-1 and cystatin C were the most markedly affected and early increased indicators among the biomarkers tested; whereas, blood urea nitrogen (BUN) and creatinine were not affected
Aqueous and ethanol extracts of <i>I. tenuipes</i> N45 (CCTCC M2011145) ( <i>P. tenuipes</i> RCEF 4339, Anhui Agricultural University, China)	In acute toxicity, neither mortality nor toxicological signs were found in mice and rats with the maximum tolerance dose of 15 g kg <sup>-1</sup> . No mortality or adverse effects was observed in the subchronic toxicity studies, in which no significant difference in bodyweight, relative organ weight or hematological parameters, and no abnormality of internal organs were found between the treatment and control groups
Water extract from the powder of fruiting bodies of <i>P. tenuipes</i> (Korea Food Research Institute, Seongnam, Korea)	The acute oral LD <sub>50</sub> to Sprague Dawley rats was estimated to be greater than 2000 mg kg <sup>-1</sup> of body weight. In the subchronic study, the oral treatment of rats with 500, 1000 or 2000 mg kg <sup>-1</sup> daily for 13 weeks did not induce any dose-related changes (body weight, food consumption, clinical observation, urinalysis, hematology, clinical chemistry and organ weight). In contrast, histopathological observation revealed that the extract induced karyomegaly in the outer medulla of the kidney in all the treated rats. In the Ames tests, the extract significantly produced His <sup>+</sup> mutants at the highest concentration of 5000 lg per plate in <i>S. typhimurium</i> TA102 and 1535 without S9 mix. In the presence of S9 mix, it also significantly produced His <sup>+</sup> mutants in TA98, 100 and 102 at 5000 lg per plate of the extract. The increase in His <sup>+</sup> mutants was dose-dependent although it was less than 2-fold in comparison to that of the control group

mycotoxins and macro biological molecules in both fungi may not have been discovered yet. Also, their toxicity is related to the genetic backgrounds of the fungal strains and the hosts, and their growing environment and culture conditions.<sup>106</sup> In fact, different fungal strains under the same culture conditions or the same strain under different culture conditions, usually produce different toxic substances.<sup>107,108</sup> Therefore, controlling

the culturing conditions and growth environments of *Isaria* fungi is very important for the quality and safety management of their fungal products.

To date, the products of *I. cicadae* and *I. tenuipes* have no standards for quality and safety control. Referring to *C. sinensis* in the Chinese Pharmacopoeia, 2015, the adenosine marker is the key parameter for quality control. Cordycepin and



cordyceps acid are considered as the bioactive components.<sup>29</sup> Their other functional compounds such as nucleosides, amino acids, fatty acids and carbohydrates are attracting researchers attention and certain detection methods have been developed.<sup>31,109</sup> However, to absolutely ensure the safety of the products of both fungi, only detecting their functional components is not sufficient. Two other aspects must be emphasized. First, identification standards for both fungi and hosts must be developed. As it is known, distinguishing fungal species based on their morphological features is very difficult, especially in commercially processed products. A good choice is may be methods based on DNA. In fact, DNA barcode-based species-specific sequence characterized amplified region (SCAR) markers to discriminate authentic herbal *Cordyceps* medicines and *Cordyceps*-derived dietary supplements from related but inauthentic species were recently reported. The ITS-based SCAR markers and the real-time PCR assay constitute a useful genetic tool for preventing the adulteration of *Cordyceps* and *Cordyceps*-related dietary supplements.<sup>110</sup> Second, the accurate detection and differentiation of toxic components (mycotoxins) in the products of both fungi must be established. Furthermore, residue standards of their toxic components should be developed. In recent years, metabonomics based on GC/HPLC-MS has been used for the analysis of *Cordyceps* compounds<sup>5,28,111</sup> and other effective methods such as HPLC fingerprint for functional substances have also been developed.<sup>31,109,112</sup> These technologies will support the detection and identification of SMs in the both *Isaria* fungi and their products.

In conclusion, both *I. cicadae* and *I. tenuipes* are important edible and medicinal fungal species with multiple pharmacological functions. They produce various SMs that affect the quality and safety of their fungal products. For *I. cicadae*, its SMs include nucleosides, amino acids, beauvericins, and myriocin, oosporein. For *I. tenuipes*, trichothecene derivatives, isariotins, cyclopenta benzopyrans and PKs were found. Among them, beauvericins, myriocin, oosporein and many trichothecene derivatives are toxic compounds. Thus, the toxicity and side-effects of the fungal products of *I. cicadae* and *I. tenuipes* may be related to these SMs. To ensure the quality and safety of their fungal products, residues standards for their SMs must be developed. Furthermore, methods for SM detection and strain biological identification must be give attention.

## Conflicts of interest

The authors declare no conflict of interests.

## Abbreviations

SMs	Secondary metabolites
NRPS	Non-ribosomal peptide synthetase
PKS	Polyketide synthase
TS	Terpenoid synthetase
PK	Polyketides
ACH	Acetylcholine esterase
PTP1B	Protein tyrosine phosphatase 1B

MN  
PCR

Micronuclei  
Polymerase chain reaction

## Acknowledgements

This research is supported by The National Key Research and Development Program of China (2016YFD0200506) and National Natural Science Foundation of China (31572053). We cordially thank Dr Sengodan Karthi for his carefully linguistic modification to this article.

## References

- 1 J. J. Luangsa-Ard, N. L. Hywel-Jones, L. Manoch and R. A. Samson, *Mycol. Res.*, 2005, **109**, 581–589.
- 2 C. Dong, S. Guo, W. Wang and X. Liu, *Mycology*, 2015, **6**, 121–129.
- 3 J. Holliday and M. Cleaver, *Int. J. Med. Mushrooms*, 2008, **10**, 219–234.
- 4 K. Yue, M. Ye, Z. J. Zhou, W. Sun and X. Lin, *J. Pharm. Pharmacol.*, 2013, **65**, 474–493.
- 5 Y. Lu, F. Luo, K. Cen, Y. Yin, S. Zhan, C. Wang, H. Zhang, G. Xiao, C. Li and Z. Li, *BMC Genomics*, 2017, **18**, DOI: 10.1186/s12864-017-4060-4.
- 6 E. Yokoyama, M. Arakawa, K. Yamagishi and A. Hara, *FEMS Microbiol. Lett.*, 2006, **264**, 182–191.
- 7 F. Takano, N. Yahagi, R. Yahagi, S. Takada, M. Yamaguchi, S. Shoda, T. Murase, S. Fushiya and T. Ohta, *Int. Immunopharmacol.*, 2005, **5**, 903–916.
- 8 B. Song, Q. Liu, T. Lin, Y. Shen, J. Li and D. Luo, *J. Fungal Res.*, 2006, **4**, 10–25.
- 9 C. Dong, W. Li, Z. Li, W. Yan, T. Li and X. Liu, *Mycosystema*, 2016, **35**, 1–15.
- 10 G. H. Cunningham, *Trans. R. Soc. N. Z.*, 1921, **53**, 372–382.
- 11 R. T. Duarte, K. C. Goncalves, D. J. Espinosa, L. F. Moreira, S. A. De Bortoli, R. A. Humber and R. A. Polanczyk, *J. Econ. Entomol.*, 2016, **109**, 594–601.
- 12 Z. Li, Z. Chen and Y. Chen, *Guobao Chongcao Jingchuanhua*, Hefei University of Technology Publishing House, China, 1st edn, 2014.
- 13 M. Y. Ahn, Y. S. Jung, S. D. Jee, C. S. Kim, S. H. Lee, C. H. Moon, S. I. Cho, B. M. Lee and K. S. Ryu, *Arch. Pharmacol. Res.*, 2007, **30**, 493–501.
- 14 M. Jeong, Y. W. Kim, J. R. Min, M. Kwon, B. S. Han, J. G. Kim and S. H. Jeong, *Food Chem. Toxicol.*, 2013, **59**, 177–186.
- 15 M. Jeong, Y.-W. Kim, J.-R. Min, M. Kwon, B.-S. Han, J.-G. Kim and S.-H. Jeong, *Toxicol. Res.*, 2012, **28**, 179–185.
- 16 P. Vega-Aquino, S. Sanchez-Pena and C. A. Blanco, *J. Invertebr. Pathol.*, 2010, **103**, 145–149.
- 17 Y. Han, J. Liang, X. Zou, X. Dong and Z. Liang, *Acta Edulis Fungi*, 2011, **18**, 89–94.
- 18 K. H. Shin, S. S. Lim, S. Lee, Y. S. Lee, S. H. Jung and S. Y. Cho, *Phytother. Res.*, 2003, **17**, 830–833.
- 19 J. Liang, Y. Han, W. Tian, Z. Liang and Z. Li, *Acta Edulis Fungi*, 2018, **25**, 113–119.



- 20 S. S. Sum and J. Ziegler, *Nutr. Clin. Pract.*, 2016, **31**, 695–697.
- 21 X. Wang, X. Gong, P. Li, D. Lai and L. Zhou, *Molecules*, 2018, **23**, 169, DOI: 10.3390/molecules23010169.
- 22 K. Zhou, X. Zhang, F. Zhang and Z. Li, *Microb. Ecol.*, 2011, **62**, 644–654.
- 23 J.-H. Hsu, B.-Y. Jhou, S.-H. Yeh, Y.-L. Chen and C.-C. Chen, *J. Nutr. Food Sci.*, 2015, **5**, 432.
- 24 S.-C. Weng, C.-J. Chou, L.-C. Lin, W.-J. Tsai and Y.-C. Kuo, *J. Ethnopharmacol.*, 2002, **83**, 79–85.
- 25 Y. Shen, M. Cai, W. Xia, J. Liu, Q. Zhang, H. Xie, C. Wang, X. Wang and S. Zheng, *Cancer Lett.*, 2007, **254**, 288–297.
- 26 K. Su, P. Zeng, W. Liang, Z. Luo, Y. Wang, X. Lv, Q. Han, M. Yan and C. Chen, *Mediators Inflammation*, 2017, 3701385, DOI: 10.1155/2017/3701385.
- 27 Y. Sun, M. Wink, P. Wang, H. Lu, H. Zhao, H. Liu, S. Wang, Y. Sun and Z. Liang, *Phytomedicine*, 2017, **36**, 217–228.
- 28 J. Zhang, H. Yu, X. Zhong, G. Zhang and X. Liu, *Chin. J. Exp. Tradit. Med. Formulae*, 2018, **24**, 23–29.
- 29 X. Zhou, Z. Gong, Y. Su, J. Lin and K. Tang, *J. Pharm. Pharmacol.*, 2009, **61**, 279–291.
- 30 L. He, S. Ma, J. Cheng, W. Li and X. Wu, *Shipin Yu Shengwu Jishu Xuebao*, 2012, **31**, 8–16.
- 31 L. Li, H. Liang, Z. Wu, Z. Zhang, L. Zhang, Y. Wang, T. Zhang and B. Wang, *Lishizhen Med. Mater. Med. Res.*, 2017, **28**, 1537–1542.
- 32 Y. Wang, Y. Guo, L. Zhang and J. Wu, *Braz. J. Microbiol.*, 2012, **43**, 449–455.
- 33 M. Siev, R. Weinberg and S. Penman, *J. Cell Biol.*, 1969, **41**, 510–520.
- 34 O. J. Olatunji, Y. Feng, J. Tang, O. O. Olatunji, Z. Ouyang and Z. Su, *Biomed. Pharmacother.*, 2016, **81**, 7–14.
- 35 K. Nakamura, K. Shinozuka and N. Yoshikawa, *J. Pharmacol. Sci.*, 2015, **127**, 53–56.
- 36 H. S. Tuli, A. K. Sharma, S. S. Sandhu and D. Kashyap, *Life Sci.*, 2013, **93**, 863–869.
- 37 B. Li, Y. Hou, M. Zhu, H. Bao, J. Nie, G. Y. Zhang, L. Shan, Y. Yao, K. Du, H. Yang, M. Li, B. Zheng, X. Xu, C. Xiao and J. Du, *Int. J. Neuropsychopharmacol.*, 2016, **19**, DOI: 10.1093/ijnp/pyv112.
- 38 Y. C. Chen, Y. H. Chen, B. S. Pan, M. M. Chang and B. M. Huang, *J. Food Drug Anal.*, 2017, **25**, 197–205.
- 39 Z. Wang, X. Zhao, F. Gao, Z. Zhao and K. Wang, *Zhong Xi Yi Jiehe Yanjiu*, 2017, **9**, 169–171.
- 40 Z. Wang, X. Zhao, L. Song, F. Gao, Z. Zhao and K. Wang, *World Journal of Integrated Traditional and Western Medicine*, 2018, **13**, 1306–1309.
- 41 H. C. Kuo, I. C. Huang and T. Y. Chen, *Int. J. Med. Mushrooms*, 2015, **17**, 1077–1085.
- 42 M. Y. Ahn, J. E. Heo, J.-H. Ryu, S. D. Ji, H. Jeong, H. C. Park and H. S. Sim, *Int. J. Indust. Entomol.*, 2008, **17**, 197–200.
- 43 S. Lin, Z. Q. Liu, Y. P. Xue, P. J. Baker, H. Wu, F. Xu, Y. Teng, M. E. Brathwaite and Y. G. Zheng, *Appl. Biochem. Biotechnol.*, 2016, **179**, 633–649.
- 44 M. Saeed, D. Babazadeh, M. Naveed, M. A. Arain, F. U. Hassan and S. Chao, *Trop. Anim. Health Prod.*, 2017, **49**, 1329–1338.
- 45 S.-Y. Chen, K.-J. Ho, Y.-J. Hsieh, L.-T. Wang and J.-L. Mau, *LWT-Food Sci. Technol.*, 2012, **47**, 274–278.
- 46 J. H. Jeong, H. J. Cha, S. C. Ha, C. Rojviriyaya and Y. G. Kim, *Biochem. Biophys. Res. Commun.*, 2014, **452**, 1098–1103.
- 47 Y. J. Kwon, M. J. Sohn, C. J. Zheng and W. G. Kim, *Org. Lett.*, 2007, **9**, 2449–2451.
- 48 P. X. Chen, S. A. Wang, S. P. Nie and M. Marcone, *J. Funct. Foods*, 2013, **5**, 550–569.
- 49 M. L. Yang, P. C. Kuo, T. L. Hwang and T. S. Wu, *J. Nat. Prod.*, 2011, **74**, 1996–2000.
- 50 X. W. Ye, W. Y. Chai, X. Y. Lian and Z. Z. Zhang, *Nat. Prod. Res.*, 2017, **31**, 1390–1396.
- 51 J. Wang, D.-M. Zhang, J.-F. Jia, Q.-L. Peng, H.-Y. Tian, L. Wang and W.-C. Ye, *Fitoterapia*, 2014, **97**, 23–27.
- 52 Q. G. Wang and L. J. Xu, *Molecules*, 2012, **17**, 2367–2377.
- 53 Y. W. Tao, Y. C. Lin, Z. G. She, M. T. Lin, P. X. Chen and J. Y. Zhang, *Anti-Cancer Agents Med. Chem.*, 2015, **15**, 258–266.
- 54 W. Watjen, A. Debbab, A. Hohlfeld, Y. Chovolou and P. Proksch, *Toxicol. Lett.*, 2014, **231**, 9–16.
- 55 X. F. Wu, R. Xu, Z. J. Ouyang, C. Qian, Y. Shen, X. D. Wu, Y. H. Gu, Q. Xu and Y. Sun, *PLoS One*, 2013, **8**, e83013, DOI: 10.1371/journal.pone.0083013.
- 56 M. J. Ruiz, P. Franzova, A. Juan-Garcia and G. Font, *Toxicon*, 2011, **58**, 315–326.
- 57 C. Luz, F. Saladino, F. B. Luciano, J. Manes and G. Meca, *Food Chem. Toxicol.*, 2017, **107**, 430–439.
- 58 X. L. Xu, L. Y. Yin, S. Y. S. Wang, H. H. Liu, J. H. Gao and S. J. Zhao, *Rec. Nat. Prod.*, 2013, **7**, 292–295.
- 59 J. Yu, Z. Mo, X. Mao, H. Xu and H. Zhu, *Yaowu Fenxi Zazhi*, 2010, **30**, 664–667.
- 60 J. Yu, H. Xu, Z. Mo, H. Zhu and X. Mao, *Anal. Sci.*, 2009, **25**, 855–859.
- 61 H. Xu, Z. Mo, J. Yu, X. Mao and H. Zhu, *Nat. Prod. Res. Dev.*, 2010, **22**(5), 794–797.
- 62 T. Fujita, K. Inoue, S. Yamamoto, T. Ikumoto, S. Sasaki, R. Toyama, K. Chiba, Y. Hoshino and T. Okumoto, *J. Antibiot.*, 1994, **47**, 208–215.
- 63 D. Kluepfel, J. Bagli, H. Baker, M. P. Charest and A. Kudelski, *J. Antibiot.*, 1972, **25**, 109–115.
- 64 T. Fujita, K. Inoue, S. Yamamoto, T. Ikumoto, S. Sasaki, R. Toyama, K. Chiba, Y. Hoshino and T. Okumoto, *J. Antibiot.*, 1994, **47**, 208–215.
- 65 T. Fujita, R. Hirose, N. Hamamichi, Y. Kitao, S. Sasaki, M. Yoneta and K. Chiba, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 1857–1860.
- 66 K. Adachi, T. Kohara, N. Nakao, M. Arita, K. Chiba, T. Mishina, S. Sasaki and T. Fujita, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 853–856.
- 67 K. Adachi, *J. Synth. Org. Chem., Jpn.*, 2011, **69**, 904–912.
- 68 M. Sanford, *Drugs*, 2014, **74**, 1411–1433.
- 69 F. Pina, F. Yagisawa, K. Obara, J. D. Gregerson, A. Kihara and M. Niwa, *J. Cell Biol.*, 2018, **217**, 495–505.
- 70 H. Takeshita and M. Anchel, *Science*, 1965, **147**, 152–153.
- 71 R. J. Cole, J. W. Kirksey, H. G. Cutler and E. E. Davis, *J. Agric. Food Chem.*, 1974, **22**, 517–520.





- 72 L. B. Jeffs and G. G. Khachatourians, *Toxicon*, 1997, **35**, 1351–1356.
- 73 R. Alurappa, M. R. M. Bojgowda, V. Kumar, N. K. Mallesh and S. Chowdappa, *Nat. Prod. Res.*, 2014, **28**, 2217–2220.
- 74 C. Seger, D. Erlebach, H. Stuppner, U. J. Griesser and H. Strasser, *Helv. Chim. Acta*, 2005, **88**, 802–810.
- 75 J. W. Choi and J. P. Park, *Biotechnol. Bioprocess Eng.*, 2018, **23**, 246–249.
- 76 F. A. E. Torres, B. R. Zaccarim, L. C. D. Novaes, A. F. Jozala, C. A. dos Santos, M. F. S. Teixeira and V. C. Santos-Ebinuma, *Appl. Microbiol. Biotechnol.*, 2016, **100**, 2511–2521.
- 77 R. Bentley, *Nat. Prod. Rep.*, 2008, **25**, 118–138.
- 78 S.-W. Zhang and L.-J. Xuan, *J. Antibiot.*, 2008, **61**, 43–45.
- 79 N. N. Yang, N. Jiang, Q. Y. Ma, F. D. Kong, Q. Y. Xie, L. M. Zhou, Z. F. Yu and Y. X. Zhao, *J. Asian Nat. Prod. Res.*, 2018, DOI: 10.1080/10286020.2018.1451518.
- 80 T. Asai, Y. M. Chung, H. Sakurai, T. Ozeki, F. R. Chang, K. Yamashita and Y. Oshima, *Org. Lett.*, 2012, **14**, 513–515.
- 81 L. Y. Song, Y. Liu and R. B. Tong, *Org. Lett.*, 2013, **15**, 5850–5853.
- 82 Y. Zheng, H. Pang, J. Wang, D. Chen, G. Shi and J. Huang, *Chem. J. Chin. Univ.*, 2012, **17**, 1665–1669.
- 83 G.-S. Oh, K.-H. Hong, H. Oh, H.-O. Pae, I.-K. Kim, N.-Y. Kim, T.-O. Kwon, M.-K. Shin and H.-T. Chung, *Biol. Pharm. Bull.*, 2001, **24**, 785–789.
- 84 O. Yoo, J.-H. Son and D.-H. Lee, *J. Biochem. Mol. Biol.*, 2005, **38**, 211–217.
- 85 K. S. Nam, Y. S. Jo, Y. H. Kim, J. W. Hyun and H. W. Kim, *Life Sci.*, 2001, **69**, 229–237.
- 86 H. Kikuchi, Y. Miyagawa, Y. Sahashi, S. Inatomi, A. Haganuma, N. Nakahata and Y. Oshima, *Tetrahedron Lett.*, 2004, **45**, 6225–6228.
- 87 H. Kikuchi, Y. Miyagawa, K. Nakamura, Y. Sahashi, S. Inatomi and Y. Oshima, *Org. Lett.*, 2004, **6**, 4531–4533.
- 88 H. Kikuchi, Y. Miyagawa, Y. Sahashi, S. Inatomi, A. Haganuma, N. Nakahata and Y. Oshima, *J. Org. Chem.*, 2004, **69**, 352–356.
- 89 S. Aupanun, S. Poapolathep, M. Giorgi, K. Imsilp and A. Poapolathep, *J. Vet. Med. Sci.*, 2017, **79**, 6–13.
- 90 Y.-P. Chen, C.-G. Yang, P.-Y. Wei, L. Li, D.-Q. Luo, Z.-H. Zheng and X.-H. Lu, *Molecules*, 2014, **19**, 1663–1671.
- 91 C. Takahashi, A. Numata, T. Yamada, K. Minoura, S. Enomoto, K. Konishi, M. Nakai, C. Matsuda and K. Nomoto, *Tetrahedron Lett.*, 1996, **37**, 655–658.
- 92 R. Haritakun, P. Srikitikulchai, P. Khoyaiklang and M. Isaka, *J. Nat. Prod.*, 2007, **70**, 1478–1480.
- 93 T. Bunyapaiboonsri, S. Yoiprommarat, K. Intereya, P. Rachtawee, N. L. Hywel-Jones and M. Isaka, *J. Nat. Prod.*, 2009, **72**, 756–759.
- 94 J. Y. Cha, Y. Huang and T. R. R. Pettus, *Angew. Chem., Int. Ed.*, 2009, **48**, 9519–9521.
- 95 T. Bunyapaiboonsri, S. Yoiprommarat, U. Srisanoh, W. Choowong, K. Tasanathai, N. L. Hywel-Jones, J. J. Luangsa-ard and M. Isaka, *Phytochem. Lett.*, 2011, **4**, 283–286.
- 96 A. Sakakura, K. Shioya, H. Katsuzaki, T. Komiya, T. Imamura, Y. Aizono and K. Imai, *Tetrahedron*, 2009, **65**, 6822–6827.
- 97 A. Sakakura, K. Suzuki, H. Katsuzaki, T. Komiya, T. Imamura, Y. Aizono and K. Imai, *Tetrahedron Lett.*, 2005, **46**, 9057–9059.
- 98 S. P. Putri, K.-i. Ishido, H. Kinoshita, S. Kitani, F. Ihara, Y. Sakihama, Y. Igarashi and T. Nihira, *J. Biosci. Bioeng.*, 2014, **117**, 557–562.
- 99 Y.-L. Chen, C.-S. Chen, S.-H. Yeh, T.-W. Lin, C.-C. Chen and C.-F. Kuo, *Int. J. Med. Mushrooms*, 2015, **17**, 771–781.
- 100 S. J. Kwack and B. M. Lee, *Toxicol. Res.*, 2009, **25**, 101–106.
- 101 J. Song, J. Xin, Y. Zhu and X. Zhang, *Chin JMAP*, 2004, **21**, 12–13.
- 102 M. Y. Ahn, K. S. Ryu, S. D. Jee, I. Kim, J. W. Kim, Y. S. Kim, H. S. Kim, I. S. Kim, S. C. Kang, H. J. Koo, Y. A. Park, S. M. Choi, E. J. Yoo, S. J. Kwack, S. D. Yoo and B. M. Lee, *J. Toxicol. Environ. Health, Part A*, 2004, **67**, 2037–2044.
- 103 P. Song, X. Liu and Q. Zhuo, *World Chin. J. Digestol.*, 1997, **5**, 22.
- 104 J.-H. Che, J.-W. Yun, E.-Y. Cho, S.-H. Kim, Y.-S. Kim, W. H. Kim, J.-H. Park, W.-C. Son, M. K. Kim and B.-C. Kang, *Regul. Toxicol. Pharmacol.*, 2014, **70**, 527–534.
- 105 L. Du, Y. Liu, C. Liu, Q. Meng, J. Song, D. Wang, J. Lu, L. Teng, Y. Zhou and L. Teng, *Comb. Chem. High Throughput Screening*, 2015, **18**, 809–818.
- 106 H. Zhang, X. Shin, T. Liu, W. Ma, Y. Zhang and X. Shi, *Global Tradit. Chin. Med.*, 2017, **10**, 297–301.
- 107 Z. Zhang, T. Chen, B. Yin, Q. Cui, S. Zhang and Y. Wang, *Drug Eval. Res.*, 2016, **39**, 797–805.
- 108 H. Zhang, X. Gao, W. Chen and K. Huang, *Acta Agric. Shanghai*, 2016, **32**, 101–104.
- 109 H. Lu, A. Yang and H. Zhang, *Pract. Pharm. Clin. Rem.*, 2015, **18**, 1466–1469.
- 110 B. C. Moon, W. J. Kim, I. Park, G. H. Sung and P. Noh, *Molecules*, 2018, **23**, 1932.
- 111 W. Zhang, L. Chen, S. Gao, F. Xu, J. Liu, R. Lu and F. Hu, *Junwu Xuebao*, 2015, **34**, 252–268.
- 112 W. Lu, J. Xu, S. Xu and Q. Yuan, *Zhejiang Zhongyiyao Daxue Xuebao*, 2013, **37**, 601–605.

