

Cite this: *RSC Adv.*, 2019, 9, 22080

# New *N*-phenylacetamide-incorporated 1,2,3-triazoles: [Et<sub>3</sub>NH][OAc]-mediated efficient synthesis and biological evaluation†

Satish V. Akolkar,<sup>a</sup> Amol A. Nagargoje,<sup>a</sup> Vagolu S. Krishna,<sup>b</sup> Dharmarajan Sriram,<sup>b</sup> Jaiprakash N. Sangshetti,<sup>c</sup> Manoj Damale<sup>d</sup> and Bapurao B. Shingate<sup>\*a</sup>

A facile, highly efficient, and greener method for the synthesis of new 1,4-disubstituted-1,2,3-triazoles was conducted using [Et<sub>3</sub>NH][OAc] as a medium by the implementation of ultrasound irradiation *via* click chemistry, affording excellent yields. The present synthetic method exhibited numerous advantages such as mild reaction conditions, excellent product yields, minimal chemical waste, operational simplicity, shorter reaction time, and a wide range of substrate scope. The synthesized compounds were further evaluated for *in vitro* antifungal activity against five fungal strains, and some of the compounds displayed equivalent or greater potency than the standard drug. A molecular docking study against the modelled three-dimensional structure of cytochrome P450 lanosterol 14 $\alpha$ -demethylase was also performed to understand the binding affinity and binding interactions of the enzyme. Furthermore, the synthesized compounds were evaluated for DPPH radical scavenging activity and antitubercular activity against *Mycobacterium tuberculosis* H37Rv strain.

Received 7th May 2019  
Accepted 27th June 2019

DOI: 10.1039/c9ra03425k

rsc.li/rsc-advances

## 1. Introduction

1,2,3-Triazole, a five-membered heterocyclic ring system, is a very well-known biologically active pharmacophore constructed by the copper-catalyzed azide-alkyne cycloaddition (CuAAC) reaction, which is popular as a click chemistry reaction.<sup>1</sup> Over the last decade, 1,2,3-triazole has become one of the key structural motifs and is used in numerous fields including polymer chemistry,<sup>2</sup> material science,<sup>3</sup> and drug discovery.<sup>4</sup> 1,2,3-triazole-based molecules display various biological activities such as anti-fungal,<sup>5,6</sup> anti-bacterial,<sup>6</sup> anti-tubercular,<sup>7</sup> anti-inflammatory,<sup>8</sup> anti-allergic,<sup>9</sup> anti-HIV,<sup>10</sup> anti-cancer,<sup>11</sup> and anti-phytopathogenic.<sup>12</sup> Some marketed drugs have the 1,2,3-triazole unit in their structure, and these include Cefatrizine (a broad-spectrum antibiotic), Tazobactam (an antibiotic), and Carboxyamidotriazole (CAI) (a calcium channel blocker) (Fig. 1).

Azole drugs have broad-spectrum activities against most yeasts and filamentous fungi and are mostly used in antifungal

chemotherapy.<sup>13</sup> Some well-known antifungal agents including fluconazole, voriconazole, ravuconazole, and itraconazole contain a 1,2,4-triazole ring in their structure, as shown in Fig. 2. However, their clinical uses have been restricted by their relatively high risk of toxicity, pharmacokinetic deficiencies, emergence of drug resistance, and undesirable side effects. These antifungal drugs inhibit CYP51, a key enzyme in the biosynthesis of ergosterol, through a mechanism in which the antifungal agent having a triazole scaffold is positioned perpendicular to the porphyrin plane with a ring nitrogen atom (N-4 of triazole) coordinated with a heme iron atom.<sup>14</sup> Over a couple of decades, the incidence of systemic fungal infections has increased due to cancer chemotherapy, organ transplantation, tuberculosis, and immunodeficiency virus (HIV) infection.<sup>15</sup> However, the extensive use of antifungal drugs has led to an increase in the resistance to these drugs.<sup>16</sup> Hence, there is an urgent need for developing new antifungal agents with effective activities, low toxicity, and minimum side effects.

Ionic liquids (ILs) are environment-friendly solvents because of their interesting properties and they can be used as alternatives to harmful organic solvents. Furthermore, they are useful in catalytic reactions<sup>17</sup> and organic synthesis<sup>18</sup> because of their




Fig. 1 Structures of drugs containing the 1,2,3-triazole unit.

<sup>a</sup>Department of Chemistry, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad 431 004, India. E-mail: bapushingate@gmail.com; Fax: +91-240-2403113; Tel: +91-240-2403313

<sup>b</sup>Department of Pharmacy, Birla Institute of Technology & Science-Hyderabad Campus, Jawahar Nagar, Hyderabad 500 078, India

<sup>c</sup>Department of Pharmaceutical Chemistry, Y. B. Chavan College of Pharmacy, Dr. Rafiq Zakaria Campus, Aurangabad 431 001, India

<sup>d</sup>Department of Pharmaceutical Chemistry, Srinath College of Pharmacy, Aurangabad 431136, MS, India

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra03425k






Fig. 2 1,2,4-Triazole-based marketed antifungal drugs.


unique properties, making them superior media for increasing selectivity, reactivity, and recyclability. Various ionic liquids have been significantly used in heterocyclic synthesis as solvents or catalysts.<sup>19</sup> The copper-catalyzed 1,3-dipolar cycloaddition of organic azides and terminal alkynes has been reported by using ionic liquids such as [bmim][BF<sub>4</sub>],<sup>20a,b</sup> [Bmim]OH,<sup>20c</sup> (SNIL-Cu(II)),<sup>20d</sup> ([Hmim]TFA),<sup>20e</sup> and [C<sub>8</sub>dabco][N(CN)<sub>2</sub>],<sup>20f</sup> and DBU-based ionic liquids.<sup>20g</sup> The ionic liquid [Et<sub>3</sub>NH][OAc] has been used extensively in various organic transformations.<sup>21</sup>

Considering the importance of ionic liquids in organic synthesis and in continuation of our work<sup>7d,22</sup> as well as for the development of new synthetic methodologies using ionic liquids,<sup>23</sup> herein, we reported the efficient synthesis of new 1,4-disubstituted-1,2,3-triazoles *via* the click chemistry approach using [Et<sub>3</sub>NH][OAc] as the medium under ultrasonic irradiation. Furthermore, the newly synthesized compounds were investigated for their antifungal, antioxidant, and antitubercular activities.

## 2. Results and discussion

### 2.1 Chemistry

There are several reports on the synthesis of 1,4-disubstituted-1,2,3-triazoles bearing amide functionality and displaying various biological activities;<sup>24</sup> recently, Ferroni *et al.* developed triazoles as non-steroidal anti-androgens for prostate cancer treatment.<sup>25</sup> On the basis of these findings, we designed small




Scheme 1 Synthesis of alkynes **2a–c** and 2-azido-*N*-phenylacetamides **5a–g**. Reagents and conditions: (a) propargyl bromide, K<sub>2</sub>CO<sub>3</sub>, DMF, 2 h, 90–95%; (b) chloroacetyl chloride, NEt<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to rt, 3–5 h, 85–95%; (c) NaN<sub>3</sub>, toluene, reflux, 5–7 h, 88–96%.

1,2,3-triazoles with amide linkage in their structures. Initially, the starting materials, *i.e.*, (prop-2-yn-1-yloxy)benzenes **2a–c** were prepared by a previously reported method.<sup>7d</sup> 2-Azido-*N*-phenylacetamides **5a–g** were synthesized from their corresponding anilines *via* chloroacetylation using chloroacetyl chloride, followed by nucleophilic substitution with sodium azide in excellent yields (Scheme 1).

After the synthesis of alkynes **2a–c** and azides **5a–g**, we considered the model substrates (prop-2-yn-1-yloxy)benzenes **2a** (1.0 mmol) and 2-azido-*N*-phenylacetamide **5a** (1.0 mmol) for the click reaction (Scheme 2).

The literature survey reveals the use of *t*-BuOH–H<sub>2</sub>O, THF–H<sub>2</sub>O, and DMF–H<sub>2</sub>O as solvents in combination with copper salts for the click reaction. Accordingly, we performed the model reactions of **2a** and **5a** by using CuSO<sub>4</sub>·5H<sub>2</sub>O, Cu(OAc)<sub>2</sub>·H<sub>2</sub>O, CuI, and CuCl with or without sodium ascorbate as a reducing agent in solvents such as *t*-BuOH and THF (Table 1). In our initial attempt, the model reaction was carried out by using the conventional method, *i.e.*, the reaction mixture was stirred at room temperature for 10 h using various copper salts in *t*-BuOH or THF as a solvent (Table 1, entries 2–7). The product 2-(4-(phenoxy)methyl)-1*H*-1,2,3-triazol-1-yl)-*N*-phenylacetamide **6a** was obtained in 28–72% yield. Furthermore, the model reaction was carried out using copper salts and *t*-BuOH/THF as a solvent under ultrasound irradiation (40 kHz, 30 °C) for 7 h, which resulted in the product **6a** in 32–83% yield (Table 1, entries 2–7). These results indicate a longer reaction time with a lower yield of the product **6a**.


Ionic liquids have unique properties as solvents and as media; herein, we attempted to use various ionic liquids for the synthesis of new amide-linked 1,4-disubstituted-1,2,3-triazole **6a** *via* the click chemistry approach. Freshly synthesized appropriate ionic liquids such as [Et<sub>3</sub>NH][OAc], [Et<sub>3</sub>NH][Cl], [Et<sub>3</sub>NH][HSO<sub>4</sub>], [HDBU][OAc], and [HDBU][HSO<sub>4</sub>] in combination with copper salts such as CuSO<sub>4</sub>·5H<sub>2</sub>O and Cu(OAc)<sub>2</sub>·H<sub>2</sub>O were used along with sodium ascorbate as a reducing agent to carry out the model reaction (Table 1, entries 9–16) at room temperature for 7 h. The maximum yield (90%) of the product **6a** was obtained by using Cu(OAc)<sub>2</sub>·H<sub>2</sub>O as the catalyst and [Et<sub>3</sub>NH][OAc] as the solvent (Table 1, entries 9 and 10). In our further attempt, the model reaction was carried out under ultrasound irradiation for appropriate times (35 and 45 min) (Table 1, entries 9–16). The product **6a** was obtained in 95% yield under ultrasound irradiation using Cu(OAc)<sub>2</sub>·H<sub>2</sub>O as the catalyst and [Et<sub>3</sub>NH][OAc] as the solvent (Table 1, entries 9 and 10). In short, it was observed from the optimization of the best reaction conditions that the Cu(OAc)<sub>2</sub>·H<sub>2</sub>O-catalyzed click reaction in [Et<sub>3</sub>NH][OAc] ionic liquid under ultrasound



Scheme 2 Standard model reaction.



Table 1 Optimization of the reaction conditions for the synthesis of 6a



Entry	Copper salt	Solvent/medium	Conventional method		US method	
			Time <sup>a</sup>	Yield <sup>b</sup> (%)	Time <sup>a</sup>	Yield <sup>b</sup> (%)
1	—	<i>t</i> -BuOH : H <sub>2</sub> O	10 h	0	6 h	0
2	CuSO <sub>4</sub> ·5H <sub>2</sub> O	<i>t</i> -BuOH : H <sub>2</sub> O	10 h	28	6 h	32
3	CuSO <sub>4</sub> ·5H <sub>2</sub> O + Na ascorbate	<i>t</i> -BuOH : H <sub>2</sub> O	10 h	70	6 h	78
4	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O	<i>t</i> -BuOH : H <sub>2</sub> O	10 h	72	6 h	82
5	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O + Na ascorbate	<i>t</i> -BuOH : H <sub>2</sub> O	10 h	72	6 h	83
6	CuI	THF	10 h	70	6 h	80
7	CuCl	THF	10 h	60	6 h	65
8	—	[Et <sub>3</sub> NH][OAc]	10 h	0	6 h	0
9	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O	[Et <sub>3</sub> NH][OAc]	7 h	90	35 min	95
10	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O + Na ascorbate	[Et <sub>3</sub> NH][OAc]	7 h	90	35 min	95
11	CuSO <sub>4</sub> ·5H <sub>2</sub> O + Na ascorbate	[Et <sub>3</sub> NH][OAc]	7 h	81	35 min	84
12	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O	[Et <sub>3</sub> NH][Cl]	7 h	Trace	35 min	10
13	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O	[Et <sub>3</sub> NH][HSO <sub>4</sub> ]	7 h	Trace	35 min	12
14	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O	[HDBU][OAc]	7 h	84	45 min	88
15	CuSO <sub>4</sub> ·5H <sub>2</sub> O + Na ascorbate	[HDBU][OAc]	7 h	81	45 min	84
16	Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O	[HDBU][HSO <sub>4</sub> ]	7 h	10	45 min	14

<sup>a</sup> Reaction conditions: 2a (1 mmol), 5a (1 mmol), and 10 mol% [Cu], 1 mL solvent, 30 °C; reaction progress monitored by TLC. <sup>b</sup> Isolated yield, US: ultrasonication.

irradiation at 30 °C efficiently gives the desired 1,4-disubstituted-1,2,3-triazole derivative 6a as compared to the conventional method with respect to the reaction time and yield (Table 1, entry 9).

In order to evaluate the amount of the Cu(OAc)<sub>2</sub>·H<sub>2</sub>O catalyst for the model reaction for different concentrations such as 5, 10, 15, and 20 mol% (Table 2, entries 1–4), the reaction was carried out in the presence of 1 mL of [Et<sub>3</sub>NH][OAc] as a solvent at 30 °C under ultrasound irradiation. It was observed that 10 mol% of Cu(OAc)<sub>2</sub>·H<sub>2</sub>O was sufficient to achieve an excellent yield of the product 6a in the shortest reaction time (Table 2, entry 2), while an excess amount of the catalyst (up to 20 mol%) did not increase the yield of the product. The above results reveal that 10 mol% of Cu(OAc)<sub>2</sub>·H<sub>2</sub>O in [Et<sub>3</sub>NH][OAc] is the sufficient catalyst amount in terms of an excellent yield and short reaction time (Table 2, entry 2).


The recyclability of the [Et<sub>3</sub>NH][OAc] solvent was also explored for the model reaction. The reaction was conducted for four successive cycles on the same recycled solvent. After completion of the first reaction with 95% yield, the reaction mass was poured into ice cold water to get the solid product, which was isolated by filtration. The filtrate was subjected to evaporation of water to get a viscous liquid, which on cooling gave the ionic liquid. Then, the recovered [Et<sub>3</sub>NH][OAc] was reused for four more successive cycles without significant loss in the yield of the product 6a. The yields achieved were 94%,

92%, 91%, and 88% after the successful runs (Fig. 3). The result demonstrates the high stability of the ionic liquid [Et<sub>3</sub>NH][OAc] under the reaction conditions, which could be recycled and reused without a significant loss in the product yield.

In a further attempt, we also examined a one-pot reaction for the selected model reaction with 2a (1.0 mmol), 4a (1.0 mmol), and sodium azide (1.5 mmol) for the synthesis of 6a, which was optimized for different copper sources (Table 3) for the one-pot, three-component click reaction in [Et<sub>3</sub>NH][OAc] as the solvent, thus producing 2-(4-(phenoxy)methyl)-1H-1,2,3-triazol-1-yl)-N-phenylacetamide 6a (Scheme 3). The optimization result of this one-pot model reaction revealed that 15 mol% of Cu(OAc)<sub>2</sub>·H<sub>2</sub>O in [Et<sub>3</sub>NH][OAc] as the solvent was a sufficient catalyst loading amount to obtain an excellent yield. However, it required a longer reaction time and higher concentration of the Cu catalyst (Table 3, entry 2) than the two-component reaction, as discussed earlier.

After optimization of the reaction parameters and to define the present methodology, as discussed earlier, the reaction was assessed with a variety of substituted alkynes 2a–c and azides 5a–g in the presence of Cu(OAc)<sub>2</sub>·H<sub>2</sub>O in [Et<sub>3</sub>NH][OAc] under ultrasound irradiation. The reaction conditions were favorable for the electron-donating and electron-withdrawing groups on alkynes 2a–c and azides 5a–g, resulting in the corresponding 1,4-disubstituted-1,2,3-triazole derivatives 6a–g, 7a–g, and 8a–g in excellent yields (Scheme 4).




**Table 2** The influence of the amount of  $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$  on the synthesis of **6a**


Entry	mol%	Time (min)	Isolated yield (%)
1	5	55	70
2	10	35	95
3	15	35	95
4	20	35	95

All the synthesized compounds, namely, **6a-g**, **7a-g**, and **8a-g** were confirmed by physical data and spectral analysis.

## 2.2 Biological activity

**2.2.1. Antifungal activity.** All the newly synthesized compounds were screened for their *in vitro* antifungal activities against five different fungal strains, namely, *Candida albicans*, *Fusarium oxysporum*, *Aspergillus flavus*, *Aspergillus niger*, and *Cryptococcus neoformans*. The minimum inhibitory concentration (MIC,  $\mu\text{g mL}^{-1}$ ) values of all the newly synthesized compounds were determined by the standard agar dilution method as per the CLSI (formerly, NCCLS) guidelines<sup>26</sup> (Table 4). DMSO was used as the negative control and Miconazole was used as the standard antifungal drug for the comparison of antifungal activities. Among the series, the triazole derivatives **6** and **7** derived from alkynes **2a** and **2b** (particularly from phenol and 4-methoxyphenol) did not show any activity against all the fungal strains. However, the triazoles derived from alkyne **2c** (from 4-nitrophenol) displayed excellent antifungal activity against all the strains. The compounds **8b**, **8c**, and **8f** having MIC of  $12.5 \mu\text{g mL}^{-1}$  were found to be more potent; compounds **8a**, **8d**, and **8g** having MIC of  $25 \mu\text{g mL}^{-1}$  were found to be equipotent as compared to the standard drug Miconazole against the fungal strain *Candida albicans*. The compounds **8b**, **8c**, **8d**, **8e**, and **8g** with MIC of  $25 \mu\text{g mL}^{-1}$  were found to be equivalent to the standard drug Miconazole and **8f** with MIC of


**Fig. 3** Reusability of  $[\text{Et}_3\text{NH}][\text{OAc}]$ .**Table 3** The effect of the amount of copper salts for the one-pot synthesis of **6a**

Entry	Catalyst	mol%	Time (h)	Isolated yield (%)
1	$\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$	10	17	90
2	$\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$	15	16	95
3	$\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$	20	16	95
4	$\text{CuSO}_4 \cdot 5\text{H}_2\text{O} + \text{Na ascorbate}$	10	18	81
5	$\text{CuSO}_4 \cdot 5\text{H}_2\text{O} + \text{Na ascorbate}$	15	17	87
6	$\text{CuI}$	10	18	50
7	$\text{CuI}$	15	17	61
8	$\text{CuCl}$	10	17	40
9	$\text{CuCl}$	15	17	45

$12.5 \mu\text{g mL}^{-1}$  was more potent than Miconazole against the fungal strain *F. oxysporum*. Compounds **8e** and **8g** with MIC of  $12.5 \mu\text{g mL}^{-1}$  exhibited equivalent activities as compared to Miconazole against *A. flavus*. For the fungal strain *A. niger*, the compounds **8b**, **8d**, **8e**, and **8g** with MIC of  $25 \mu\text{g mL}^{-1}$  displayed equivalent antifungal activities as compared to Miconazole. However, **8f** having MIC of  $12.5 \mu\text{g mL}^{-1}$  was more active than Miconazole. For the fungal strain *C. neoformans*, the compounds **8e**, **8f**, and **8g** with MIC of  $12.5 \mu\text{g mL}^{-1}$  were more potent than the standard drug and **8c** with MIC of  $25 \mu\text{g mL}^{-1}$  was equivalent to Miconazole.

The remaining compounds displayed moderate to low antifungal activities against all the tested fungal strains. The antifungal activity depends on the various substituents present on phenyl rings. Among the series, the compounds with nitro groups (**8a-g**) were the most active against all the tested fungal strains. Compounds **8f** and **8g** were the most active in the series as they possessed a nitro group at R and chloro groups at R<sup>2</sup> and R<sup>3</sup>; the chloro group at the R<sup>1</sup> position for **8e** resulted in less activity against one fungal strain. The methyl group at the R<sup>1</sup>/R<sup>2</sup>/R<sup>3</sup> position and the nitro group at the R position in **8b**, **8c**, and **8d** resulted in good activity than that of the others in the series.

**2.2.2. Antitubercular activity.** All the newly synthesized 1,4-disubstituted-1,2,3-triazoles (**6a-g**, **7a-g**, and **8a-g**) were screened for their *in vitro* antitubercular activities against the MTB H37Rv (ATCC 27294) strain using the microplate Alamar Blue assay method with the determination of MIC in triplicate; the results are mentioned in Table 4. The newly synthesized 1,2,3-triazoles exhibited MICs ranging from  $25$  to  $\geq 25 \mu\text{g mL}^{-1}$ . Among the series, compounds **6a**, **6c**, and **6f** exhibited good antitubercular activities with MIC of  $25 \mu\text{g mL}^{-1}$  as compared to other compounds.

**Scheme 3** Synthesis of 1,4-disubstituted-1,2,3-triazole **6a** via the one-pot three-component click reaction.


Scheme 4 Synthesis of 1,4-disubstituted-1,2,3-triazole derivatives **6a-g**, **7a-g**, and **8a-g**.

Table 4 *In vitro* biological evaluation of the synthesized compounds 6a–g, 7a–g, and 8a–g MIC ( $\mu\text{g mL}^{-1}$ )<sup>a</sup>

Entry	Antifungal activity					Anti TB MTB H37Rv	DPPH IC <sub>50</sub> ( $\mu\text{g mL}^{-1}$ )
	CA	FO	AF	AN	CN		
6a	87.5	100	175	125	187.5	25	55.37
6b	150	125	150	150	*	>25	26.64
6c	175	200	150	175	200	25	20.54
6d	187.5	162.5	187.5	200	*	>25	*
6e	200	*	*	187.5	175	>25	29.64
6f	150	175	162.5	200	*	25	74.28
6g	162.5	175	175	187.5	150	>25	74.41
7a	137.5	125	100	137.5	125	>25	11.44
7b	175	200	100	175	200	>25	16.79
7c	*	*	200	*	175	>25	12.30
7d	200	175	150	150	200	>25	19.46
7e	175	175	200	*	*	>25	70.33
7f	100	150	100	100	75	>25	*
7g	75	100	87.5	100	200	>25	49.60
8a	25	37.5	25	75	75	>25	97.29
8b	12.5	25	25	25	37.5	>25	*
8c	12.5	25	25	50	25	>25	48.02
8d	25	25	37.5	25	50	>25	22.17
8e	50	25	12.5	25	12.5	>25	62.01
8f	12.5	12.5	25	12.5	12.5	>25	*
8g	25	25	12.5	25	12.5	>25	16.99
MA	25	25	12.5	25	25	NA	NA
INH	NA	NA	NA	NA	NA	0.1	NA
RIF	NA	NA	NA	NA	NA	0.2	NA
EMB	NA	NA	NA	NA	NA	1.56	NA
CIP	NA	NA	NA	NA	NA	1.56	NA
BHT	NA	NA	NA	NA	NA	NA	16.47

<sup>a</sup> CA: *Candida albicans*; FO: *Fusarium oxysporum*; AF: *Aspergillus flavus*; AN: *Aspergillus niger* and CN: *Cryptococcus neoformans*. MA: Miconazole, DPPH: 2,2-diphenyl-1-picrylhydrazyl; INH: Isoniazid; RIF: Rifampicin; EMB: Ethambutol; CIP: Ciprofloxacin; BHT: butylated hydroxytoluene; \*no activity was observed, NA: not applicable.

**2.2.3. Antioxidant activity.** The antioxidant activities of the synthesized compounds 6a–g, 7a–g, and 8a–g were screened using a 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay.<sup>27</sup> The DPPH radical scavenging assay is the most commonly used procedure for screening the antioxidant activities of different natural and synthetic compounds. A lower IC<sub>50</sub> value demonstrates more antioxidant activity. The IC<sub>50</sub> (concentration required to scavenge 50% of the radicals) values were calculated to assess the potential antioxidant activities.

Butylated hydroxytoluene (BHT) was used as the standard drug for the comparison of antioxidant activities; the observed results are summarized in Table 4. According to the DPPH assay, compounds 7a and 7c exhibited excellent radical scavenging activities with IC<sub>50</sub> values of 11.44 and 12.30  $\mu\text{g mL}^{-1}$ , respectively, compared to the synthetic antioxidant BHT. Similarly, compounds 7b, 7d, and 8g having IC<sub>50</sub> values of 16.79, 19.46, and 16.99  $\mu\text{g mL}^{-1}$ , respectively, showed comparable antioxidant activities with that of BHT; in contrast, the

Table 5 Molecular docking score for the compounds 6a–g, 7a–g, and 8a–g

Molecular docking score							
Entry	Total score ( $-\log K_i$ )	Crash score	Polar score	Entry	Total score ( $-\log K_i$ )	Crash score	Polar score
6a	7.0377	-0.6933	1.9115	7e	6.4127	-1.2494	1.4225
6b	6.4564	-2.9621	0.0004	7f	7.2492	-1.4724	2.5093
6c	6.8229	-0.7726	2.0682	7g	7.2394	-1.4033	3.0192
6d	6.0468	-1.4517	1.3586	8a	7.0868	-1.1801	3.6314
6e	6.5873	-0.9241	2.8329	8b	7.5441	-2.8665	0.9019
6f	6.7343	-0.8582	3.0543	8c	7.4658	-1.0993	0.7391
6g	7.0332	-1.0463	1.6968	8d	7.3945	-1.5222	1.9024
7a	6.0045	-0.7865	2.8105	8e	7.3314	-0.9292	0.7808
7b	6.6907	-0.9183	4.5244	8f	7.9971	-1.9946	0.0031
7c	6.5748	-2.7834	0.0090	8g	7.5972	-1.3842	3.0509
7d	7.0239	-1.2982	3.2802	MA	6.4895	-1.4009	1.0311





and Bruker Advance III 125 MHz NMR in DMSO-*d*<sub>6</sub>, CDCl<sub>3</sub> + DMSO-*d*<sub>6</sub>, and CDCl<sub>3</sub> + CD<sub>3</sub>OD using tetramethylsilane (TMS) as the internal standard; the chemical shifts ( $\delta$ ) were recorded in parts per million (ppm). The splitting pattern abbreviations are designed as singlet (s), doublet (d), double doublet (dd), triplet (t), quartet (q), and multiplet (m). High-resolution mass spectra (HRMS) were recorded on an Agilent 6520 (QTOF) ESI-HRMS instrument and Agilent Technologies 1200 series electron spray ionization (ESI) was used for the LC-MS data.

## 4.2 General experimental procedure for the synthesis of *N*-phenylacetamide-incorporated 1,2,3-triazole derivatives (6a-g, 7a-g, and 8a-g)

The mixture of appropriate (prop-2-yn-1-yloxy) benzenes **2a-c** (1 mmol), substituted 2-azido-*N*-phenylacetamide **5a-g** (1 mmol) and Cu(OAc)<sub>2</sub>·H<sub>2</sub>O (10 mol%) in [NEt<sub>3</sub>H][OAc] (1 mL) was placed in a round bottom flask. The mixture was sonicated (40 kHz) at 30 °C for an appropriate time until the completion of the reaction. The progress of the reaction was monitored by TLC. After the completion of the reaction, the mixture was quenched with crushed ice. The obtained solid was filtered and washed with water. The crude solid was crystallized in ethanol to afford the corresponding pure product. The synthesized compounds **6a-g**, **7a-g**, and **8a-g** were characterized by IR, <sup>1</sup>H NMR, <sup>13</sup>C NMR, and mass spectroscopy.

**4.2.1. 2-(4-(Phenoxymethyl)-1H-1,2,3-triazol-1-yl)-*N*-phenylacetamide (6a).** The compound **6a** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5a** and alkyne **2a** in 35 min as a white solid; mp: 166–168 °C; FT-IR (cm<sup>-1</sup>): 3263 (N–H stretching), 1667 (C=O stretching); <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 5.17 (s, 2H, –NCH<sub>2</sub>CO–), 5.36 (s, 2H, –OCH<sub>2</sub>), 6.96 (t, *J* = 8.0 Hz, 1H, Ar–H), 7.08 (t, *J* = 8.0 Hz, 3H, Ar), 7.28–7.38 (m, 4H, Ar–H), 7.59 (d, *J* = 8.0 Hz, 2H, Ar–H), 8.26 (s, 1H, triazole), and 10.49 (s, 1H, NH); <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 51.8, 60.5, 114.3, 118.9, 120.5, 123.5, 125.9, 128.6, 129.2, 138.0, 142.2, 157.7, and 163.8; HRMS calcd for C<sub>17</sub>H<sub>17</sub>N<sub>4</sub>O<sub>2</sub> [M + H]<sup>+</sup>: 309.1346 and found 309.1340.

**4.2.2. 2-(4-(Phenoxymethyl)-1H-1,2,3-triazol-1-yl)-*N*-(*o*-tolyl)acetamide (6b).** The compound **6b** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5b** and alkyne **2a** in 37 min as a white solid; mp: 184–186 °C; FT-IR (cm<sup>-1</sup>): 3271 (N–H stretching), 1665 (C=O stretching); <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 2.24 (s, 3H, –CH<sub>3</sub>), 5.17 (s, 2H, –NCH<sub>2</sub>CO–), 5.41 (s, 2H, –OCH<sub>2</sub>), 6.95 (t, *J* = 8.0 Hz, 1H, Ar–H), 7.05 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.10–7.35 (m, 5H, Ar–H), 7.43 (d, *J* = 8.0 Hz, 1H, Ar–H), 8.26 (s, 1H, triazole), and 9.84 (s, 1H, NH); <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 17.5, 51.3, 60.5, 114.3, 120.4, 124.8, 125.8, 129.1, 130.1, 131.2, 135.1, 142.2, 157.7, and 164.0; HRMS calcd for C<sub>18</sub>H<sub>19</sub>N<sub>4</sub>O<sub>2</sub> [M + H]<sup>+</sup>: 323.1508 and found 323.1510.

**4.2.3. 2-(4-(Phenoxymethyl)-1H-1,2,3-triazol-1-yl)-*N*-(*m*-tolyl)acetamide (6c).** The compound **6c** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5c** and alkyne **2a** in 38 min as a yellow solid; mp: 130–132 °C; FT-IR (cm<sup>-1</sup>): 3302 (N–H stretching), 1676 (C=O stretching); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub> + DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 2.21 (s, 3H, –CH<sub>3</sub>), 5.10 (s, 2H, –NCH<sub>2</sub>CO–), 5.13 (s, 2H, –OCH<sub>2</sub>), 6.79–6.93 (m,

4H, Ar–H), 7.08 (d, *J* = 8.0 Hz 1H), 7.13–7.30 (m, 3H, Ar–H), 7.34 (s, 1H, Ar–H), 7.85 (s, 1H, triazole), and 9.70 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub> + DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 21.1, 52.7, 61.4, 114.3, 116.7, 120.2, 124.8, 124.9, 128.3, 129.1, 137.3, 138.4, 143.7, 157.8, and 163.0; LC-MS for C<sub>18</sub>H<sub>18</sub>N<sub>4</sub>O<sub>2</sub>: [M]<sup>+</sup> 323.1.

**4.2.4. 2-(4-(Phenoxymethyl)-1H-1,2,3-triazol-1-yl)-*N*-(*p*-tolyl)acetamide (6d).** The compound **6d** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5d** and alkyne **2a** in 36 min as a yellow solid; mp: 202–204 °C; FT-IR (cm<sup>-1</sup>): 3271 (N–H stretching), 1665 (C=O stretching).

**4.2.5. 2-(4-(Phenoxymethyl)-1H-1,2,3-triazol-1-yl)-*N*-(2-chlorophenyl)-acetamide (6e).** The compound **6e** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5e** and alkyne **2a** in 36 min as a white solid; mp: 178–180 °C; FT-IR (cm<sup>-1</sup>): 3267 (N–H stretching), 1671 (C=O stretching); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub> + DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 5.19 (s, 2H, –NCH<sub>2</sub>CO–), 5.31 (s, 2H, –OCH<sub>2</sub>), 6.94 (d, *J* = 8.0 Hz, 3H, Ar–H), 7.05 (d, *J* = 8.0 Hz, 1H, Ar–H), 7.16–7.34 (m, 4H, Ar–H), 7.89 (s, 1H, triazole), 8.02 (d, *J* = 8.0 Hz, 1H, Ar–H), and 9.01 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub> + DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 52.8, 61.3, 114.3, 120.8, 123.1, 124.3, 125.5, 127.1, 129.0, 133.7, 141.9, 157.7, and 162.6. LC-MS for C<sub>17</sub>H<sub>15</sub>ClN<sub>4</sub>O<sub>2</sub>: [M]<sup>+</sup> 343.

**4.2.6. 2-(4-(Phenoxymethyl)-1H-1,2,3-triazol-1-yl)-*N*-(3-chlorophenyl)-acetamide (6f).** The compound **6f** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5f** and alkyne **2a** in 37 min as a white solid; mp: 140–142 °C; FT-IR (cm<sup>-1</sup>): 3257 (N–H stretching), 1672 (C=O stretching); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub> + DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 5.45 (s, 2H, –NCH<sub>2</sub>CO–), 5.46 (s, 2H, –OCH<sub>2</sub>), 7.15–7.57 (m, 7H Ar–H), 7.67 (s, 1H, Ar–H), 7.92 (s, 1H, triazole), 8.15 (s, 1H, Ar–H), and 10.16 (s, 1H, NH). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub> + DMSO-*d*<sub>6</sub>,  $\delta$  ppm): 53.1, 61.7, 114.7, 117.9, 120.0, 121.2, 124.5, 129.5, 129.9, 134.4, 138.9, 158.1, and 163.5; LC-MS for C<sub>17</sub>H<sub>15</sub>ClN<sub>4</sub>O<sub>2</sub>: [M]<sup>+</sup> 343.1.

**4.2.7. 2-(4-((4-Methoxyphenoxy)methyl)-1H-1,2,3-triazol-1-yl)-*N*-phenylacetamide (7a).** The compound **7a** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5a** and alkyne **2b** in 37 min as a white solid; mp: 180–182 °C; FT-IR (cm<sup>-1</sup>): 3271 (N–H stretching), 1674 (C=O stretching); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>,  $\delta$  ppm): 3.76 (s, 3H, OCH<sub>3</sub>), 5.19 (s, 2H, –NCH<sub>2</sub>CO–), 5.20 (s, 2H, –OCH<sub>2</sub>), 6.83 (d, *J* = 8.0 Hz, 2H, Ar–H), 6.92 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.14 (t, *J* = 8.0 Hz, 1H, Ar–H), 7.31 (t, *J* = 8.0 Hz, 2H, Ar–H), 7.43 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.81 (s, 1H, triazole), and 7.87 (s, 1H, NH); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>,  $\delta$  ppm): 53.9, 55.7, 62.5, 114.8, 115.9, 120.3, 125.4, 129.1, 137.0, 139.3, 152.1, and 162.3; LC-MS for C<sub>18</sub>H<sub>18</sub>N<sub>4</sub>O<sub>3</sub>: [M]<sup>+</sup> 339.1.

**4.2.8. 2-(4-((4-Methoxyphenoxy)methyl)-1H-1,2,3-triazol-1-yl)-*N*-(*o*-tolyl)acetamide (7b).** The compound **7b** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5b** and alkyne **2b** in 35 min as a white solid; mp: 176–178 °C; FT-IR (cm<sup>-1</sup>): 3266 (N–H stretching), 1674 (C=O stretching); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>,  $\delta$  ppm): 2.10 (s, 3H, CH<sub>3</sub>), 3.75 (s, 3H, –OCH<sub>3</sub>), 5.21 (s, 2H, –NCH<sub>2</sub>CO–), 5.22 (s, 2H, –OCH<sub>2</sub>), 6.82 (d, *J* = 8.0 Hz, 2H, Ar–H), 6.91 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.08 (t, *J* = 8.0 Hz, 1H, Ar–H), 7.15 (d, *J* = 8.0 Hz, 1H, Ar–H), 7.20 (t, *J* =





8.0 Hz, 1H, Ar-H), 7.80 (s, 1H, triazole), and 7.82 (s, 1H, NH);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ ,  $\delta$  ppm): 17.4, 53.9, 55.7, 62.4, 114.8, 115.9, 122.3, 124.5, 125.9, 126.9, 127.9, 137.6, and 162.8; LC-MS for  $\text{C}_{19}\text{H}_{20}\text{N}_4\text{O}_3$ :  $[\text{M}]^+$  353.1.

**4.2.9.** **2-(4-((4-Methoxyphenoxy)methyl)-1H-1,2,3-triazol-1-yl)-N-(m-tolyl)acetamide (7c).** The compound **7c** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5c** and alkyne **2b** in 35 min as a yellow solid; mp: 140–142 °C; FT-IR ( $\text{cm}^{-1}$ ): 3271 (N–H stretching), 1665 (C=O stretching);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 2.18 (s, 3H,  $\text{CH}_3$ ), 3.62 (s, 3H,  $\text{OCH}_3$ ), 5.01 (s, 2H,  $-\text{NCH}_2\text{CO}-$ ), 5.10 (s, 2H,  $-\text{OCH}_2$ ), 6.68 (d,  $J = 6$  Hz, 2H, Ar-H), 6.79 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.04 (t,  $J = 8.0$  Hz, 1H, Ar-H), 7.18 (d,  $J = 8.0$  Hz, 1H, Ar-H), 7.26 (d,  $J = 8.0$  Hz, 1H, Ar-H), 7.31 (s, 1H, Ar-H), 7.79 (s, 1H, triazole), and 9.73 (s, 1H, NH);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 21.3, 52.9, 55.5, 62.4, 114.5, 115.7, 116.8, 120.4, 124.4, 125.1, 128.5, 137.6, 138.5, 144.1, 152.2, 153.9, and 163.3. LC-MS for  $\text{C}_{19}\text{H}_{20}\text{N}_4\text{O}_3$ :  $[\text{M}]^+$  353.1.

**4.2.10.** **2-(4-((4-Methoxyphenoxy)methyl)-1H-1,2,3-triazol-1-yl)-N-(p-tolyl)acetamide (7d).** The compound **7d** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5d** and alkyne **2b** in 36 min as a white solid; mp: 196–198 °C; FT-IR ( $\text{cm}^{-1}$ ): 3273 (N–H stretching), 1675 (C=O stretching);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 2.10 (s, 3H,  $\text{CH}_3$ ), 3.55 (s, 3H,  $\text{OCH}_3$ ), 4.94 (s, 2H,  $-\text{NCH}_2\text{CO}-$ ), 5.02 (s, 2H,  $-\text{OCH}_2$ ), 6.62 (d,  $J = 8.0$  Hz, 2H, Ar-H), 6.72 (d,  $J = 8.0$  Hz, 2H, Ar-H), 6.89 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.24 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.73 (s, 1H, triazole), and 9.68 (s, 1H, NH);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 20.4, 52.6, 55.2, 62.1, 114.2, 115.4, 119.5, 124.2, 128.9, 133.5, 134.9, 151.9, 153.6, and 162.8; LC-MS for  $\text{C}_{19}\text{H}_{20}\text{N}_4\text{O}_3$ :  $[\text{M}]^+$  353.1.

**4.2.11.** **N-(2-Chlorophenyl)-2-(4-((4-methoxyphenoxy)methyl)-1H-1,2,3-triazol-1-yl)acetamide (7e).** The compound **7e** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5e** and alkyne **2b** in 35 min as a white solid; mp: 168–170 °C; FT-IR ( $\text{cm}^{-1}$ ): 3260 (N–H stretching), 1683 (C=O stretching);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 3.89 (s, 3H,  $\text{OCH}_3$ ), 5.32 (s, 2H,  $-\text{NCH}_2\text{CO}-$ ), 5.47 (s,  $-\text{OCH}_2$ ), 6.96 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.06 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.23 (t,  $J = 8.0$  Hz, 1H, Ar-H), 7.39 (t,  $J = 8.0$  Hz, 1H, Ar-H), 7.50 (d,  $J = 8.0$  Hz, 1H, Ar-H), 8.03 (s, 1H, triazole), 8.26 (d,  $J = 8.0$  Hz, 1H, Ar-H), and 8.98 (s, 1H, NH);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 53.0, 55.4, 62.3, 114.4, 115.6, 122.8, 124.3, 125.6, 127.3, 129.1, 133.4, 144.7, 152.0, 153.9, and 163.3; LC-MS for  $\text{C}_{18}\text{H}_{17}\text{ClN}_4\text{O}_3$ :  $[\text{M}]^+$  373.

**4.2.12.** **N-(2-Chlorophenyl)-2-(4-((4-methoxyphenoxy)methyl)-1H-1,2,3-triazol-1-yl)acetamide (7f).** The compound **7f** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5f** and alkyne **2b** in 35 min as a green solid; mp: 160–162 °C; FT-IR ( $\text{cm}^{-1}$ ): 3266 (N–H stretching), 1675 (C=O stretching);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 3.57 (s, 3H,  $\text{OCH}_3$ ), 4.96 (s, 2H,  $-\text{NCH}_2\text{CO}-$ ), 5.05 (s, 2H,  $-\text{OCH}_2$ ), 6.63 (d,  $J = 8.0$  Hz, 2H, Ar-H), 6.73 (d,  $J = 8.0$  Hz, 2H, Ar-H), 6.87 (d,  $J = 8.0$  Hz, 1H, Ar-H), 7.03 (t,  $J = 8.0$  Hz, 1H, Ar-H), 7.24 (d,  $J = 8.0$  Hz, 1H, Ar-H), 7.53 (s, 1H, Ar-H), 7.73 (s, 1H, triazole), and 10.04 (s, 1H, NH);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$  +  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 52.4, 55.1, 62.0, 114.1, 115.27, 117.3, 119.3, 123.7, 124.1, 129.4, 133.8, 138.7, 151.8, 153.5, and 163.2; LC-MS for  $\text{C}_{18}\text{H}_{17}\text{ClN}_4\text{O}_3$   $[\text{M} + \text{H}]^+$  373.

**4.2.13.** **N-(4-Chlorophenyl)-2-(4-((4-methoxyphenoxy)methyl)-1H-1,2,3-triazol-1-yl)acetamide (7g).** The compound **7g** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5g** and alkyne **2b** in 36 min as a white solid; mp: 192–194 °C; FT-IR ( $\text{cm}^{-1}$ ): 3266 (N–H stretching), 1672 (C=O stretching).

**4.2.14.** **2-(4-((4-Nitrophenoxy)methyl)-1H-1,2,3-triazol-1-yl)-N-phenylacetamide (8a).** The compound **8a** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5a** and alkyne **2c** in 40 min as a yellow solid; mp: 180–182 °C; FT-IR ( $\text{cm}^{-1}$ ): 3250 (N–H stretching), 1682 (C=O stretching);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$  +  $\text{CD}_3\text{OD}$ ,  $\delta$  ppm): 5.18 (s, 2H,  $-\text{NCH}_2\text{CO}-$ ), 5.27 (s, 2H,  $-\text{OCH}_2$ ), 7.05 (d,  $J = 8.0$  Hz, 3H, Ar-H), 7.27 (t,  $J = 8.0$  Hz, 3H, Ar-H), 7.49 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.98 (s, 1H, triazole), and 8.17 (d,  $J = 8$  Hz, 2H, Ar-H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$  +  $\text{CD}_3\text{OD}$ ,  $\delta$  ppm): 52.7, 61.9, 114.5, 125.1, 125.7, 128.8, 137.5, 141.7, and 162.8; LC-MS for  $\text{C}_{17}\text{H}_{15}\text{N}_5\text{O}_4$ :  $[\text{M}]^+$  354.

**4.2.15.** **2-(4-((4-Nitrophenoxy)methyl)-1H-1,2,3-triazol-1-yl)-N-(o-tolyl)acetamide (8b).** The compound **8b** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5b** and alkyne **2c** in 39 min as a white solid; mp: 178–180 °C; FT-IR ( $\text{cm}^{-1}$ ): 3257 (N–H stretching), 1668 (C=O stretching);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$  +  $\text{CD}_3\text{OD}$ ,  $\delta$  ppm): 2.15 (s, 3H,  $\text{CH}_3$ ), 5.23 (s, 2H,  $-\text{NCH}_2\text{CO}-$ ), 5.25 (s, 2H,  $-\text{OCH}_2$ ), 7.03–7.13 (m, 6H, Ar-H), 7.47 (s, 1H, NH), 7.98 (s, 1H, triazole), and 8.14 (s, 2H, Ar-H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$  +  $\text{CD}_3\text{OD}$ ,  $\delta$  ppm): 17.2, 52.6, 62.1, 114.9, 124.9, 125.4, 125.9, 126.4, 126.5, 127.4, 130.7, 131.2, and 163.7; LC-MS for  $\text{C}_{18}\text{H}_{17}\text{N}_5\text{O}_4$   $[\text{M}]^+$  368.1.

**4.2.16.** **2-(4-((4-Nitrophenoxy)methyl)-1H-1,2,3-triazol-1-yl)-N-(m-tolyl)acetamide (8c).** The compound **8c** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5c** and alkyne **2c** in 38 min as a white solid; mp: 168–170 °C; FT-IR ( $\text{cm}^{-1}$ ): 3280 (N–H stretching), 1662 (C=O stretching);  $^1\text{H}$  NMR (200 MHz,  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 2.28 (s, 3H,  $\text{CH}_3$ ), 5.36 (s, 4H,  $-\text{NCH}_2\text{CO}$ ,  $-\text{OCH}_2$ ), 6.91 (d,  $J = 8.0$  Hz, 1H, Ar-H), 7.19 (d,  $J = 8.0$  Hz, 1H, Ar-H), 7.27 (t,  $J = 8$  Hz, 1H, Ar-H), 7.33 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.42 (s, 1H, Ar-H), 8.24 (d,  $J = 8.0$  Hz, 2H, Ar-H), 8.32 (s, 1H, triazole), and 10.42 (s, 1H, NH);  $^{13}\text{C}$  NMR (50 MHz,  $\text{DMSO}-d_6$ ,  $\delta$  ppm): 21.1, 52.2, 61.9, 115.3, 116.4, 119.8, 124.5, 125.9, 126.7, 128.7, 138.1, 138.3, 141.0, 141.5, 163.3, and 164.0; HRMS calcd for  $\text{C}_{18}\text{H}_{18}\text{N}_5\text{O}_4$   $[\text{M} + \text{H}]^+$ : 368.1359 and found 368.1364.

**4.2.17.** **2-(4-((4-Nitrophenoxy)methyl)-1H-1,2,3-triazol-1-yl)-N-(p-tolyl)acetamide (8d).** The compound **8d** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5d** and alkyne **2c** in 40 min as a white solid; mp: 210–212 °C; FT-IR ( $\text{cm}^{-1}$ ): 3260 (N–H stretching), 1696 (C=O stretching);  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$  +  $\text{CD}_3\text{OD}$ ,  $\delta$  ppm): 2.26 (s, 3H,  $\text{CH}_3$ ), 5.14 (s, 2H,  $-\text{NCH}_2\text{CO}-$ ), 5.26 (s, 2H,  $-\text{OCH}_2$ ), 7.0 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.07 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.35 (d,  $J = 8.0$  Hz, 2H, Ar-H), 7.96 (s, 1H, triazole), and 8.16 (d,  $J = 8.0$  Hz, 2H, Ar-H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$  +  $\text{CD}_3\text{OD}$ ,  $\delta$  ppm): 17.7, 53.1, 62.6, 115.4, 124.8, 125.9, 126.5, 126.9, 131.2, 143.5, 152.0, and 164.2; LC-MS for  $\text{C}_{18}\text{H}_{17}\text{N}_5\text{O}_4$ :  $[\text{M} + \text{H}]^+$  368.1.

**4.2.18.** **N-(2-Chlorophenyl)-2-(4-((4-nitrophenoxy)methyl)-1H-1,2,3-triazol-1-yl)acetamide (8e).** The compound **8e** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5e**



and alkyne **2c** in 39 min as a red solid; mp: 169–171 °C; FT-IR (cm<sup>-1</sup>): 3243 (N–H stretching), 1672 (C=O stretching); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub> + CD<sub>3</sub>OD, δ ppm): 5.31 (s, 2H, –NCH<sub>2</sub>CO–), 5.33 (s, 2H, –OCH<sub>2</sub>), 7.08 (d, *J* = 8.0 Hz, 3H, Ar–H), 7.29 (d, *J* = 8.0 Hz, 2H), 7.37 (d, *J* = 8.0 Hz, 1H, Ar–H), 7.96 (s, 1H, triazole), 8.15 (s, 1H, NH), and 8.20 (d, *J* = 8.0 Hz, 2H, Ar–H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub> + CD<sub>3</sub>OD, δ ppm): 53.2, 62.1, 114.8, 125.1, 126.0, 126.1, 127.7, 129.4, 133.5, 136.5, 146.4, 152.3, and 163.0; LC-MS for C<sub>17</sub>H<sub>14</sub>ClN<sub>5</sub>O<sub>4</sub>: [M]<sup>+</sup> 388.

**4.2.19. N-(3-Chlorophenyl)-2-(4-((4-nitrophenoxy)methyl)-1H-1,2,3-triazol-1-yl)acetamide (8f).** The compound **8f** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5f** and alkyne **2c** in 40 min as a white solid; mp: 190–192 °C; FT-IR (cm<sup>-1</sup>): 3327 (N–H stretching), 1683 (C=O stretching); <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>, δ ppm): 5.36 (s, 2H, –NCH<sub>2</sub>CO–), 5.40 (s, 2H, –OCH<sub>2</sub>), 7.15 (d, *J* = 8.0 Hz, 1H, Ar–H), 7.29 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.42 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.78 (s, 1H, Ar–H), 8.23 (d, *J* = 8.0 Hz, 2H, Ar–H), 8.33 (s, 1H, triazole), and 10.81 (s, 1H, NH); <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>, δ ppm): 52.3, 61.9, 115.4, 117.7, 118.9, 123.6, 125.9, 126.8, 130.7, 133.3, 139.9, 141.1, 141.7, 163.4, and 164.7; HRMS calcd for C<sub>17</sub>H<sub>15</sub>ClN<sub>5</sub>O<sub>4</sub> [M + H]<sup>+</sup>: 388.0813 and found 388.0814.

**4.2.20. N-(4-Chlorophenyl)-2-(4-((4-nitrophenoxy)methyl)-1H-1,2,3-triazol-1-yl)acetamide (8g).** The compound **8g** was obtained *via* a 1,3-dipolar cycloaddition reaction between azide **5g** and alkyne **2c** in 37 min as a white solid; mp: 188–190 °C; FT-IR (cm<sup>-1</sup>): 3243 (N–H stretching), 1672 (C=O stretching); <sup>1</sup>H NMR (200 MHz, DMSO-*d*<sub>6</sub>, δ ppm): 5.36 (s, 4H, –NCH<sub>2</sub>CO, –OCH<sub>2</sub>), 7.29 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.39 (d, *J* = 8.0 Hz, 2H, Ar–H), 7.61 (d, *J* = 8.0 Hz, 2H, Ar–H), 8.23 (d, *J* = 8.0 Hz, 2H, Ar–H), 8.32 (s, 1H, triazole), and 10.64 (s, 1H, NH); <sup>13</sup>C NMR (50 MHz, DMSO-*d*<sub>6</sub>, δ ppm): 52.2, 61.9, 115.3, 120.8, 125.8, 126.7, 127.4, 128.8, 137.3, 141.0, 141.6, 163.3, and 164.3; HRMS calcd for C<sub>17</sub>H<sub>15</sub>ClN<sub>5</sub>O<sub>4</sub> [M + H]<sup>+</sup>: 388.0813 and found 388.0819.

### 4.3 Experimental protocol for biological activity

**4.3.1. Antifungal activity.** Antifungal activity was determined by the standard agar dilution method as per the CLSI (formerly, NCCLS) guidelines.<sup>26</sup> The newly synthesized compounds were screened for their *in vitro* antifungal activity against five human pathogenic fungal strains including *Candida albicans* (NCIM 3471), *Fusarium oxysporum* (NCIM 1332), *Aspergillus flavus* (NCIM 539), *Aspergillus niger* (NCIM 1196), and *Cryptococcus neoformans* (NCIM 576). The synthesized compounds and the standard Miconazole were dissolved in DMSO. The medium yeast nitrogen base was dissolved in phosphate buffer of pH 7 and was autoclaved at 110 °C for 10 min. With each set, a growth control without the antifungal agent and solvent control DMSO were included. The fungal strains were freshly subcultured onto sabouraud dextrose agar (SDA) and incubated at 25 °C for 72 h. The fungal cells were suspended in sterile distilled water and diluted to get 10<sup>5</sup> cells per mL. Ten μL of the standardized suspension was inoculated onto the control plates and the media incorporated with the antifungal agents. The inoculated plates were incubated at 25 °C for 48 h. The readings were taken at the end of 48 and 72 h. MIC is the lowest

concentration of the drug preventing the growth of macroscopically visible colonies on the drug-containing plates when there is visible growth on the drug-free control plates.

**4.3.2. In vitro Mtb MABA assay.** The newly synthesized compounds were screened for their *in vitro* antitubercular activity against *Mtb H37Rv* (ATCC 27294) by using the Microplate Alamar Blue Assay (MABA)<sup>30</sup> for the determination of MIC in triplicate. The MIC (in μg mL<sup>-1</sup>) was recorded as the lowest concentration/highest dilution of the compounds/control drugs that completely inhibited the growth of the *Mtb* cultures. The MIC values of the compounds were compared with those of the standard drugs (Isoniazid, Rifampicin, Ethambutol, and Ciprofloxacin). The experimental method for antitubercular activity is briefly described as follows.

Initially, the inoculum was prepared from a fresh LJ medium re-suspended in the 7H9-S medium (7H9 broth, 0.1% casitone, 0.5% glycerol, supplemented oleic acid, albumin, dextrose, and catalase [OADC]), adjusted to a McFarland tube no. 1, and diluted 1 : 20; 100 μL was used as the inoculum. Each drug stock solution was thawed and diluted in 7H9-S at four-fold the final highest concentration tested. Serial two-fold dilutions of each drug were prepared directly in a sterile 96-well microtiter plate using 100 μL 7H9-S. A growth control containing no antibiotic and a sterile control was also prepared on each plate. Sterile water was added to all perimeter wells to avoid evaporation during incubation. The plate was covered, sealed in plastic bags, and incubated at 37 °C in normal atmosphere. After 7 days incubation, 30 μL of alamar blue solution was added to each well, and the plate was re-incubated overnight. A change in the colour from blue (oxidised state) to pink (reduced) indicated the growth of bacteria, and the MIC was defined as the lowest concentration of the drug that prevented this change in colour.

**4.3.3. Antioxidant activity (DPPH radical scavenging).** The hydrogen atom or electron donation ability of some of the compounds was measured from the bleaching of the purple coloured methanol solution of 1,1-diphenyl-1-picrylhydrazyl (DPPH). The spectrophotometric assay uses the stable radical DPPH as a reagent. One mL of various concentrations of the test compounds (5, 10, 25, 50, and 100 μg mL<sup>-1</sup>) in methanol was added to 4 mL of 0.004% IJ w/v methanol solution of DPPH. The reaction mixture was incubated at 37 °C. The scavenging activity on DPPH was determined by measuring the absorbance at 517 nm after 30 min. All tests were performed in triplicate, and the mean values were obtained. The percent of inhibition of free radical production from DPPH was calculated by using the following equation:

$$\% \text{ of scavenging} = [(A_{\text{control}} - A_{\text{sample}}) / (A_{\text{sample}} \times 100)],$$

where  $A_{\text{control}}$  is the absorbance of the control (DPPH radical without test sample) and  $A_{\text{sample}}$  is the absorbance of the test sample (DPPH radical with test sample). The control contained all the reagents except the test samples.

**4.3.4. Homology modeling.** The three-dimensional homology model structure of cytochrome P450 lanosterol 14 $\alpha$ -demethylase of *C. albicans* was predicted using the theoretical methods of protein structure prediction, *i.e.*, comparative



modelling using the Biopredicta module of VLifeMDS 4.3 Pro-Model. The protein sequence of cytochrome P450 lanosterol 14 $\alpha$ -demethylase for *C. albicans* was retrieved from the protein knowledge database Universal Protein Resource (UniProtKB) (<http://www.uniprot.org/>) (accession code: P10613) and its homologous template was identified from the protein structural database, *i.e.*, Protein Data Bank (PDB) using Basic Local Alignment Search Tool for proteins (BLASTp). The homologous template identified using BLASTp solved the three-dimensional crystal structure of human lanosterol 14 $\alpha$ -demethylase (CYP51) complex with azole (PDB ID: 3LD6). To construct a basic framework and to build the loop regions of *C. albicans* cytochrome P450 lanosterol 14 $\alpha$ -demethylase, pairwise sequence alignment and loop refinement of CA-CYP51 (P10613) and human CYP51 (3LD6\_B) were performed. Three-dimensional modelled protein structures were subjected to structural quality check programs such as PRO-CHECK and its other structural validation parameters.<sup>31</sup>

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

One of the authors, SVA, is very much grateful to the Council of Scientific and Industrial Research (CSIR), New Delhi for the award of research fellowship. Authors are also thankful to the Head, Department of Chemistry, Dr Babasaheb Ambedkar Marathwada University Aurangabad for providing laboratory facility.

## Notes and references

- (a) D. Dheer, V. Singh and R. Shankar, *Bioorg. Chem.*, 2017, **71**, 30; (b) A. Massarotti, S. Aprile, V. Mercalli, E. D. Grosso, G. Grosa, G. Sorba and G. C. Tron, *ChemMedChem*, 2014, **9**, 2497; (c) S. G. Agalave, S. R. Maujan and V. S. Pore, *Chem.-Asian J.*, 2011, **6**, 2696; (d) M. Whiting, J. Muldoon, Y. C. Lin, S. M. Silverman, W. Lindstrom, A. J. Olson, H. C. Kolb, M. G. Finn, K. B. Sharpless, J. H. Elder and V. V. Fokin, *Angew. Chem., Int. Ed.*, 2006, **45**, 1435; (e) V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2002, **41**, 2596; (f) C. W. Tornoe, C. Christensen and M. Meldal, *J. Org. Chem.*, 2002, **67**, 3057.
- (a) K. Kempe, A. Krieg, C. R. Becer and U. S. Schubert, *Chem. Soc. Rev.*, 2012, **41**, 176; (b) N. Akeroyd and B. Klumperman, *Eur. Polym. J.*, 2011, **47**, 1207.
- (a) J. J. Bryant and U. H. F. Bunz, *Chem.-Asian J.*, 2013, **8**, 1354; (b) K. D. Hanni and D. A. Leigh, *Chem. Soc. Rev.*, 2010, **39**, 1240.
- (a) G. C. Tron, T. Pirali, R. A. Billington, P. L. Canonico, G. Sorba and A. A. Genazzani, *Med. Res. Rev.*, 2008, **28**, 278; (b) A. D. Moorhouse and J. E. Moses, *ChemMedChem*, 2008, **3**, 715; (c) J. F. Lutz and Z. Zarafshani, *Adv. Drug Delivery Rev.*, 2008, **60**, 958.
- (a) N. Fu, S. Wang, Y. Zhang, C. Zhang, D. Yang, L. Weng, B. Zhao and L. Wang, *Eur. J. Med. Chem.*, 2017, **136**, 596; (b) H. J. Chen, Y. J. Jiang, Y. Q. Zhange, Q. W. Jingb, N. L. Y. Wang, W. N. Zhangb and C. Q. Shenga, *Chin. Chem. Lett.*, 2017, **28**, 913; (c) Z. C. Dai, Y. F. Chen, M. Zhang, S. K. Li, T. T. Yang, L. Shen, J. X. Wang, S. S. Qian, H. L. Zhu and Y. H. Ye, *Org. Biomol. Chem.*, 2015, **13**, 477; (d) Z. Jiang, J. Gu, C. Wang, S. Wang, N. Liu, Y. Jiang, G. Dong, Y. Wang, Y. Liu, J. Yao, Z. Miao, W. Zhang and C. Sheng, *Eur. J. Med. Chem.*, 2014, **82**, 490; (e) N. G. Aher, V. S. Pore, N. N. Mishra, A. Kumar, P. K. Shukla, A. Sharma and M. K. Bhat, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 759.
- (a) B. Zhang, *Eur. J. Med. Chem.*, 2019, **168**, 357; (b) X. L. Wang, K. Wan and C. H. Zhou, *Eur. J. Med. Chem.*, 2010, **45**, 4631.
- (a) K. S. Raju, S. AnkiReddy, G. Sabitha, V. S. Krishna, D. Sriram, K. B. Reddy and S. R. Sagurthi, *Bioorg. Med. Chem. Lett.*, 2019, **29**, 284; (b) S. Zhan, Z. Xu, C. Gao, Q. C. Ren, L. Chang, Z. S. Lv and L. S. Feng, *Eur. J. Med. Chem.*, 2017, **138**, 501; (c) Z. Xu, X. F. Song, Y. Q. Hu, M. Qiang and Z. S. Lv, *Eur. J. Med. Chem.*, 2017, **138**, 66; (d) M. H. Shaikh, D. D. Subhedar, L. Nawale, D. Sarkar, F. A. K. Khan, J. N. Sangshetti and B. B. Shingate, *Med. Chem. Commun.*, 2015, **6**, 1104; (e) S. R. Patpi, L. Pulipati, P. Yogeewari, D. Sriram, N. Jain, B. Sridhar, R. Murthy, D. T. Anjana, S. V. Kalivendi and S. Kantevar, *J. Med. Chem.*, 2012, **55**, 3911.
- (a) P. S. Rao, C. Kurumurthy, B. Veeraswamy, G. S. Kumar, Y. Poornachandra, C. G. Kumar, S. B. Vasamsetti, S. Kotamraju and B. Narsaiah, *Eur. J. Med. Chem.*, 2014, **80**, 184; (b) R. D. Simone, M. G. Chini, I. Bruno, R. Riccio, D. Mueller, O. Werz and G. Bifulco, *J. Med. Chem.*, 2011, **54**, 1565.
- D. R. Buckle, D. J. Outred, C. J. M. Rockell, H. Smith and B. A. Spicer, *J. Med. Chem.*, 1983, **26**, 251.
- (a) Y. Tian, Z. Liu, J. Liu, B. Huang, D. Kang, H. Zhang, E. D. Clercq, D. D. mans, C. Pannecouque, K. H. Lee, C. H. Chen, P. Zhan and X. Liu, *Eur. J. Med. Chem.*, 2018, **151**, 339; (b) G. Wu, W. A. Zalloum, M. E. Meuser, L. Jing, D. Kang, C. Chen, Y. Tian, F. Zhang, S. Cocklin, K. H. Lee, X. Liu and P. Zhan, *Eur. J. Med. Chem.*, 2018, **158**, 478; (c) M. J. Giffin, H. Heaslet, A. Brik, Y. C. Lin, G. Cauvi, C. H. Wong, D. E. McRee, J. H. Elder, C. D. Stout and B. E. Torbett, *J. Med. Chem.*, 2008, **51**, 6263.
- (a) M. Allam, A. K. D. Bhavani, A. Mudiraj, N. Ranjan, M. Thippaana and P. P. Babu, *Eur. J. Med. Chem.*, 2018, **156**, 43; (b) V. Prachayasittikul, R. Pingaew, N. Anuwongcharoen, A. Worachartcheewan, C. Nantasenamat, S. Prachayasittikul, S. Ruchirawat and V. Prachayasittikul, *SpringerPlus*, 2015, **4**, 571; (c) Y. C. Duan, Y. C. Ma, E. Zhang, X. J. Shi, M. M. Wang, X. W. Ye and H. M. Liu, *Eur. J. Med. Chem.*, 2013, **62**, 11; (d) Y. C. Zheng, Y. C. Duan, J. L. Ma, R. M. Xu, X. Zi, W. L. Lv, M. M. Wang, X. W. Ye, S. Zhu, D. Mobley, Y. Y. Zhu, J. W. Wang, J. F. Li, Z. R. Wang, W. Zhao and H. M. Liu, *J. Med. Chem.*, 2013, **56**, 8543.



- 12 X. Wang, Z. C. Dai, Y. F. Chen, L. L. Cao, W. Yan, S. K. Li, J. X. Wang, Z. G. Zhang and Y. H. Ye, *Eur. J. Med. Chem.*, 2017, **126**, 171.
- 13 R. Cha and J. D. Sobel, *Expert Rev. Anti-Infect. Ther.*, 2004, **2**, 357.
- 14 N. H. Georgopapadakou and T. J. Walsh, *Antimicrob. Agents Chemother.*, 1996, **40**, 279.
- 15 D. A. Enoch, H. A. Ludlam and N. M. Brown, *J. Med. Microbiol.*, 2006, **55**, 809.
- 16 (a) G. I. Lepesheva, N. G. Zaitseva, W. D. Nes, *et al.*, *J. Biol. Chem.*, 2006, **281**, 3577; (b) L. Jeu, F. J. Piacenti, A. G. Lyakhovetskiy and H. B. Fung, *Clin. Ther.*, 2003, **25**, 1321; (c) M. A. Pfaller, S. A. Messer, R. J. Hollis and R. N. Jones, *Antimicrob. Agents Chemother.*, 2001, **45**, 2862.
- 17 (a) T. L. Greaves and C. J. Drummond, *Chem. Rev.*, 2008, **108**, 206; (b) R. Sheldon, *Chem. Commun.*, 2001, 2399; (c) P. Wasserscheid and W. Keim, *Angew. Chem., Int. Ed.*, 2000, **39**, 3772; (d) T. Welton, *Chem. Rev.*, 1999, **99**, 2071.
- 18 (a) C. Rub and B. Konig, *Green Chem.*, 2012, **14**, 2969; (b) W. Miao and T. H. Chan, *Org. Lett.*, 2003, **5**, 5003.
- 19 M. A. P. Martins, C. P. Frizzo, A. Z. Tier, D. N. Moreira, N. Zanatta and H. G. Bonacorso, *Chem. Rev.*, 2014, **114**, PR1–PR70.
- 20 (a) A. A. Ali, M. Konwar, M. Chetia and D. Sarma, *Tetrahedron Lett.*, 2016, **57**, 5661; (b) M. Tavassoli, A. L. Isfahani, M. Moghadam, S. Tangestaninejad, V. Mirkhani and I. M. Baltork, *ACS Sustainable Chem. Eng.*, 2016, **4**, 1454; (c) M. Dabiri, S. K. Movahed and D. I. MaGee, *Res. Chem. Intermed.*, 2015, **41**, 3335; (d) H. Singh, J. Sindhu, J. M. Khurana, C. Sharma and K. R. Aneja, *Eur. J. Med. Chem.*, 2014, **77**, 145; (e) A. Z. Ahmady, F. Heidarizadeh and M. Keshavarz, *Synth. Commun.*, 2013, **43**, 2100; (f) A. Marra, A. Vecchi, C. Chiappe, B. Melai and A. Dondoni, *J. Org. Chem.*, 2008, **73**, 2458; (g) Y. B. Zhao, Z. Y. Yan and Y. M. Liang, *Tetrahedron Lett.*, 2006, **47**, 1545.
- 21 (a) N. J. Parmar, H. A. Barad, B. R. Pansuriya and N. P. Talpada, *RSC Adv.*, 2013, **3**, 8064; (b) N. J. Parmar, B. R. Pansuriya, H. A. Barad, B. D. Parmar, R. Kant and V. K. Gupta, *Monatsh. Chem.*, 2013, **144**, 865; (c) N. J. Parmar, R. A. Patel, B. D. Parmar and N. P. Talpada, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 1656; (d) P. Attri and M. Pal, *Green Chem. Lett. Rev.*, 2010, **3**, 249; (e) T. R. Sutariya, B. M. Labana, N. J. Parmar, R. Kant, V. K. Gupta, G. B. Plata and J. M. Padroón, *New J. Chem.*, 2015, **39**, 2657; (f) A. K. Verma, P. Attri, V. Chopra, R. K. Tiwari and R. Chandra, *Monatsh. Chem.*, 2008, **139**, 1041.
- 22 (a) A. B. Danne, A. S. Choudhari, S. Chakraborty, D. Sarkar, V. M. Khedkar and B. B. Shingate, *Med. Chem. Commun.*, 2018, **9**, 1114; (b) S. P. Khare, T. R. Deshmukh, J. N. Sangshetti, V. S. Krishna, D. Sriram, V. M. Khedkar and B. B. Shingate, *ChemistrySelect*, 2018, **3**, 13113; (c) A. B. Danne, A. S. Choudhari, D. Sarkar, J. N. Sangshetti, V. M. Khedkar and B. B. Shingate, *Res. Chem. Intermed.*, 2018, **44**, 6283; (d) M. H. Shaikh, D. D. Subhedar, M. Arkile, V. M. Khedkar, N. Jadhav, D. Sarkar and B. B. Shingate, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 561; (e) M. H. Shaikh, D. D. Subhedar, B. B. Shingate, F. A. K. Khan, J. N. Sangshetti, V. M. Khedkar, L. Nawale, D. Sarkar, G. R. Navale and S. S. Shinde, *Med. Chem. Res.*, 2016, **25**, 790; (f) M. H. Shaikh, D. D. Subhedar, F. A. K. Khan, J. N. Sangshetti and B. B. Shingate, *Chin. Chem. Lett.*, 2016, **27**, 295; (g) M. H. Shaikh, D. D. Subhedar, V. M. Khedkar, P. C. Jha, F. A. K. Khan, J. N. Sangshetti and B. B. Shingate, *Chin. Chem. Lett.*, 2016, **27**, 1058.
- 23 (a) D. D. Subhedar, M. H. Shaikh, B. B. Shingate, L. Nawale, D. Sarkar, V. M. Khedkar, F. A. K. Khan and J. N. Sangshetti, *Eur. J. Med. Chem.*, 2017, **125**, 385; (b) D. D. Subhedar, M. H. Shaikh, F. A. K. Khan, J. N. Sangshetti, V. M. Khedkar and B. B. Shingate, *New J. Chem.*, 2016, **40**, 3047; (c) D. D. Subhedar, M. H. Shaikh, B. B. Shingate, L. Nawale, D. Sarkar and V. M. Khedkar, *Med. Chem. Commun.*, 2016, **7**, 1832; (d) D. D. Subhedar, M. H. Shaikh, L. Nawale, A. Yeware, D. Sarkar, F. A. K. Khan, J. N. Sangshetti and B. B. Shingate, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 2278; (e) D. D. Subhedar, M. H. Shaikh, M. A. Arkile, A. Yeware, D. Sarkar and B. B. Shingate, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 1704; (f) M. H. Shaikh, D. D. Subhedar, F. A. Kalam Khan, J. N. Sangshetti and B. B. Shingate, *Res. Chem. Intermed.*, 2016, **42**, 5115; (g) D. D. Subhedar, M. H. Shaikh, L. Nawale, A. Yeware, D. Sarkar and B. B. Shingate, *Res. Chem. Intermed.*, 2016, **42**, 6607.
- 24 (a) S. T. Dhumal, A. R. Deshmukh, K. R. Kharat, B. R. Sathe, S. S. Chavan and R. A. Mane, *New J. Chem.*, 2019, **43**, 7663; (b) C. P. Kaushik, R. Luxmi, M. Kumar, D. Singh, K. Kumar and A. Pahwa, *Synth. Commun.*, 2019, **49**, 118; (c) G. Wang, Z. Peng, J. Wang, X. Li and J. Li, *Eur. J. Med. Chem.*, 2017, **125**, 423; (d) P. Neeraja, S. Srinivas, K. Mukkanti, P. K. Dubey and S. Pal, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 5212; (e) G. Wang, Z. Peng, J. Wang, J. Li and X. Li, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 5719.
- 25 C. Ferroni, A. Pepe, Y. Sang Kim, S. Lee, A. Guerrini, M. D. Parenti, A. Tesei, A. Zamagni, M. Cortesi, N. Zaffaroni, M. De Cesare, G. L. Beretta, J. B. Trepel, S. V. Malhotra and G. Varchi, *J. Med. Chem.*, 2017, **60**, 3082.
- 26 (a) K. Khan, In vitro and In vivo Screening Techniques for Bioactivity Screening and Evaluation, *Proc. Int. Workshop UNIDO-CDRI*, 1997, p. 210; (b) A. H. Collins, *Microbiological Methods*, Butterworth, London, 2nd edn, 1976.
- 27 M. Burits and F. Bucar, *Phytother. Res.*, 2000, **14**, 323.
- 28 S. Li, D. Li, T. Xiao, S. Zhang, Z. Song and H. Ma, *J. Agric. Food Chem.*, 2016, **23**, 8927.
- 29 V. S. Dofe, A. P. Sarkate, D. K. Lokwani, S. H. Kathwate and C. H. Gill, *Res. Chem. Intermed.*, 2017, **43**, 15.
- 30 S. G. Franzblau, R. S. Witzig and J. C. McLaughlin, *J. Clin. Microbiol.*, 1998, **36**, 362.
- 31 (a) J. N. Sangshetti, F. A. Kalam Khan, R. S. Chouthi, M. G. Damale and D. B. Shinde, *Chin. Chem. Lett.*, 2014, **25**, 1033; (b) R. W. Hooft, G. Vriend, C. Sander and E. E. Abola, *Nature*, 1996, **381**, 272.

