


 Cite this: *RSC Adv.*, 2019, 9, 6935

Correction: Reusable bentonite clay: modelling and optimization of hazardous lead and *p*-nitrophenol adsorption using a response surface methodology approach

 Mohamed Zbair,^{*a} Zakaria Anfar^b and Hassan Ait Ahsaine^b

DOI: 10.1039/c9ra90016k

www.rsc.org/advances

 Correction for 'Reusable bentonite clay: modelling and optimization of hazardous lead and *p*-nitrophenol adsorption using a response surface methodology approach' by Mohamed Zbair *et al.*, *RSC Adv.*, 2019, 9, 5756–5769.

In the published article there was an error in the naming of the pollutant Pb²⁺ in Table 9 on p. 5763. The correct version of this table is shown here below.

 Table 9 Thermodynamic parameters for Pb²⁺ and *p*-nitrophenol adsorption onto BC-500

	ΔH (kJ mol ⁻¹)	ΔS (kJ mol ⁻¹ K ⁻¹)		ΔG (kJ mol ⁻¹)		
				293 K	313 K	333 K
Pb ²⁺	-104.499	-0.201	K_C	184 568 580 -46.365	3 415 381.2 -39.148	1 126 960.8 -38.580
<i>p</i> -Nitrophenol	-132.467	-0.303	K_C	123 915 710.3 -463.657	362 868.435 -33.313	199 963.6695 -33.792

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aLaboratory of Catalysis and Corrosion of Materials, Chouaïb Doukkali University, Faculty of Sciences, El Jadida, BP. 20, El Jadida 24000, Morocco. E-mail: zbair.mohamed@gmail.com

^bMaterials and Environment Laboratory, Ibn Zohr University, Faculty of Sciences, Agadir, Morocco

