 Cite this: *RSC Adv.*, 2019, 9, 22576

DOI: 10.1039/c9ra90055a

www.rsc.org/advances

Correction: Theoretical calculation of a full-dimensional *ab initio* potential energy surface and prediction of infrared spectra for Xe–CS₂

 Miao Qin,^{ab} Xiuchan Xiao^{*ab} and Hua Zhu^c

 Correction for 'Theoretical calculation of a full-dimensional *ab initio* potential energy surface and prediction of infrared spectra for Xe–CS₂' by Miao Qin et al., *RSC Adv.*, 2019, 9, 20925–20930.

The authors regret that eqn (4) was displayed incorrectly in the PDF version of the original article. The correct version of eqn (4) is presented below:

$$\hat{H} = -\frac{1}{2\mu} \frac{\partial^2}{\partial R^2} + \frac{\hat{j}^2}{2I_{v_1+v_3}} + \frac{(\hat{J}-\hat{j})^2}{2\mu R^2} + V_{v_1+v_3}(R, \theta)$$

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aSchool of Architectural and Environmental Engineering, Chengdu Technological University, Chengdu 611730, China. E-mail: shawailsa@sina.cn

^bCenter of Big Data for Smart Environmental Protection, Chengdu Technological University, Chengdu 611730, China

^cSchool of Chemistry, Sichuan University, Chengdu 610064, China