

Cite this: *Chem. Sci.*, 2019, 10, 1117

All publication charges for this article have been paid for by the Royal Society of Chemistry

A low-valent dinuclear ruthenium diazadiene complex catalyzes the oxidation of dihydrogen and reversible hydrogenation of quinones†

Xiuxiu Yang,^a Thomas L. Gianetti,^b Michael D. Wörle,^a Nicolaas P. van Leest,^c Bas de Bruin^c and Hansjörg Grützmacher^a

Received 28th June 2018
Accepted 1st November 2018

DOI: 10.1039/c8sc02864h

rsc.li/chemical-science

The dinuclear ruthenium complex [Ru₂H(μ-H)(Me₂dad)(dbcot)₂] contains a 1,4-dimethyl-diazabuta-1,3-diene (Me₂dad) as a non-innocent bridging ligand between the metal centers to give a [Ru₂(Me₂dad)] core. In addition, each ruthenium is bound to one dibenzo[a,e]cyclooctatetraene (dbcot) ligand. This Ru dimer converts H₂ to protons and electrons. It also catalyzes reversibly under mild conditions the selective hydrogenation of vitamins K₂ and K₃ to their corresponding hydroquinone equivalents without affecting the C=C double bonds. Mechanistic studies suggest that the [Ru₂(Me₂dad)] moiety, like hydrogenases, reacts with H₂ and releases electrons and protons stepwise.

Hydrogenase enzymes are the most efficient biological catalysts for the mutual interconversion of hydrogen to protons and electrons, H₂ ⇌ H⁺ + H⁻.¹⁻⁴ A detailed understanding of the mechanisms of these reactions is necessary for the development of efficient artificial catalytic systems for the use of H₂ as a renewable energy source.⁵ Currently, there are three classes of hydrogenases known which either contain a bimetallic core, [Fe,Fe] or [Fe,Ni], or a single Fe center as active sites.^{3,4} Intensive spectroscopic investigations, including the determination of the structures of several hydrogenases by single-crystal X-ray diffraction methods,⁶⁻⁸ allowed extraction of the essential features needed for activity: (i) redox active metal centers; (ii) an electron reservoir; (iii) a cooperating ligand⁹⁻¹¹ participating reversibly in the heterolytic cleavage/formation of H₂; and (iv) a free coordination site for substrate binding (see a simplified sketch of the active [Fe,Fe] core at the top of Fig. 1).

All hydrogenases contain CO and some CN as archetypical σ-donor/π-acceptor ligands that keep iron in a low spin state and link these enzymes to classical organometallic chemistry. Consequently, the synthesis of hydrogenase model complexes is an intensively investigated topic of organometallic chemistry and some recent relevant examples **A–G** are shown in Fig. 1. Complexes **A** and **B** employ redox non-innocent ligands like

bipyridine¹² or phosphole¹³ in order to mimic Fe₄S₄ ferredoxins as ubiquitous electron reservoirs in enzymes.^{14,15} Both complexes, **A** and **B**, are active electrocatalysts for the production of H₂ from acidic media. To date, the closest model to natural [Fe,Fe] hydrogenases is complex **C**, which can catalyze the oxidation of H₂ to H⁺ in the presence of an oxidant and a base.¹⁶ The discovery that basic sites in a chelating diphosphane ligand greatly enhance the efficiency of heterolytic H₂ splitting and the electrochemical oxidation of H₂^{17,18} has led to the development of iron or nickel complexes like **D** as functional hydrogenase models which achieve truly impressive turnover frequencies (TOFs) of up to 100 000 s⁻¹ for the electrocatalytic production of H₂.¹⁹⁻²³ When tethered to conducting support materials, derivatives of **D** allow fabrication of membrane-electrode assemblies at which H₂ is produced at a very low overpotential.¹⁸ The structural and functional model **E** for [Ni,Fe] hydrogenase was reported, which likewise generates H₂ from mildly acidic solutions with high rates.²⁴ The mononuclear iron model complex **F** in combination with an apoenzyme was used to prepare an [Fe] hydrogenase model.²⁵ Remarkably, this semiartificial enzyme, like its natural counterpart, is able to reversibly hydrogenate methylene tetrahydromethanopterin. Due to the enhanced stability of Ru hydrides, the replacement of Fe with Ru in artificial enzymes has also been investigated.²⁶⁻²⁹ Several dinuclear ruthenium complexes were proposed as hydrogenase mimics.³⁰⁻³² For example, Rauchfuss *et al.* achieved the photochemical addition of H₂ across the Ru–Ru bond in complex **G** and moreover could demonstrate that the terminal hydride ligand in the resulting diruthenium dihydride complex is more easily protonated than the bridging hydride.³⁰

Recently, diazadiene olefin complexes [M(trop₂dad)]ⁿ with low-valent iron or ruthenium centers became accessible

^aDepartment of Chemistry and Applied Biosciences, ETH Zürich, Vladimir-Prelog-Weg 1, 8093 Zürich, Switzerland. E-mail: hgruetzmacher@ethz.ch

^bDepartment of Chemistry and Biochemistry, The University of Arizona, Tucson, Arizona 85721, USA. E-mail: tgianetti@email.arizona.edu

^cVan't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam (UvA), Science Park 904, 1098 XH Amsterdam, The Netherlands

† Electronic supplementary information (ESI) available. CCDC 1535510, 1535543, 1535559, 1535561 and 1852336. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8sc02864h






Fig. 1 Structure of the [Fe,Fe] hydrogenase active site (top) and some hydrogenase mimics (A–G), as well as the illustration of diazadiene olefin complexes related to this work (bottom).

(Fig. 1).^{33,34} In these complexes, the trop_2dad ligand combines the well-established chemical and redox non-innocence of diazadienes (dads) and related ligands^{35–37} with the σ -donor/ π -acceptor properties of olefins.³⁰ Some of these low-valent metal complexes have remarkable properties. For example, $[\text{Ru}^0(\text{trop}_2\text{dad})]$ was found to be an efficient catalyst for the clean conversion of aqueous basic methanol or formaldehyde solutions into H_2 and carbonate.^{34,38}

Herein we report the synthesis of the dinuclear complex $[\text{Ru}_2(\text{Me}_2\text{dad})(\text{dbcot})_2]$ (**3**) which under 1 bar of H_2 is converted to $[\text{Ru}_2\text{H}(\mu\text{-H})(\text{Me}_2\text{dad})(\text{dbcot})_2]$ (**3**($\mu\text{-H}$)) as a fully artificial but functional [Fe,Fe] hydrogenase model (Fig. 2).³⁹ This complex catalyzes the oxidation of hydrogen to protons and electrons as well as the reversible and selective hydrogenation of vitamin K_3 (VK_3) or vitamin K_2 (VK_2) forming dihydrovitamin K_3 (VK_3H_2) or VK_2 hydroquinone (VK_2H_2) without affecting the $\text{C}=\text{C}$ double bonds of these vitamins.

Refluxing dbcot with ruthenium(III) chloride trihydrate in ethanol and THF quantitatively produces the brown coordination polymer $[\text{RuCl}_2(\text{dbcot})]_x$ (**1**), which reacts further with N,N' -dimethyldiaminoethane (Me_2en) in THF and forms the mononuclear complex $[\text{RuCl}_2(\text{Me}_2\text{en})(\text{dbcot})]$ (**2**) in 92% yield (Fig. 2).

The deprotonation of **2** by 2.05 eq. of $\text{KO}t\text{Bu}$ produced a tetranuclear complex, $[\text{K}][\text{Ru}_4(\mu\text{-H})(\text{Me}_2\text{dad})_2(\text{dbcot})_4]$ ($[\text{K}][3_2(\mu\text{-H})]$), in 40% yield as the major species (for detailed spectroscopic data of all isolated complexes reported in this work see the ESI†).³⁰

Reducing the tetranuclear $[\text{K}][3_2(\mu\text{-H})]$ with 3.4 equivalents of K_2C_8 generates the dianionic dinuclear $[\text{K}]_2[\text{Ru}_2(\text{Me}_2\text{dad})(\text{dbcot})_2]$ ($[\text{K}]_2[3]$) in 85% yield. The 2 : 1 ratio of dbcot to Me_2dad and lack of hydride signals in the ^1H NMR spectrum suggests the presence of a Ru dimer complex with no hydrides. The low-frequency shift of the Me_2dad and olefinic protons in $[\text{K}]_2[3]$ compared to $[\text{K}][3_2(\mu\text{-H})]$ is consistent with a more electron rich complex (see ESI Table 2†). The oxidation of $[\text{K}]_2[3]$ with 2 eq. of $[\text{Fc}][\text{PF}_6]$ gives a neutral dinuclear complex $\text{Ru}_2(\text{Me}_2\text{dad})(\text{dbcot})_2$ (**3**) in 77% yield. The ^1H NMR spectra of **3** show sharp resonances at 0 °C, indicating the structure shown in Fig. 2, in which one Ru center is coordinated in a $\kappa^2\text{N,N}$ fashion, and the other in an η^4 fashion by the bridging Me_2dad ligand. But broad signals for the aromatic and olefinic protons are observed at room temperature, indicating molecular dynamics phenomena. Various NMR experiments show that two dynamic processes occur: (i) exchange of the $\kappa^2\text{N,N}/\eta^4$






Fig. 2 Preparation of $[\text{Ru}_2\text{H}(\mu\text{-H})(\text{Me}_2\text{dad})(\text{dbcot})_2]$ ($3(\mu\text{-H})\text{H}$). Illustration of the *dbcot* and *Me*₂*dad* ligand, and six-step synthesis of binuclear complex $3(\mu\text{-H})\text{H}$.

coordination mode of the *Me*₂*dad* ligand between the two Ru centers and (ii) rotation of the *dbcot* ligands (see ESI Fig. 3 for details†). Finally, exposure of **3** to 1 bar of H₂ quickly forms a dinuclear dihydride complex, $\text{Ru}_2\text{H}(\mu\text{-H})(\text{Me}_2\text{dad})(\text{dbcot})_2$ ($3(\mu\text{-H})\text{H}$), in quantitative yield. The two Ru–H resonances at -0.74 ppm and -2.74 ppm in THF-*d*₈ show a coupling of $^2J_{\text{HH}} = 8.7$ Hz, which suggests that both hydrides coordinate to the same Ru atom.⁴⁰ The NOESY spectrum allows us to propose the structure of $3(\mu\text{-H})\text{H}$ as shown in Fig. 2 with a bridging hydride ($\delta = -0.74$ ppm) and a terminal hydride ($\delta = -2.74$ ppm) (see ESI Fig. 5†).

Single crystals of $[\text{K}][3_2(\mu\text{-H})]$ and $[\text{K}]_2[3]$ were obtained from a THF/DME/hexane mixture in the presence of 18-crown-6. Crystals of $3(\mu\text{-H})\text{H}$ were grown by slow evaporation of a saturated benzene solution. All structures were investigated by X-ray diffraction methods and plots are shown in Fig. 3. $[\text{K}][3_2(\mu\text{-H})]$ is a tetranuclear complex which contains two bimetallic ruthenium moieties bridged by a hydride.⁴¹ The complex $[\text{K}]_2[3]$ is best described as an ion pair $[\text{K}]^+[\text{K}][3]^-$ in which the $[\text{K}][3]^-$ anion consists of a sandwich complex with a $(\text{dbcot})\text{RuN}_2\text{C}_2$ unit as the central deck to which a $\text{Ru}(\text{dbcot})$ fragment binds in an η^5 -fashion to one side and a $\text{K}(18\text{-crown-6})$ fragment to the other side. The dinuclear dihydride $3(\mu\text{-H})\text{H}$ contains a bridging and a terminal hydride located on a plane of symmetry including also the two Ru centers, similar to the complex $\text{Ru}_2(\text{S}_2\text{C}_3\text{H}_6)(\mu\text{-H})(\text{H})(\text{CO})_3(\text{PCy}_3)_2$ **G** reported by Rauchfuss *et al.*³⁰ Note that in the Ru_2 complex fragments, the $\kappa^2\text{N,N}/\eta^4$ coordination mode of the diazadiene ligand centers is retained in all complexes. This structural motif is known for related dimeric $[\text{Ru}_2(\text{CO})_5(\text{R}_2\text{dad})]$ complexes.⁴²

The Ru–Ru distances in $[\text{K}][3_2(\mu\text{-H})]$, $[\text{K}]_2[3]$ and $3(\mu\text{-H})\text{H}$ are 2.6947(5), 2.8239(8) and 2.7300(3) Å, respectively, similar to the ones observed in the Ru(I) dimer complexes (2.632–2.937 Å),^{24,42,43} suggesting the presence of Ru–Ru bonds. The longer Ru–Ru distance in $[\text{K}]_2[3]$ (2.8239(8) Å) reflects the highly reduced state of this species. Also, the olefinic bonds in $[\text{K}]_2[3]$ (average of 1.450(3) Å) are longer than the ones in $[\text{K}][3_2(\mu\text{-H})]$ (average of 1.433(3) Å) and $3(\mu\text{-H})\text{H}$ (average of 1.424(3) Å), indicating strong back donation from the Ru–Ru unit into the

π^* -orbitals of the coordinated $\text{C}=\text{C}_{\text{dbcot}}$ bonds. This effect increases with increasing anionic charge of the complex (in free *dbcot*, the average olefin bond length is 1.321 Å). Note that the $\text{C}=\text{C}_{\text{dbcot}}$ bond lengths coordinated to Ru2 are longer than those bound to Ru1, indicating a higher electron density at Ru2. Of special interest are the C–N and C–C bond lengths of the diazadiene ligand because they reflect the oxidation state of the ligand and consequently also of the metal. The neutral diimine form, $\text{RN}=\text{CH}-\text{CH}=\text{NR}$, binds to a low-valent metal center, M^n , and is characterized by short C–N bonds (≈ 1.29 Å) and a long C–C bond (≈ 1.46 Å). With increasing shift of electron density from the metal center to the ligand, the C–N bonds are lengthened while the C–C bond shortens: in the diazadiene radical anion $[\text{RN}=\text{CH}-\text{CH}=\text{NR}]^{\cdot-}$ coordinated to M^{n+1} , C–N ≈ 1.33 Å and C–C ≈ 1.39 Å; in the dianionic bisamido olefin form $[\text{RN}-\text{CH}=\text{CH}-\text{NR}]^{2-}$ coordinated to M^{n+2} , C–N ≈ 1.38 Å and C–C ≈ 1.35 Å.^{36,44–46} The C–N and C–C bonds of the *Me*₂*dad* in $[\text{K}][3_2(\mu\text{-H})]$, $[\text{K}]_2[3]$ and $3(\mu\text{-H})\text{H}$ are approx. 1.38 Å and 1.39 Å, respectively, indicating a reduced form of the ligand. Consequently, the oxidation states at the Ru centers vary between 0 and +1. These data illustrate the redox non-innocent behavior of the *Me*₂*dad* ligand in these complexes.^{36,46} The bridging coordination modes of $2e^-$ reduced *dad* and the closely related pyridine-diimine ligands have been reported before.^{47,48}

The ability to split H₂ into protons and electrons like that of hydrogenases was investigated using the “Rauchfuss test”.¹⁶ In the presence of ten equivalents of PPh_3 and $[\text{Fc}][\text{PF}_6]$, $3(\mu\text{-H})\text{H}$ catalytically splits hydrogen into protons and electrons under 1 bar H₂ at 40 °C, forming protonated triphenylphosphane $[\text{Ph}_3\text{PH}][\text{PF}_6]$, and Cp_2Fe in THF within 1.5 hours (Fig. 4a). More compellingly, $3(\mu\text{-H})\text{H}$ is able to catalyze reversibly the hydrogenation of the biologically relevant vitamins VK_3 and VK_2 (Fig. 4b).^{49,50} Hydrogenation of VK_3 was investigated under 1 bar or 15 bar H₂ pressure, at 40 °C or 70 °C (ESI Table 3, entries 1–3†). With 1 bar H₂, a TON of 252 was achieved. Remarkably, $3(\mu\text{-H})\text{H}$ remained active even after 2 months. At 70 °C and 15 bar H₂, 0.13 mol% $3(\mu\text{-H})\text{H}$ converts 65% of VK_3 to VK_3H_2 to give a TON of 220. Under 15 bar H₂, the TOF values at 40 °C and






Fig. 3 Structures (ORTEP plots) of $[K][3_2(\mu\text{-H})]$ (top), $[K]_2[3]$ (bottom left) and $3(\mu\text{-H})\text{H}$ (bottom right). $[K]$ represents a K^+ cation either coordinated to 18-crown-6 or THF molecules. Thermal ellipsoids are drawn at the 50% probability level. Non-relevant hydrogen atoms, solvent molecules, crown ethers, and potassium cations, which are not part of the anionic Ru_2 complex, have been removed for clarity. Ct_{1-4} are the centroids of the coordinated $C5=C6$, $C13=C14$, $C21=C22$, and $C19=C20$ bonds, respectively. Selected bond distances (Å). $[K][3_2(\mu\text{-H})]$, $Ru1-Ru2$ 2.6947(5), $Ru1-N_{av}$ 2.107(2), $Ru1-ct_{av}$ 2.014(2), $Ru2-N_{av}$ 2.173(2), $Ru2-ct_{av}$ 1.991(3), $N1-C2$ 1.381(3), $C2-C3$ 1.382(4), $C3-N2$ 1.379(2), $C-C_{olefin\ av}$ 1.440(3); $[K]_2[3]$, $Ru1-Ru2$ 2.8239(8), $Ru1-N_{av}$ 2.066(2), $Ru1-ct_{av}$ 1.985(2), $Ru2-N2$ 2.476(2), $Ru2-N2$ 2.293(2), $Ru2-ct_{av}$ 1.970(3), $N1-C2$ 1.399(4), $C2-C3$ 1.406(4), $C3-N2$ 1.390(3), $C-C_{olefin\ av}$ 1.452(4); $3(\mu\text{-H})\text{H}$ (average values from two molecules in one cell), $Ru1-Ru2$ 2.7300(3), $Ru1-N_{av}$ 2.093(2), $Ru1-ct_{av}$ 2.025, $Ru2-N_{av}$ 2.180(2), $Ru2-ct_{av}$ 2.018, $N1-C2$ 1.378(3), $C2-C3$ 1.398(3), $C3-N2$ 1.371(3), $C-C_{olefin\ av}$ 1.425(3).

70 °C are 2.2 and 10 h^{-1} , respectively. VK_2 is a more delicate substrate because it tautomerizes.⁵¹ At room temperature, the hydrogenation of VK_2 is more selectively achieved (ESI Table 3, entry 4†). In addition, the $C=C$ double bonds of VK_2 remain intact and the hydrogenation occurs selectively at the quinone moiety of the substrate. These catalytic reactions can be reversed. The dehydrogenation of VK_3H_2 and VK_2H_2 was tested at 40 °C under Ar (ESI Table 3, entries 5–7†). The TON values of VK_3H_2 and VK_2H_2 are 70 (in 48 h) and 24 (in 40 h), respectively. The TOFs for hydrogenation and dehydrogenation reactions are comparable. A kinetic isotope effect of $k_H/k_D = 1.9(2)$ was measured experimentally for the hydrogenation of VK_3 by measuring k_H and k_D in separate experiments (ESI, part 7.1†).

Stoichiometric reactions were performed and monitored by NMR spectroscopy in order to gain some insights into possible reaction mechanisms. $3(\mu\text{-H})\text{H}$ can be rapidly and quantitatively oxidized at room temperature by ferrocenium salts, $[Fc][X]$ ($X = PF_6$ or OTf), to give the complex $[Ru_2(\mu\text{-H})(Me_2dad)(dbcot)_2]^+[PF_6]^-$ or $[Ru_2(\mu\text{-H})(OTf)(Me_2dad)(dbcot)_2]$ with a bridging hydride (Fig. 5a) in good yield. The complex $[Ru_2(\mu\text{-H})(Me_2dad)(dbcot)_2][PF_6]$ $[3(\mu\text{-H})][PF_6]$ ($Ru-H$, $\delta = -10.34$ ppm)

was characterized by NMR in $THF-d_8$, while $[Ru_2(\mu\text{-H})(OTf)(Me_2dad)(dbcot)_2]$ $[3(\mu\text{-H})(OTf)]$ was isolated in crystalline form. Characterization of this complex by single crystal X-ray diffraction reveals the presence of a structure closely related to $3(\mu\text{-H})\text{H}$ with a triflate anion bound to $Ru1$ instead of the terminal hydride (see Fig. 2 and ESI Fig. 2†). In THF solution, partial dissociation occurs to give $[3(\mu\text{-H})][OTf]$ ($Ru-H$, $\delta = -10.34$ ppm) and $[3(\mu\text{-H})(OTf)]$ ($Ru-H$, $\delta = -10.18$ ppm) in a 1 : 9 ratio. These results indicate that the radical cation salt $[3(\mu\text{-H})\text{H}]^+[X]^-$ as a primary oxidation product rapidly loses hydrogen to give $[3(\mu\text{-H})][X]$ (Fig. 6). The cyclic voltammogram of $3(\mu\text{-H})\text{H}$ shows only one irreversible oxidation peak in THF (ESI Fig. 14†). Our attempts to characterize $[3(\mu\text{-H})\text{H}]^+[X]^-$ by EPR spectroscopy failed so far and oxidation of $3(\mu\text{-H})\text{H}$ with $[Fc][OTf]$ or reduction of $[3(\mu\text{-H})(OTf)]$ with Cp_2Co at 20 K in MeTHF glass afforded only very weak EPR signals characteristic of (a mixture of) metal-centered radical species with g -values in the range of 2.6–1.8 and 2.5–1.7, respectively, which are unlikely to stem from any of the paramagnetic on-cycle catalytic intermediates (such as $[3(\mu\text{-H})\text{H}]^+$ or $[3(\mu\text{-H})]^\cdot$; *vide infra*). Note that






Fig. 6 Proposed mechanisms of H_2 splitting by $3(\mu\text{-H})\text{H}$ using PPh_3 and $[\text{Fc}][\text{PF}_6]$ as the proton and electron acceptors (Cycle A, left), and hydrogenation of vitamin K_n ($n = 2$ or 3) to their hydroquinone analogs by $3(\mu\text{-H})\text{H}$ (Cycle B, right).




Fig. 7 DFT calculation for the activation of H_2 from $3(\mu\text{-H})^+$ in the presence of PPh_3 . The transition state for the deprotonation of the H_2 complex by PPh_3 was not found; however, the direct reaction path from $3(\mu\text{-H})(\text{H}_2)^+$ ($-3.5 \text{ kcal mol}^{-1}$) to $3(\mu\text{-H})\text{H}$ + $[\text{PPh}_3\text{H}]^+$ ($-11.3 \text{ kcal mol}^{-1}$) cannot be excluded.






Fig. 8 DFT calculation for the activation of H₂ from [3(μ-H)]⁺ in the presence of VK₃H⁻.

is still unclear if these radical species are formed *via* direct hydrogen transfer from 3(μ-H)H to VK_n, or *via* initial electron transfer to form the oxidized complex [3(μ-H)H]⁺ and the reduced quinone [VK_n]⁻, followed by proton transfer to form [3(μ-H)][•] and [VK_nH][•]. Subsequent radical recombination with net reduction of the semiquinone radical forms the observed complex [3(μ-H)(VK_nH)]. This complex is subsequently hydrogenated by H₂ to give the product VK_nH₂ and 3(μ-H)H, which is the resting state in this cycle as observed during a single turnover experiment (ESI, Fig. 13†). In this mechanism, the observation of 3(μ-H)H as the only species present after a single turnover experiment is consistent with a less acidic product VK_nH₂ compared to [Ph₃PH]⁺.

The activation steps of H₂ by the Ru₂(Me₂dad) core, in both catalytic reactions, were investigated by DFT calculations without using structural simplification (ΔG values are given here, see Fig. 7 and ESI part 8†). In the absence of other substrates, H₂ adds to [3(μ-H)]⁺ to give [3(μ-H)(H)₂]⁺ in a weakly exergonic reaction. This complex can react with a basic substrate like PPh₃ to give the products 3(μ-H)H and [Ph₃PH]⁺ in an overall exergonic reaction (-11.3 kcal mol⁻¹) although we did not succeed in finding any transition states for this bimolecular reaction. Alternatively, heterolytic cleavage across one Ru-N bond is energetically accessible *via* TS-1 at 15.8 kcal mol⁻¹ to give [3(μ-H)(H)₂]_a⁺. This complex can either rearrange into complex [3(μ-H)(H)₂]_b⁺ to give a hydrogenated C=N bond of the Me₂dad ligand or the acidic NH group is deprotonated to give the products 3(μ-H)H and [Ph₃PH]⁺. Note that the direct heterolytic cleavage of H₂ across a Ru C vector *via* TS-2 and the hydrogenation of one of the C=C_{dbcot} bonds *via* TS-3 are both highly unfavorable processes. With [3(μ-H)]⁺, PPh₃ forms

a stable complex, [3(μ-H)(PPh₃)]⁺ (-22.3 kcal mol⁻¹), which agrees with our assumption that this is the resting state in the catalytic splitting of H₂ with PPh₃ and [Fe][X]. The DFT calculations, for the reaction with VK₃ (Fig. 8), confirm that VK₃H⁻ forms a stable complex, [3(μ-H)(VK₃H)] (-38.9 kcal mol⁻¹), in the reaction with [3(μ-H)]⁺. But in this case, the final products VK₃H₂ and 3(μ-H)H (-45.0 kcal mol⁻¹) are more stable. That is, the adduct [3(μ-H)(VK₃H)] is not the resting state in this cycle but 3(μ-H)H. The cleavage of H₂ involves a classical sigma-bond metathesis transition state TS-4 at 32 kcal mol⁻¹ in which hydrogen is activated across the oxygen atom of the coordinated VK₃H and the ruthenium metal center. Subsequently, TS-4 collapses in a strongly exergonic reaction to give the products VK₃H₂ and 3(μ-H)H. This finding agrees with the measured kinetic isotope effect (1.89), which indicates that the cleavage of H₂ is likely involved in the rate determining step. The activation of H₂ may also occur across a Ru-N bond and involves the second Ru center situated far from the Ru-VK₃H unit but this activation barrier is significantly higher (*E*^a = 42.5 kcal mol⁻¹).

Conclusions

The complex [Ru₂H(μ-H)(Me₂dad)(dbcot)₂], 3(μ-H)H, can be regarded as a fully artificial mimic of binuclear [Fe,Fe] hydrogenases. The Ru₂(Me₂dad) core serves as an electron reservoir and [K]₂[Ru₂(Me₂dad)(dbcot)₂], [K]₂[3], is the most reduced species and [Ru₂(Me₂dad)(dbcot)₂], 3, is the most oxidized species. The Me₂dad ligand binds in an N,N-κ²-fashion to one Ru center, resulting in a five-membered RuN₂C₂ heterocycle with a conjugated π-electron system. Formally, this RuN₂C₂ heterocycle coordinates to a second Ru center as a heteroatom



analogue of archetypical cyclopentadienyl ligands, a structural feature known for dinuclear Ru₂(dad) complexes. However, the Ru₂(dad) complexes reported in the literature carry CO as additional ligands and are likely too unstable to serve as efficient catalysts. In this study, the very rigid and concave shaped dibenzocyclooctatriene, dbcot, was used as a neutral four electron π -donor and π^* -acceptor ligand, which stabilizes all complexes. A Ru–Ru interaction of about 2.7 Å is observed in all complexes, structurally characterized by single crystal X-ray diffraction methods, which is remarkably invariant. Furthermore, the C=C_{dbcot} units bound to the Ru centers do not differ much in length (by 0.05 Å between [K]₂[3] (longest) and [3(μ-H)] [PF₆] (shortest)), indicating that the electron densities at the Ru centers in the various complexes are rather similar. Small structural and electronic variances at the metal centers are also a feature of [Fe,Fe] and [Ni,Fe] hydrogenases.^{3,24,52–58} The complex 3(μ-H)H is a catalyst that splits H₂ into protons and electrons in the presence of PPh₃ and [Fc][PF₆], forming [Ph₃PH][PF₆] and Cp₂Fe (Rauchfuss test for hydrogenase activity), and reversibly and selectively hydrogenates vitamins VK₃ or VK₂, which are natural substrates for the enzyme hydrogen:quinone oxidoreductase. Spectroscopic data strongly suggest that like in hydrogenases, multiple coupled electron and proton transfer steps might be involved in these reactions. Clearly, the observed activities and efficiencies must be significantly improved. But this investigation demonstrates that redox and chemically non-innocent ligands may be key components and their variation may allow further improvements and uncovering of new bearings in synthesizing small molecular hydrogenase mimics.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Dr Reinhard O. Kissner and Jean-Pierre H. Oudsen for their kind help with EPR measurements. This work was supported by the Swiss National Science Foundation and the ETH Zürich. X. Y. is grateful for financial support from the China Scholarship Council. T. L. G. was supported by the ETH Zürich Postdoctoral Fellowship Program, co-funded by the ETH Zurich-Marie Curie action for people (FEL-14 15-1).

Notes and references

- M. Senger, K. Laun, F. Wittkamp, J. Duan, M. Haumann, T. Happe, M. Winkler, U.-P. Apfel and S. T. Stripp, *Angew. Chem., Int. Ed.*, 2017, **56**, 16503–16506.
- M. Winkler, M. Senger, J. Duan, J. Esselborn, F. Wittkamp, E. Hofmann, U.-P. Apfel, S. T. Stripp and T. Happe, *Nat. Commun.*, 2017, **8**, 16115.
- W. Lubitz, H. Ogata, O. Rudiger and E. Reijerse, *Chem. Rev.*, 2014, **114**, 4081–4148.
- C. Tard and C. J. Pickett, *Chem. Rev.*, 2009, **109**, 2245–2274.
- R. Cammack, M. Frey and R. Robson, *Hydrogen as a fuel: Learning from nature*, CRC Press, London and New York, 2002.
- A. Volbeda, M.-H. Charon, C. Piras, E. C. Hatchikian, M. Frey and J. C. Fontecilla-Camps, *Nature*, 1995, **373**, 580–587.
- J. W. Peters, W. N. Lanzilotta, B. J. Lemon and L. C. Seefeldt, *Science*, 1998, **282**, 1853–1858.
- S. Shima, O. Pilak, S. Vogt, M. Schick, M. S. Stagni, W. Meyer-Klaucke, E. Warkentin, R. K. Thauer and U. Ermler, *Science*, 2008, **321**, 572–575.
- J. C. Fontecilla-Camps, A. Volbeda, C. Cavazza and Y. Nicolet, *Chem. Rev.*, 2007, **107**, 4273–4303.
- O. R. Luca and R. H. Crabtree, *Chem. Soc. Rev.*, 2013, **42**, 1440–1459.
- M. Trincado and H. Grützmacher, *Cooperative Catalysis: Designing Efficient Catalysts for Synthesis*, Wiley-VCH Verlag GmbH & Co., Weinheim, 2015, ch. 3, pp. 67–105.
- S. Roy, T. L. Groy and A. K. Jones, *Dalton Trans.*, 2013, **42**, 3843–3853.
- R. Becker, S. Amirjalayer, P. Li, S. Woutersen and J. N. H. Reek, *Sci. Adv.*, 2016, **2**, e1501014.
- R. Lill, *Nature*, 2009, **460**, 831–838.
- S. C. Lee, W. Lo and R. H. Holm, *Chem. Rev.*, 2014, **114**, 3579–3600.
- J. M. Camara and T. B. Rauchfuss, *Nat. Chem.*, 2012, **4**, 26–30.
- C. J. Curtis, A. Miedaner, R. Ciancanelli, W. W. Ellis, B. C. Noll, M. Rakowski DuBois and D. L. DuBois, *Inorg. Chem.*, 2003, **42**, 216–227.
- A. Le Goff, V. Artero, B. Jusselme, P. D. Tran, N. Guillet, R. Métayé, A. Fihri, S. Palacin and M. Fontecave, *Science*, 2009, **326**, 1384–1387.
- T. B. Liu, D. L. DuBois and R. M. Bullock, *Nat. Chem.*, 2013, **5**, 228–233.
- M. L. Helm, M. P. Stewart, R. M. Bullock, M. R. DuBois and D. L. DuBois, *Science*, 2011, **333**, 863–866.
- S. E. Smith, J. Y. Yang, D. L. DuBois and R. M. Bullock, *Angew. Chem., Int. Ed.*, 2012, **51**, 3152–3155.
- A. Dutta, D. L. DuBois, J. A. S. Roberts and W. J. Shaw, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 16286–16291.
- N. Priyadarshani, A. Dutta, B. Ginovska, G. W. Buchko, M. O'Hagan, S. Raugei and W. J. Shaw, *ACS Catal.*, 2016, **6**, 6037–6049.
- D. Brazzolotto, M. Gennari, N. Queyriaux, T. R. Simmons, J. Pecaute, S. Demeshko, F. Meyer, M. Orto, V. Artero and C. Duboc, *Nat. Chem.*, 2016, **8**, 1054–1060.
- S. Shima, D. F. Chen, T. Xu, M. D. Wodrich, T. Fujishiro, K. M. Schultz, J. Kahnt, K. Ataka and X. L. Hu, *Nat. Chem.*, 2015, **7**, 995–1002.
- T. Liu, M. R. DuBois, D. L. DuBois and R. M. Bullock, *Energy Environ. Sci.*, 2014, **7**, 3630–3639.
- G. Gezer, S. Verbeek, M. A. Siegler and E. Bouwman, *Dalton Trans.*, 2017, **46**, 13590–13596.
- T. Matsumoto, Y. Nakaya, N. Itakura and K. Tatsumi, *J. Am. Chem. Soc.*, 2008, **130**, 2458–2459.
- G. M. Chambers, R. Angamuthu, D. L. Gray and T. B. Rauchfuss, *Organometallics*, 2013, **32**, 6324–6329.



- 30 A. K. Justice, R. C. Linck, T. B. Rauchfuss and S. R. Wilson, *J. Am. Chem. Soc.*, 2004, **126**, 13214–13215.
- 31 C. Sommer, C. P. Richers, W. Lubitz, T. B. Rauchfuss and E. J. Reijerse, *Angew. Chem., Int. Ed.*, 2018, **57**, 5429–5432.
- 32 M. Yuki, K. Sakata, Y. Hirao, N. Nonoyama, K. Nakajima and Y. Nishibayashi, *J. Am. Chem. Soc.*, 2015, **137**, 4173–4182.
- 33 C. Lichtenberg, L. Viciu, M. Vogt, R. E. Rodriguez-Lugo, M. Adelhardt, J. Sutter, M. M. Khusniyarov, K. Meyer, B. de Bruin, E. Bill and H. Grützmacher, *Chem. Commun.*, 2015, **51**, 13890–13893.
- 34 R. E. Rodríguez-Lugo, M. Trincado, M. Vogt, F. Tewes, G. Santiso-Quinones and H. Grützmacher, *Nat. Chem.*, 2013, **5**, 342–347.
- 35 F. Richard Keene, *Coord. Chem. Rev.*, 1999, **187**, 121–149.
- 36 K. G. Caulton, *Eur. J. Inorg. Chem.*, 2012, 435–443.
- 37 D. L. J. Broere, R. Plessius and J. I. van der Vlugt, *Chem. Soc. Rev.*, 2015, **44**, 6886–6915.
- 38 M. Trincado, V. Sinha, R. E. Rodriguez-Lugo, B. Pribanic, B. de Bruin and H. Grützmacher, *Nat. Commun.*, 2017, **8**, 14990.
- 39 F. Wittkamp, M. Senger, S. T. Stripp and U. P. Apfel, *Chem. Commun.*, 2018, **54**, 5934–5942.
- 40 D. Schott, C. J. Sleight, J. P. Lowe, S. B. Duckett, R. J. Mawby and M. G. Partridge, *Inorg. Chem.*, 2002, **41**, 2960–2970.
- 41 D. A. Viciu, T. J. Anderson, J. A. Cowan and A. J. Schultz, *J. Am. Chem. Soc.*, 2004, **126**, 8132–8133.
- 42 K. Vrieze, *J. Organomet. Chem.*, 1986, **300**, 307–326.
- 43 E. A. Seddon and K. R. Seddon, *The chemistry of ruthenium*, Elsevier, 1984.
- 44 A. Mederos, S. Domínguez, R. Hernández-Molina, J. n. Sanchiz and F. Brito, *Coord. Chem. Rev.*, 1999, **193–195**, 913–939.
- 45 C. Mealli, A. Ienco, A. D. Phillips and A. Galindo, *Eur. J. Inorg. Chem.*, 2007, **2007**, 2556–2568.
- 46 K. P. Butin, E. K. Beloglazkina and N. V. Zyk, *Russ. Chem. Rev.*, 2005, **74**, 531–553.
- 47 C. Tejel, M. A. Ciriano, M. P. del Río, F. J. van den Bruele, D. G. H. Hetterscheid, N. Tsihchlis i Spithas and B. de Bruin, *J. Am. Chem. Soc.*, 2008, **130**, 5844–5845.
- 48 G. V. Koten and K. Vrieze, in *Advances in Organometallic Chemistry*, ed. F. G. A. Stone and R. West, Academic Press, 1982, vol. 21, pp. 151–239.
- 49 S. Jiang, T. Y. Zhang, X. Zhang, G. H. Zhang and B. Li, *Dalton Trans.*, 2015, **44**, 16708–16712.
- 50 T. Matsumoto, H.-C. Chang, M. Wakizaka, S. Ueno, A. Kobayashi, A. Nakayama, T. Taketsugu and M. Kato, *J. Am. Chem. Soc.*, 2013, **135**, 8646–8654.
- 51 A. M. Swartz, M. Barra and D. Kuntz, *J. Org. Chem.*, 2004, **69**, 3198–3201.
- 52 Y. Higuchi, H. Ogata, K. Miki, N. Yasuoka and T. Yagi, *Structure*, 1999, **7**, 549–556.
- 53 T. Krämer, M. Kampa, W. Lubitz, M. van Gastel and F. Neese, *ChemBioChem*, 2013, **14**, 1898–1905.
- 54 M.-E. Pandelia, H. Ogata and W. Lubitz, *ChemPhysChem*, 2010, **11**, 1127–1140.
- 55 P. Amara, A. Volbeda, J. C. Fontecilla-Camps and M. J. Field, *J. Am. Chem. Soc.*, 1999, **121**, 4468–4477.
- 56 Y. Nicolet, A. L. de Lacey, X. Vernède, V. M. Fernandez, E. C. Hatchikian and J. C. Fontecilla-Camps, *J. Am. Chem. Soc.*, 2001, **123**, 1596–1601.
- 57 S. Stripp, O. Sanganas, T. Happe and M. Haumann, *Biochemistry*, 2009, **48**, 5042–5049.
- 58 P. E. M. Siegbahn, J. W. Tye and M. B. Hall, *Chem. Rev.*, 2007, **107**, 4414–4435.

