Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2019, 10, 2501

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 16th October 2018 Accepted 21st December 2018

DOI: 10.1039/c8sc04601h

rsc.li/chemical-science

A tandem dearomatization/rearomatization strategy: enantioselective N-heterocyclic carbenecatalyzed α-arylation†

Zijun Wu and Jian Wang 回 *

In this study, the first example of the carbene-catalyzed tandem dearomatization/rearomatization reaction of azonaphthalenes with α -chloroaldehydes is described. This protocol enables the efficient assembly of chiral dihydrocinnolinone derivatives in good yields with excellent enantioselectivities (up to 99% ee). Moreover, this strategy enables not only the highly enantioselective NHC-catalyzed nucleophilic aromatic substitution, but also a formal Csp²-Csp³ bond formation.

Advances in reaction development for the stereoselective construction of medicinally important scaffolds or high valueadded chemicals depend on utilizing innovative and longstanding synthesis strategies.1 Over the past decade, a large number of organic transformations driven by N-heterocyclic carbene (NHC) catalysis have been rapidly developed that enabled certain types of reaction manifolds and asymmetric versions of important compounds.² Since the seminar report of the Breslow intermediate in 1958, there has been an everincreasing demand in recruiting novel NHC-bound intermediates for their applications in organic synthesis.³ Among these achievements, the exploration of NHC-bound enolates for new bond formations has attracted considerable attention from the synthetic community due to its high chemical reactivity and remarkable stereo-control. As shown in Fig. 1a, several readily available starting materials (e.g., enals, aldehydes, ketenes, carboxylic acids, esters) have proven to be effective reactants for the *in situ* generation of azolium enolates, thus resulting in a large number of distinct reactions and diversified core skeletons. In 2006, Bode et al. first reported an elegant example of the protonation of electron-deficient enals to produce azolium enolates and the subsequent trapping by N-sulfonyl, α,β unsaturated imines to yield dihydropyridinones.⁴ Rovis et al. have pioneered in using *α*-chloroaldehydes as reactants to prepare azolium enolates for further protonation^{5a} or internal redox reaction.^{5b} Meanwhile, Smith et al. also confirmed that αaroyloxyaldehydes as precursors could convert to azolium enolates in NHC-catalyzed redox esterification, amination, and cycloaddition reactions.⁶ Moreover, the Smith et al.^{7a} and Ye

School of Pharmaceutical Sciences, Key Laboratory of Bioorganic Phosphorous Chemistry and Chemical Biology, Ministry of Education, Tsinghua University, Beijing, 100084, China. E-mail: wangjian2012@tsinghua.edu.cn

[†] Electronic supplementary information (ESI) available. CCDC 1858695. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8sc04601h

This journal is © The Royal Society of Chemistry 2019

(a) Background



To find suitable electrophilic substrates

• To achieve the tandem dearomatizing and rearomatizng chemistry

Fig. 1 Representative transformations of azolium enolates catalyzed by NHCs.



View Article Online

View Journal | View Issue

et al.^{7b,c} have documented that azolium enolates could be readily prepared from ketenes for further annulation. Rovis *et al.*, Chi *et al.*, Scheidt *et al.*, and Ye *et al.* have utilized aliphatic aldehydes,^{sa} activated esters,^{8b} carboxylic acids^{8c,d} and anhydrides^{8e} as suitable reactants to successfully achieve azolium enolates, respectively. In brief, the synthetic utilities of azolium enolates have widely extended to various [2 + 2],^{7a-c,9} [2 + 2 + 2],^{9f,10} $[3 + 2]^{11}$ and $[4 + 2]^{4,6,7c,8a-c,12}$ cycloadditions. These broad diversities may be attributed to the special property of absence of polarity reversal of azolium enolates. Nonetheless, NHC-catalyzed α arylation still remains a formidable challenge for synthetic chemists.

Given the success of *in situ* generated azolium enolates, we anticipated that the desirable NHC-catalyzed α -arylation reaction might partially be attributed to the recruiting of suitable substrates. In 2015, Glorius *et al.* found that the NHC-bound enolates could react with aromatic azomethine imines to generate pyrazolo[5,1-*a*]isoquinolines *via* a direct dearomatization process.¹³ Inspired by this prominent result, we speculated that a matching electrophile (*e.g.*, electron-deficient aromatic compound) may facilitate the initial dearomatization step *via* a nucleophilic attack and C–C bond formation, followed by a rearomatization process, if successfully, to offer the desired α -arylated molecules (Fig. 1b). We also envisioned that the thermodynamics of this intramolecular cycloaddition process probably helps to drive the nucleophilic dearomatization step.

In the process of recruiting suitable substrates, we particularly focused on the azobenzene scaffolds because it has the basic required characteristics of being the matching substrate. First, the azobenzene-type structures can be used as electrophiles in the critical addition step. Second, the azo group can assist in accomplishing the tandem dearomatization-rearomatization process via intermolecular tautomerization. In fact, azobenzene derivatives as synthons have been broadly utilized in metal-catalyzed transformations, as illustrated by rhodiumcatalyzed [3 + 2] cyclization of electron-deficient alkenes with azobenzenes14 (Fig. 2a). Until recently, the Tan et al.15 reported an elegant example of chiral phosphoric acid-catalyzed [3 + 2]cyclization of azonaphthalenes with 2,3-disubstituted indoles (Fig. 2b). In these successful examples, the azo motif within the azonaphthalenes not only works as an electron-withdrawing group to activate the aromatic ring, but also plays a critical role in triggering the rearomatization process. Meanwhile, the NHC-bounded homoenolates, azolium enolates or acyl anions have proven to be effective nucleophiles in catalytic nucleophilic dearomatization.13,16 Inspired by these achievements,15,17 we then set out to explore the feasibility of NHC-catalyzed nucleophilic dearomatization of azonaphthalenes with α chloroaldehyde. Unambiguously, the following challenges need to be overcome in this design: (1) controlling the regioselectivity as the nucleophilic addition of azobenzene derivatives can probably occur at either the N=N double bond^{7c,12h,i,18} or the aromatic ring. (2) Check the compatibility of reactive azonaphthalenes with the nucleophilic NHC-bounded enolate intermediates in the dearomatization reaction. (3) Identify the efficient catalytic system for achieving high enantioselectivity. Herein, we report an unprecedented example of NHC-catalyzed

(a) Glorius, Kim, et al.



(c) This work : tandem dearomatization/rearomatization



Fig. 2 Anticipated cyclization reactions of azobenzene molecules.

 α -arylation of azonaphthalenes with α -chloroaldehydes. It should be mentioned that this process contains a formal Csp²–Csp³ bond formation and starts from an organocatalytic nucleophilic aromatic substitution (Fig. 2c).

To validate the feasibility of the hypothesis, our initial reaction development was conducted with α-chloroaldehyde (1a) and azonaphthalene (2a) in the presence of 20 mol% NHC precatalysts and DIPEA in THF at room temperature (Table 1). Only a trace amount or a moderate yield of 3a was achieved using NHC precatalyst A or B (entries 1 and 2). Gratifyingly, the N-Ph-substituted chiral triazolium catalyst (C2) afforded 3a in 73% yield with 90% ee (Table 1, entry 4). Further investigation indicated that the N-2,4,6-(Me)₃C₆H₂- or N-2,6-(Et)₂C₆H₃substituted chiral triazolium catalyst C1 or C3 gave excellent enantioselectivities, but tolerated moderate yields (entries 3 and 5). Further fine-tuning of other parameters (e.g., base, solvent, and catalyst loading, see ESI[†] for more details) revealed that the optimal condition is a combination of room temperature, 10 mol% C1, 200 mol% DIPEA, and 1.0 mL of ^tBuOMe (Table 1, entry 19).

After establishing the optimal reaction conditions, we examined the substrate scope with respect to various α -chlor-oaldehydes **1** (Fig. 3). The reaction was applicable to α -chlor-oaldehyde derivatives with either aromatic rings or alkyl chains. The electronic natures or substituted patterns on the aromatic



^a Reaction conditions: α-chloroaldehyde 1a (0.20 mmol, 2.0 equiv.), azonaphthalene 2a (0.10 mmol, 1.0 equiv.), cat. (0.02 mmol), base (0.20 mmol), solvent (1.0 mL), 10 h, room temperature. ^b Yields of isolated products after column chromatography. ^c The ee values were determined by HPLC using a chiral stationary phase. d C1 (10 mol%), 16 h. ^e C1 (5 mol%), 24 h.

ring appeared to have limited effects on reaction results, affording the corresponding dihydrocinnolines in yields of 73-88% and ee values between 94 and 99% (3a-3i). When the aldehydes containing alkyl R-substituents located at the acarbon were used, the corresponding dihydrocinnoline derivatives were also afforded with good yields and excellent enantiocontrols (3j-3l). Notably, when α -chloroaldehydes bore functional alkyl subunits (e.g., alkene, chlorine, ether) at the side chain, good yields with excellent enantioselectivities were also regularly observed (3m–3r). Pleasingly, 2-chloro-2phenylacetaldehyde also delivered the corresponding product (3s) in a moderate yield, albeit with a relatively low ee value.

Further investigation on the synthetic feasibility of azonaphthalenes (2) was performed with α -chloroaldehyde (1a), as shown in Fig. 4. The results indicated that this reaction tolerated a diverse array of azonaphthalenes, and afforded the dihydrocinnolines (4) in good to high yields (76-93%) and with high levels of enantio-controls (93-99% ee). When R' substituents were alkyl oxide groups (4a-4d), amino group (4e) or sulfonyl group (4f), the products were obtained in high yields with excellent enantioselectivities. Specifically, the electronic nature or the substitution pattern of the aromatic rings showed limited effects on the reactivity (4g-4l). When an anthracene moiety replaced the naphthalene ring in azonaphthalenes, 76%



mmol), 2a (0.10 mmol), cat. C1 (10 mol%), DIPEA (0.20 mmol), ^tBuOMe (1.0 mL), 12-16 h, room temperature.

yield and 95% ee were still achieved (4m). The absolute configuration of 3r was determined by single-crystal X-ray diffraction analysis,19 and those of other products were assigned by analogy.

Furthermore, to demonstrate the practicality of this catalytic transformation, a gram-scale synthesis of 3a was conducted under the optimal condition. There was almost no change in the reaction yield and enantioselectivity (Fig. 5, eqn (1), 90%, 97% ee), implying that the catalytic tandem reaction of

8





Fig. 4 Scope of azonaphthalenes. ^aReaction conditions: **1a** (0.20 mmol), **2** (0.10 mmol), cat. **C1** (10 mol%), DIPEA (0.20 mmol), ^tBuOMe (1.0 mL), 16 h, room temperature.



Fig. 5 Scale-up synthesis and further transformation.

azonaphthalenes with α -chloroaldehydes can be scaled up. Moreover, the mediated cleavage of **3a** mediated by RANEY®-Ni led to the formation of α -substituted chiral amide (5) in 65% yield with the same ee value (Fig. 5, eqn (2)).

As illustrated in Scheme 1, a postulated mechanism is described. Initial addition of catalyst C1 to α -chloroaldehyde 1 followed by 1,2-H migration generates NHC-bounded enolate I. Nucleophilic addition of enolate I to azonaphthalene (2) results in a formal tandem reaction of dearomatization^{13,16} and



rearomatization along with C–H cleavage and N–H formation to give the thermodynamically stable intermediate **III**. Finally, intramolecular *N*-acylation of **III** leads to the final product **3** or **4** and the NHC catalyst **C1** is released for the next catalytic cycle. Surely, a concerted [4 + 2] annulation mechanism cannot be completely ruled out in this case.

In summary, a novel NHC-catalyzed tandem dearomatization/rearomatization reaction of azonaphthalenes with α -chloroaldehydes has been developed.²⁰ This protocol allows for the rapid assembly of the dihydrocinnolinone scaffolds in good to high yields with high to excellent enantiose-lectivities. Further investigations on the construction of other relevant frameworks as well as a detailed mechanistic study are currently underway in our laboratory.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Generous financial support for this study was provided by the National Natural Science Foundation of China (No. 21672121, 21871160), the "Thousand Plan" Youth Program of China, the Tsinghua University, the Bayer Investigator fellow, the Fellowship of Tsinghua-Peking Centre for Life Sciences (CLS).

Notes and references

- 1 (a) E. N. Jacobsen, A. Pfaltz and H. Yamamoto, *Comprehensive asymmetric catalysis*, Springer, New York, 1999; (b) G. Shang, W. Li, X. Zhang and I. Ojima, *Catalytic asymmetric synthesis*, John Wiley & Sons, New York, 2010.
- 2 For selected recent reviews on NHC catalysis: (a) D. Enders,
 O. Niemeier and A. Henseler, *Chem. Rev.*, 2007, 107, 5606-

5655; (b) A. T. Biju, N. Kuhl and F. Glorius, Acc. Chem. Res., 2011, 44, 1182-1195; (c) D. T. Cohen and K. A. Scheidt, Chem. Sci., 2012, 3, 53-57; (d) X. Bugaut and F. Glorius, Chem. Soc. Rev., 2012, 41, 3511-3522; (e) A. Grossmann and D. Enders, Angew. Chem., Int. Ed., 2012, 51, 314-325; (f) S. J. Ryan, L. Candish and D. W. Lupton, Chem. Soc. Rev., 2013, 42, 4906-4917; (g) S. J. Connon, Angew. Chem., Int. Ed., 2014, 53, 1203-1205; (h) M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, Nature, 2014, 510, 485-496; (i) J. Mahatthananchai and J. W. Bode, Acc. Chem. Res., 2014, 47, 696-707; (j) R. S. Menon, A. T. Biju and V. Nair, Chem. Soc. Rev., 2015, 44, 5040-5052; (k) D. M. Flanigan, F. Romanov-Michailidis, N. A. White and T. Rovis, Chem. Rev., 2015, 15, 9307-9387; (l) R. S. Menon, A. T. Biju and V. Nair, Beilstein J. Org. Chem., 2016, 12, 444-461; (m) M. H. Wang and K. A. Scheidt, Angew. Chem., Int. Ed., 2016, 55, 14912-14922; (n) C. Zhang, J. F. Hooper and D. W. Lupton, ACS Catal., 2017, 7, 2583-2596.

- 3 (*a*) R. Breslow, *J. Am. Chem. Soc.*, 1958, **80**, 3719–3726; experimental evidence for Breslow intermediate, see: (*b*) V. Nair, S. Bindu, V. Sreekumar and N. P. Rath, *Org. Lett.*, 2003, **5**, 665; for isolation and characterization of Breslow intermediate, see: (*c*) A. Berkessel, V. R. Yatham, S. Elfert and J.-M. Neudöfl, *Angew. Chem., Int. Ed.*, 2013, **52**, 11158.
- 4 M. He, J. R. Struble and J. W. Bode, Highly enantioselective azadiene Diels–Alder reactions catalyzed by chiral N-heterocyclic carbenes, *J. Am. Chem. Soc.*, 2006, **128**, 8418–8420.
- 5 (a) N. T. Reynolds, J. Read de Alaniz and T. Rovis, J. Am. Chem. Soc., 2004, **126**, 9518–9519; (b) N. T. Reynolds and T. Rovis, J. Am. Chem. Soc., 2005, **127**, 16406–16407.
- 6 (a) A. Chan and D. Smith, *Chem. Commun.*, 2011, 47, 373– 375; for reviews on α -functionalized aldehydes, see: (b) P.-C. Chiang and J. W. Bode, *N*-Heterocyclic carbine catalysed reactions of α -functionalized aldehydes, ed. B. List and K. Maruoka, Science of Synthesis, Asymmetric Organocatalysis, Thieme, Germany, 2012.
- 7 (a) N. Duguet, C. D. Campbell, A. M. Z. Slavin and A. D. Smith, Org. Biomol. Chem., 2008, 6, 1108-1113; (b)
 Y. R. Zhang, L. He, X. Wu, P. L. Shao and S. Ye, Org. Lett., 2008, 10, 277-280; (c) X.-L. Huang, L. He, P.-L. Shao and S. Ye, Angew. Chem., Int. Ed., 2009, 48, 192-195.
- 8 For pioneering work on aliphatic aldehydes, see: (a) X. Zhao, K. E. Ruhl and T. Rovis, Angew. Chem., Int. Ed., 2012, 51, 12330–12333; for pioneering work on activated esters, see: (b) L. Hao, Y. Du, H. Lv, X. Chen, H. Jiang, Y. Shao and Y. R. Chi, Org. Lett., 2012, 14, 2154–2157; for pioneering work on carboxylic acids, see: (c) A. Lee, A. Younai, C. K. Price, J. Izquierdo, R. K. Mishra and K. A. Scheidt, J. Am. Chem. Soc., 2014, 136, 10589–10592; (d) X.-Y. Chen, Z.-H. Gao, C.-Y. Song, C.-L. Zhang, Z.-X. Wang and S. Ye, Angew. Chem., Int. Ed., 2014, 53, 11611–11615; for pioneering work on anhydrides, see: (e) Z. Jin, S. Chen, Y. Wang, P. Zheng, S. Yang and Y. R. Chi, Angew. Chem., Int. Ed., 2014, 53, 13506–13509.
- 9 For [2 + 2] cycloadditions with azolium enolates, see: (a) L. He, H. Lv, Y.-R. Zhang and S. Ye, J. Org. Chem., 2008, 73,

8101-8103; (b) X.-L. Huang, X.-Y. Chen and S. Ye, J. Org. Chem., 2009, 74, 7585-7587; (c) X.-N. Wang, P.-L. Shao, H. Lv and S. Ye, Org. Lett., 2009, 11, 4029-4031; (d) X.-N. Wang, Y.-Y. Zhang and S. Ye, Adv. Synth. Catal., 2010, 352, 1892-1895; (e) T. Wang, X.-L. Huang and S. Ye, Org. Biomol. Chem., 2010, 8, 5007-5011; (f) X.-N. Wang, L.-T. Shen and S. Ye, Org. Lett., 2011, 13, 6382-6385; (g) T.-Y. Jian, L. He, C. Tang and S. Ye, Angew. Chem., Int. Ed., 2011, 50, 9104-9107.

- 10 X.-N. Wang, L.-T. Shen and S. Ye, *Chem. Commun.*, 2011, 47, 8388–8390.
- 11 For [3 + 2] cycloadditions with azolium enolates, see: (a)
 P.-L. Shao, X.-Y. Chen and S. Ye, Angew. Chem., Int. Ed., 2010, 49, 8412–8416; (b) L. Li, D. Du, J. Ren and Z. Wang, Eur. J. Org. Chem., 2011, 614–618; (c) Q. Ni, H. Zhang, A. Grossmann, C. C. J. Loh, C. Merkens and D. Enders, Angew. Chem., Int. Ed., 2013, 52, 13562–13566; (d)
 Z.-F. Zhang, C.-L. Zhang, Z.-Y. Song, Z.-H. Gao and S. Ye, Chem.-Eur. J., 2018, 24, 8302–8305.
- 12 For selective recent work on [4 + 2] cycloadditions with azolium enolates, see: (a) M. He, G. J. Uc and J. W. Bode, J. Am. Chem. Soc., 2006, 128, 15088-15089; (b) T.-Y. Jian, P.-L. Shao and S. Ye, Chem. Commun., 2011, 47, 2381-2383; (c) J. Kaeobamrung, M. C. Kozlowski and J. W. Bode, Proc. Natl. Acad. Sci. U. S. A., 2010, 107, 20661-20665; (d) X. Fang, X. Chen and Y. R. Chi, Org. Lett., 2011, 13, 4708-4711; (e) Z. Wu, X. Wang, F. Li and J. Wang, Org. Lett., 2015, 17, 3588-3591; (f) T.-Y. Jian, L.-H. Sun and S. Ye, Chem. Commun., 2012, 48, 10907-10909; (g) L. Yang, F. Wang, P. J. Chua, Y. Lv, L.-J. Zhong and G. Zhong, Org. Lett., 2012, 14, 2894–2897; (h) J. E. Taylor, D. S. B. Daniels and A. D. Smith, Org. Lett., 2013, 15, 6058-6061; (i) L. Yang, F. Wang, R. Lee, Y. Lv, K.-W. Huang and G. Zhong, Org. Lett., 2014, 16, 3872-3875; (j) A. Lee and K. A. Scheidt, Chem. Commun., 2015, 51, 3407-3410; (k) X. Chen, R. Song, Y. Liu, C. Y. Ooi, Z. Jin, T. Zhu, H. Wang, L. Hao and Y. R. Chi, Org. Lett., 2018, 19, 5892-5895; (1) S. Lu, J.-Y. Ong, S. B. Poh, T. Tsang and Y. Zhao, Angew. Chem., Int. Ed., 2018, 57, 5714-5719.
- 13 C. Guo, M. Fleige, D. Janssen-Müller, C. G. Daniliuc and F. Glorius, *Nat. Chem.*, 2015, 7, 842–847.
- 14 (a) U. R. Aulwurm, J. U. Melchinger and H. Kisch, Organometallics, 1995, 14, 3385–3395; (b) D. Zhao, S. Vasquez-Cespedes and F. Glorius, Angew. Chem., Int. Ed., 2015, 54, 1657–1661; (c) S. H. Han, N. K. Mishra, H. Jo, Y. Oh, M. Jeon, S. Kim, W. J. Kim, J. S. Lee, H. S. Kim and I. S. Kim, Adv. Synth. Catal., 2017, 359, 2396–2401.
- 15 L.-W. Qi, J.-H. Mao, J. Zhang and B. Tan, *Nat. Chem.*, 2018, **10**, 58–64.
- 16 For NHC-catalyzed dearomatization, see (a) V. Nair, M. Poonoth, S. Vellalath, E. Suresh and R. Thirumalai, J. Org. Chem., 2006, 71, 8964–8965; (b) D. Janssen-Müller, M. Fleige, D. Schlüns, M. Wollenburg, C. G. Daniliuc, J. Neugebauer and F. Glorius, ACS Catal., 2016, 6, 5735– 5739; (c) J.-H. Xu, S.-C. Zheng, J.-W. Zhang, X.-Y. Liu and B. Tan, Angew. Chem., Int. Ed., 2016, 55, 11834–11839; (d) D. M. Flanigan and T. Rovis, Chem. Sci., 2017, 8, 6566–

6569; (e) S. Bera, C. G. Daniliuc and A. Studer, *Angew. Chem., Int. Ed.*, 2017, **56**, 7402–7406; (f) G. D. Carmine, D. Ragno, O. Bortolini, P. P. Giovannini, A. Mazzanti, A. Massi and M. Fogagnolo, *J. Org. Chem.*, 2018, **83**, 2050–2057.

- 17 (a) C. Ma, J.-Y. Zhou, Y.-Z. Zhang, G.-J. Mei and F. Shi, Angew. Chem., Int. Ed., 2018, 57, 5398–5402; (b) C. Li, D.-N. Xu,
 C. Ma, G.-J. Mei and F. Shi, J. Org. Chem., 2018, 83, 9190– 9200.
- 18 A. Chan and K. A. Scheidt, *J. Am. Chem. Soc.*, 2008, **130**, 2740–2741.
- 19 CCDC 1858695 (3r) contains the supplementary crystallographic data for this paper.†
- 20 (a) J. Kaeobamrung, J. Mahatthananchai, P. Zheng and J. W. Bode, *J. Am. Chem. Soc.*, 2010, 132, 8810–8812; (b) J. Kaeobamrung, J. Mahatthananchai and J. W. Bode, *ACS Catal.*, 2012, 2, 494–503.