



Cite this: *Chem. Sci.*, 2019, 10, 6149

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 31st March 2019
Accepted 8th May 2019

DOI: 10.1039/c9sc01574d

rsc.li/chemical-science

Ionization of gold (γ -methoxy)vinyl complexes generates reactive gold vinyl carbene complexes†

Nana Kim and Ross A. Widenhoefer *

Cationic gold vinyl carbene/allylic cation complexes of the form (E) - $[(L)AuC(H)C(H)CAr_2]^+ OTf^-$ { L = IPr, Ar = Ph [(E)-5a], L = IPr, Ar = 4- C_6H_4OMe [(E)-5b], L = $P(t-Bu)_2o$ -biphenyl, Ar = 4- C_6H_4OMe [(E)-5c]} were generated in solution *via* Lewis acid-mediated ionization of the corresponding gold (γ -methoxy)vinyl complexes (E) - $(L)AuC(H)C(H)C(OMe)Ar_2$ at or below -95 °C. Complexes (E)-5b and (E)-5c were fully characterized in solution employing multinuclear NMR spectroscopy, which established the predominant contribution of the aurred allylic cation resonance structure and the significant distribution of positive charge into the γ -anisyl rings. Complex (E)-5b reacted rapidly at -95 °C with neutral two-electron, hydride, and oxygen atom donors exclusively at the C1 position of the vinyl carbene moiety and with *p*-methoxystyrene to form the corresponding vinylcyclopropane. In the absence of nucleophile (E)-5a decomposed predominantly *via* intermolecular carbene dimerization whereas formation of 1-aryl-5-methoxy indene upon ionization of (Z)-(IPr)AuC(H)C(H)C(OMe)(4- C_6H_4OMe) $_2$ [(Z)-6b] implicated an intramolecular Friedel–Crafts or electrocyclic Nazarov pathway for the decomposition of the unobserved vinyl carbene complex (Z)- $[(IPr)AuC(H)C(H)C(4-C_6H_4OMe)_2]^+ OTf^-$ [(Z)-5b].

Introduction

Cationic gold carbene complexes have been invoked as intermediates in a diverse range of gold(I)-catalyzed transformations^{1–3} and, for this reason, considerable effort has been directed toward understanding the structure and reactivity of these complexes.⁴ A particularly important subset of the reactive gold carbene complexes is the gold vinyl carbene/allylic cation complexes, with the potential for both the delocalization of positive charge on the Au, C1, and/or C3 atoms and diverse reaction pathways (Fig. 1). Cationic gold vinyl carbene

complexes have been invoked as intermediates in range of gold-catalyzed transformations employing cyclopropenes,^{5–8} propargyl carboxylates,^{9,10} propargyl ethers,¹¹ enynes,¹² ene-allenes,¹³ vinyl diazo compounds,¹⁴ and norcaradienes as vinyl carbene precursors.¹⁵

Notwithstanding the insight provided through the computational analysis of Toste and Goddard,¹⁶ information regarding the structure and reactivity of cationic gold vinyl carbene complexes is scarce, owing in large part to the absence of suitable model complexes for interrogation.⁴ Only three cationic gold vinyl carbene/allylic cation complexes (1–3) have been characterized, each of which contains either a C1 or multiple C3 oxygen atoms (Fig. 2).^{17,18} As such, neither the electronic structure nor reactivity of complexes 1–3 is likely to mimic that of the reactive vinyl carbenes generated under gold(I) catalysis.^{1–3,5–15} Indeed, well defined, two-coordinate gold carbene complexes that display reactivity analogous to their catalytic counterparts, gold to alkene carbene transfer in




Fig. 1 Potential resonance contributors and composite representation of an unstabilized gold vinyl carbene/allylic cation complex.

French Family Science Center, Duke University, Durham, North Carolina, USA. E-mail: rwidenho@chem.duke.edu

† Electronic supplementary information (ESI) available: Experimental procedures, characterization data, and scans of NMR spectra. CCDC 1897597. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc01574d




Fig. 2 Gold vinyl carbene/allylic cation complexes possessing π -conjugated oxygen atoms, depicted in the form of the predominant oxonium resonance contributor.




particular, are exceedingly rare.^{19,20} Although weakly-stabilized vinyl carbene complexes are known for a range of transition metals,²¹ the properties of cationic gold(I), including high d-electron count, electronegativity²² and poor d → π back bonding,⁴ likely render the electronic structure and reactivity of cationic gold vinyl carbene complexes distinct from these related complexes.

We have recently reported the synthesis of the cationic, two-coordinate gold(I) diphenylallenylidene complex **4** and related complexes, generated *via* the Lewis acid-mediated ionization of the corresponding gold (γ-methoxy)acetylide complexes (Scheme 1),^{23,24} and which represent rare examples of cationic gold carbene complexes lacking π-conjugated heteroatoms.^{25,26} The efficiency and facility of the γ-ionization process suggested that a similar approach might provide access to particularly reactive gold vinyl carbene/allylic cation complexes. Indeed, here we report that ionization of gold (γ-methoxy-γ,γ-diaryl) vinyl complexes generates cationic, two-coordinate gold vinyl carbene complexes that react rapidly at −95 °C with a range of nucleophiles, including alkenes, and that decompose in the absence of nucleophile *via* inter- or intramolecular pathways. The spectroscopy of these complexes points to strong contribution of the auroated allylic cation resonance form and stabilization by the γ-aryl groups.

Results and discussion

Guided by our investigation of cationic gold allenylidene complexes,²³ we targeted the gold γ,γ-(diphenyl)vinylcarbene/allylic cation complex (*E*)-[(IPr)AuC(H)C(H)CPh₂]⁺ OTf[−] [(*E*)-**5a**] for initial investigation (Scheme 2). The requisite gold (γ,γ-diphenyl)vinyl complex (*E*)-(IPr)AuC(H)=C(H)C(OMe)Ph₂ [(*E*)-**6a**] was synthesized in two steps from propargylic ether **7a** *via* zirconium-catalyzed hydroboration²⁷ followed by transmetallation of the resulting boronic ester (*E*)-**8a** with (IPr)AuCl (Scheme 2).²⁸ Treatment of (*E*)-**6a** with trimethylsilyl trifluoromethanesulfonate (TMSOTf) in CD₂Cl₂ at −95 °C led to immediate (<1 min) formation of a bright orange/red solution that became colorless over the course of ~10 s. ¹H NMR analysis of the resulting solution at −95 °C revealed formation of trimethylsilyl methyl ether (TMSOMe) in 92 ± 5% yield, which established that efficient ionization of (*E*)-**6a** had occurred, but no resonances were observed that could be




Scheme 2 Synthesis and thermal decomposition of gold vinyl carbene/allylic cation complex (*E*)-**5a**.

attributed to (*E*)-**5a**. Rather ¹H NMR analysis revealed formation of a ~11 : 1 mixture of (*E*)- and (*Z*)-1,1,6,6-tetraphenylhexa-1,3,5-triene in 70% combined yield, presumably generated *via* homocoupling of the vinyl carbene groups of (*E*)-**5a** (Scheme 2).^{20,29} In an effort to directly observe vinyl carbene (*E*)-**5a**, a solution of (*E*)-**6a** was treated with TMSOTf at −110 °C in CDFCl₂/CD₂Cl₂ and analysed immediately by ¹H NMR spectroscopy. The resulting spectrum was broadened and largely uninterpretable, but displayed a pair of mutually-coupled doublets at δ 10.83 and 8.18 (*J* = 17.5 Hz), assigned to the C1 and C2 vinylic protons, respectively, of (*E*)-**5a** (Table 1).

Table 1 Selected NMR data for gold vinyl carbene complexes (*E*)-**5**. δ Values reported in ppm; *J* values reported in Hz


	(<i>E</i>)- 5a ^a	(<i>E</i>)- 5b ^b	(<i>E</i>)- 5c ^c
δH1	10.83	9.51	8.89
δH2	8.18	7.71	7.83
δC1	—	243.7	252.3
δC2	—	146.9	148.2
δC3	—	185.7	185.6
<i>J</i> _{H1,H2}	17.5	17.8	17.4
<i>J</i> _{C1,C2}	—	48	—
<i>J</i> _{C2,C3}	—	48	—

^a −110 °C in CDFCl₂/CD₂Cl₂. ^b −95 °C in CD₂Cl₂. ^c −90 °C in CD₂Cl₂.



Scheme 1 Synthesis of gold allenylidene complex **4** [IPr = 1,3-bis(2,6-diisopropylphenyl)imidazol-2-ylidene].






Scheme 5 Proposed mechanism of the conversion of (Z)-5b to 9.

end, the gold vinyl complex (Z)-6b was synthesized in two steps from propargyl ether 7b via rhodium-catalyzed, anti-selective hydroboration³⁴ followed by transmetalation of boronic ester (Z)-8b (Scheme 4).

Ionization of (Z)-6b with TMSOTf at $-95\text{ }^{\circ}\text{C}$ formed a pale orange solution that displayed no resonances in the ^1H NMR spectrum that could be attributed to vinyl carbene complex (Z)-5b.[§] Rather, warming this solution above $-50\text{ }^{\circ}\text{C}$ led to formation of the 1-aryl-5-methoxy indene 9 in 58% yield (^1H NMR; Scheme 4). Because 1-aryl indenenes are less stable than are the isomeric 3-aryl indenenes,³⁵ 9 must be formed kinetically, which points to a mechanism for the conversion of (Z)-5b to 9 involving intramolecular Friedel-Crafts alkylation or electrocyclic Nazarov-type cyclization followed by selective protonation of the γ -position of gold allyl intermediate II through an S_{E}' pathway (Scheme 5). Interestingly, (3-phenyl-1*H*-inden-1-yl)gold(i) complexes very similar to intermediate II have been recently isolated via thermal rearrangement of neutral (3,3-diphenylcycloprop-1-en-1-yl)gold(i) complexes and were shown to undergo protodeauration with HCl predominantly through an S_{E}' pathway.^{36,37} The facility of the intramolecular Friedel-Crafts/Nazarov reaction of (Z)-5b and the failure to form 9 in the decomposition of (E)-5b therefore suggests that the energy barrier for *E/Z* isomerization of (E)-5b exceeds that of the lowest energy decomposition pathway available to (E)-5b ($\Delta G^{\ddagger} \approx 16\text{ kcal mol}^{-1}$).

Nucleophilic trapping of a cationic gold vinyl carbene complex represents one of the most common terminating steps of catalytic transformations involving these intermediates.^{5,7,8} In a similar manner, treatment of (E)-5b with either tetrahydrothiophene (THT) or pyridine at $-95\text{ }^{\circ}\text{C}$ in CD_2Cl_2 led to immediate (<1 min) disappearance of the distinctive red color to form pale yellow solutions of the gold allyl carbenoid



Scheme 6 Reactions of (E)-5b with THT and pyridine and interconversion of 10 and 11. Yields determined by ^1H NMR spectroscopy versus internal standard.

complexes 10 and 11, respectively, in >90% yield (^1H NMR; Scheme 6). These complexes were characterized in solution, and in the case of 11, by X-ray crystallography (Fig. 5). For example, selective attack of THT at the C1 vinyl carbene carbon atom of (E)-5b was established by a pair of mutually-coupled doublets at δ 3.22 and 5.60 ($J = 11.5\text{ Hz}$) assigned to the aliphatic C1 and vinylic C2 protons, respectively of 10. In the solid state, 11 adopts a slightly distorted linear conformation ($\text{C1-Au-C28} = 177^{\circ}$) with the C1 allylic carbon atom adopting a distorted sp^3 geometry with a larger Au-C28-C29 angle (115°) and smaller Au-C28-N3 and N3-C28-C29 angles (108°) (Fig. 5). Treatment of 10 (15 mM) with pyridine (1 equiv.) at $0\text{ }^{\circ}\text{C}$ for 2 h led to formation of an equilibrium 1.25 : 1 mixture of 11 : 10 ($K_{\text{eq}} = 1.8 \pm 0.1$; Scheme 6). Kinetic analysis of the conversion of 10 to 11 established first-order approach to equilibrium through 2.5 half-lives with forward and reverse rate constants $k_f = 4.1 \pm 0.9 \times 10^{-4}\text{ s}^{-1}$ and $k_r = 2.3 \pm 0.5 \times 10^{-4}\text{ s}^{-1}$. The first-order kinetic behaviour points to a dissociative mechanism for the interconversion of 10 and 11 involving free (E)-5b.

Oxygen atom transfer to the C1 position of a gold vinyl carbene intermediate has been invoked in the gold-catalyzed reactions of cyclopropenes with diphenylsulfide.^{7,38,39} In a similar manner, treatment of (E)-5b with 4-picoline *N*-oxide (4-






Fig. 5 ORTEP diagram of 11. One of two crystallographically independent molecules is depicted with ellipsoids shown at the 50% probability level with counterion and hydrogen atoms omitted for clarity. Selected bond distances (Å) and bond angles (deg) for 11: Au-C1 = 2.005(4), Au-C28 = 2.067(5), C28-C29 = 1.495(5), C29-C30 = 1.343(7), C28-N3 = 1.515(6), C30-C36 = 1.501(8), C30-C43 = 1.55(1), C1-Au-C28 = 177.4(2), Au-C28-C29 = 115.6(3), Au-C28-N3 = 107.8(3), N3-C28-C29 = 108.3(4), C28-C29-C30 = 125.5(5), C29-C30-C43 = 126.1(9), C29-C30-C36 = 121.6(5), C36-C30-C43 = 112.0(9).






Scheme 7 Reactions of (*E*)-**5b** with 4-picoline-*N*-oxide and HSiEt_3 . Yields determined by ^1H NMR spectroscopy versus internal standard.



Scheme 8 Reaction of (*E*)-**5b** with DSiEt_3 and reaction (*E*)-**5b**- d_1 with HSiEt_3 .

PNO) (1.9 equiv.) in CD_2Cl_2 at -95°C for ~ 1 min led to formation of 3-arylcinnamaldehyde **12** in 88% yield and the 4-picoline gold complex $[(\text{IPr})\text{Au}(4\text{-pic})]^+ \text{OTf}^-$ in quantitative yield (Scheme 7). This outcome is consistent with selective oxygen atom transfer to the C1 carbon atom of the vinyl carbene moiety of (*E*)-**5b** followed by elimination from the unobserved α -oxopyridinium intermediate **III**. Related to this, treatment of



Scheme 9 Reaction of (*E*)-**5b** with *p*-methoxystyrene and independent synthesis of π -(vinyl cyclopropane) complex **15**.

(*E*)-**5b** with triethylsilane at -95°C led to immediate (< 1 min) and selective formation of the cationic gold π -allyl silane complex **13**, presumably *via* hydride addition to the C1 position of (*E*)-**5b** followed by silyldeauration of the resulting gold allyl intermediate through an S_E pathway.⁴⁰ Warming this solution above -40°C led to decomplexation and formation of allylic silane **14** in quantitative yield. Both the reaction of (*E*)-**5b** with DSiEt_3 and the reaction of the deuterated isotopomer (*E*)-**5b**- d_1 with HSiEt_3 at -95°C led selective incorporation of deuterium into the allylic position to form **13**- d_1 as a $\sim 1:1$ mixture of diastereomeric isotopomers, pointing to a stepwise deauration/complexation process (Scheme 8).

Despite the dearth of examples of stoichiometric gold to alkene carbene transfer,^{19,20} perhaps no process is more closely associated with the reactive gold carbene complexes, including vinyl carbene complexes, generated as catalytic intermediates.^{1,2} It is therefore significant that vinyl carbene complex (*E*)-**5b** underwent facile gold to alkene carbene transfer with *p*-methoxystyrene. For example, treatment of (*E*)-**5b** with *p*-methoxystyrene (3 equiv.) at -95°C led to immediate (< 3 min) consumption of (*E*)-**5b** to form the gold π -vinyl cyclopropane complex **15** (Scheme 9). Although ^1H NMR analysis of this solution at -95°C was complicated by the presence of free *p*-methoxystyrene and byproducts, a doublet was observed at δ 4.99 ($J = 10.2$ Hz), assigned to the vinyl proton of **15**, which corresponded to a 75% yield of **15** from (*E*)-**5b** (Scheme 9). Warming this solution of **15** above -40°C led to decomplexation to form free vinyl cyclopropane **16** in 67% NMR yield from (*E*)-**5b** as a $\sim 3:1$ mixture of *cis/trans* isomers (Scheme 9). Assignment of the δ 4.99 resonance to the vinyl proton of **15** was confirmed by independent synthesis of **15** from reaction of **16** (*cis/trans* = 3:1) with a mixture of $(\text{IPr})\text{AuCl}$ and AgSbF_6 in CD_2Cl_2 at -80°C . \blacksquare

Conclusions

We have generated highly reactive gold vinyl carbene/allylic cation complexes *via* low-temperature ionization of gold γ -methoxy-(γ,γ -diaryl)vinyl complexes. Spectroscopic analysis of vinyl carbene/allylic cation complex (*E*)-**5b** established the predominant contribution of the aurated allylic cation canonical form to the electronic structure of the vinyl carbene complex (**C**; Fig. 1) with significant delocalization of positive charge into the γ -anisyl groups. Indeed, the significantly greater stability of the γ,γ -bis(anisyl) vinyl carbene complex (*E*)-**5b** relative to the γ,γ -diphenyl vinyl carbene complex (*E*)-**5a** points to the importance of the *p*-methoxy groups in the stabilization of (*E*)-**5b**.⁴¹ Complex (*E*)-**5b** reacted rapidly at -95°C with neutral two-electron, hydride, and oxygen atom donors exclusively at the C1 position of the vinyl carbene/allylic cation moiety and with *p*-methoxystyrene to form vinyl cyclopropane. These transformations are close or direct analogues to processes attributed to gold vinyl carbene intermediates generated under catalytic conditions.^{1-3,5-15} Two distinct pathways for decomposition of these gold vinyl carbene complexes in the absence of nucleophile were identified; complex (*E*)-**5a** decomposed predominantly *via* intermolecular carbene



homocoupling whereas formation of indene **9** upon ionization of (*Z*)-**6b** directly implicated an intramolecular Friedel–Crafts or electrocyclic Nazarov pathway for the decomposition of the unobserved vinyl carbene complex (*Z*)-**5b**.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank the NSF (CHE-1800273) for support of this research. X-ray crystallography measurements were made by Dr Roger Sommer in the Molecular Education, Technology, and Research Innovation Center (METRIC) at NC State University.

Notes and references

‡ Fürstner has previously reported that attempted synthesis of the PCy₃-analogue of (*Z*)-**5a** via reaction of 1,1-diphenylcyclopropene with (PCy₃)PAuNTf₂ led instead to rapid oligomerization of the cyclopropene.¹⁹ In our hands, attempted synthesis of (*Z*)-**5b** via reaction of 1,1-(bisanisyl)cyclopropene with either (IPr)AuOTf or (IPr)AuCl/AgSbF₆ produced a similar outcome.

§ Two discrete intermediates were detected by ¹H NMR spectroscopy over the temperature range –95 to –30 °C. However, broadening, overlapping resonances, and the dearth of potentially diagnostic resonances precluded extraction of any meaningful information from these data.

¶ Reaction of **16** (*cis/trans* = 3 : 1, 1 equiv.) with a 1 : 1 mixture of (IPr)AuCl and AgSbF₆ in CD₂Cl₂ at –80 °C generated a single resonance [δ 4.99 (br d, *J* = 10.2 Hz)] that could be attributed to **15** and which constituted ~25% of the reaction mixture, the remainder being free **16**. This suggests that either (1) the vinylic ¹H resonances for the various diastereomers of **15** are coincidental or (2) only a single diastereomer of **15** is formed in detectable concentrations from this reaction. Therefore, the 75% yield determined for the conversion of (*E*)-**5b** to **15** (determined by integrating the δ 4.99 resonance relative to internal standard) represents a lower limit for the yield of **15** formed from (*E*)-**5b** and the 67% yield determined for **16** (determined by integrating the vinylic resonances of *cis*- and *trans*-**16** relative to internal standard after warming to room temperature and filtering through silica gel) points to decomposition of **16** under reaction conditions, which was confirmed experimentally. Treatment of (*E*)-**5b** with 1-(*p*-methoxyphenyl)-1,3-butadiene at –95 °C led to immediate consumption of (*E*)-**5b** but produced no products that could be attributed to either cyclopropanation or [4 + 3] cycloaddition.

1 D. Qian and J. Zhang, *Chem. Soc. Rev.*, 2015, **44**, 677–698.

2 (a) L. Liu and J. Zhang, *Chem. Soc. Rev.*, 2016, **45**, 506–516; (b) M. R. Fructos, M. M. Diáz-Requejo and P. J. Pérez, *Chem. Commun.*, 2016, **52**, 7326–7335.

3 (a) R. Dorel and A. M. Echavarren, *Chem. Rev.*, 2015, **115**, 9028–9072; (b) H. Yeom and S. Shin, *Acc. Chem. Res.*, 2014, **47**, 966–977; (c) L. Zhang, *Acc. Chem. Res.*, 2014, **47**, 877–888.

4 (a) R. J. Harris and R. A. Widenhofer, *Chem. Soc. Rev.*, 2016, **45**, 4533–4551; (b) Y. Wang, M. E. Muratore and A. M. Echavarren, *Chem.–Eur. J.*, 2015, **21**, 7332–7339.

5 (a) F. Miege, C. Meyer and J. Cossy, *Beilstein J. Org. Chem.*, 2011, **7**, 717–734; (b) A. Archambeau, F. Miege, C. Meyer and J. Cossy, *Acc. Chem. Res.*, 2015, **48**, 1021–1031.

6 (a) C. Li, Y. Zeng and J. Wang, *Tetrahedron Lett.*, 2009, **50**, 2956–2959; (b) Z.-B. Zhu and M. Shi, *Chem.–Eur. J.*, 2008, **14**, 10219–10222.

7 (a) J. T. Bauer, M. S. Hadfield and A.-L. Lee, *Chem. Commun.*, 2008, 6405–6407; (b) M. S. Hadfield, J. T. Bauer, P. E. Glen and A.-L. Lee, *Org. Biomol. Chem.*, 2010, **8**, 4090–4095.

8 (a) P.-L. Zhu, X.-Y. Tang and M. Shi, *Chem. Commun.*, 2016, **52**, 7245–7248; (b) X. Cheng, L. Zhu, M. Lin, J. Chen and X. Huang, *Chem. Commun.*, 2017, **53**, 3745–3748; (c) R. J. Mudd, P. C. Young, J. A. Jordan-Hore, G. M. Rosair and A.-L. Lee, *J. Org. Chem.*, 2012, **77**, 7633–7639; (d) E. Seraya, E. Slack, A. Ariafard, B. F. Yates and C. J. T. Hyland, *Org. Lett.*, 2010, **12**, 4768–4771; (e) P. C. Young, M. S. Hadfield, L. Arrowsmith, K. M. Macleod, R. J. Mudd, J. A. Jordan-Hore and A.-L. Lee, *Org. Lett.*, 2012, **14**, 898–901; (f) M. S. Hadfield and A.-L. Lee, *Chem. Commun.*, 2011, **47**, 1333–1335.

9 (a) L. Fensterbank and M. Malacria, *Acc. Chem. Res.*, 2014, **47**, 953–965; (b) R. K. Shiroodi and V. Gevorgyan, *Chem. Soc. Rev.*, 2013, **42**, 4991–5001; (c) X.-Z. Shu, D. Shu, C. M. Schienebeck and W. Tang, *Chem. Soc. Rev.*, 2012, **41**, 7698–7711.

10 (a) M. J. Johansson, D. J. Gorin, S. T. Staben and F. D. Toste, *J. Am. Chem. Soc.*, 2005, **127**, 18002–18003; (b) M. Petrović, D. Scarpi, B. Fiser, E. Gómez-Bengoa and E. G. Occhiato, *Eur. J. Org. Chem.*, 2015, 3943–3956; (c) W. Zi, H. Wu and F. D. Toste, *J. Am. Chem. Soc.*, 2015, **137**, 3225–3228; (d) W. Rao and P. W. H. Chan, *Chem.–Eur. J.*, 2014, **20**, 713–718; (e) T. Lauterbach, W. Ganschow, M. W. Hussong, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Adv. Synth. Catal.*, 2014, **356**, 680–686; (f) C. H. Oh, J. H. Kim, L. Piao, J. Yu and S. Y. Kim, *Chem.–Eur. J.*, 2013, **19**, 10501–10505; (g) W. Rao, M. J. Koh, D. Li, H. Hirao and P. W. H. Chan, *J. Am. Chem. Soc.*, 2013, **135**, 7926–7932; (h) T. Lauterbach, S. Gatzweiler, P. Nösel, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Adv. Synth. Catal.*, 2013, **355**, 2481–2487; (i) J. Liu, M. Chen, L. Zhang and Y. Liu, *Chem.–Eur. J.*, 2015, **21**, 1009–1013.

11 (a) E. Jiménez-Núñez, M. Raducan, T. Lauterbach, K. Molawi, C. R. Solorio and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2009, **48**, 6152–6155; (b) A. Rinaldi, M. Petrovic, S. Magnolfi, D. Scarpi and E. G. Occhiato, *Org. Lett.*, 2018, **20**, 4713–4717.

12 (a) M. Gaydou, R. E. Miller, N. Delpont, J. Cecon and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2013, **52**, 6396–6399; (b) J. Carreras, M. Livendahl, P. R. McGonigal and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2014, **53**, 4896–4899; (c) P. Calleja, Ó. Pablo, B. Ranieri, M. Gaydou, A. Pitaval, M. Moreno, M. Raducan and A. M. Echavarren, *Chem.–Eur. J.*, 2016, **22**, 13613–13618.

13 (a) J. H. Lee and F. D. Toste, *Angew. Chem., Int. Ed.*, 2007, **46**, 912–914; (b) B. D. Mokar, P. D. Jadhav, Y. B. Pandit and R.-S. Liu, *Chem. Sci.*, 2018, **9**, 4488–4492.

14 (a) J. Barluenga, G. Lonzi, M. Tomás and L. A. López, *Chem.–Eur. J.*, 2013, **19**, 1573–1576; (b) E. López, G. Lonzi and L. A. López, *Organometallics*, 2014, **33**, 5924–5927; (c) D. Zhang, G. Xu, D. Ding, C. Zhu, J. Li and J. Sun, *Angew. Chem., Int. Ed.*, 2014, **53**, 11070–11074; (d) G. Xu, C. Zhu, W. Gu, J. Li and J. Sun, *Angew. Chem., Int. Ed.*, 2015, **54**, 883–887; (e) J. F. Briones and H. M. L. Davies, *J. Am. Chem. Soc.*, 2013, **135**, 13314–13317; (f) G. Lonzi and L. A. López,



- Adv. Synth. Catal.*, 2013, **355**, 1948–1954; (g) V. V. Pagar, A. M. Jadhav and R.-S. Liu, *J. Am. Chem. Soc.*, 2011, **133**, 20728–20731; (h) A. M. Jadhav, V. V. Pagar and R.-S. Liu, *Angew. Chem., Int. Ed.*, 2012, **51**, 11809–11813; (i) V. V. Pagar and R.-S. Liu, *Angew. Chem., Int. Ed.*, 2015, **54**, 4923–4926; (j) V. V. Pagar, A. M. Jadhav and R.-S. Liu, *J. Org. Chem.*, 2013, **78**, 5711–5716; (k) S. N. Karad and R.-S. Liu, *Angew. Chem., Int. Ed.*, 2014, **53**, 5444–5448.
- 15 (a) B. Herlé, P. M. Holstein and A. M. Echavarren, *ACS Catal.*, 2017, **7**, 3668–3675; (b) X. Yin, M. Mato and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2017, **56**, 14591–14595.
- 16 D. Benitez, N. D. Shapiro, E. Tkatchouk, Y. Wang, W. A. Goddard III and F. D. Toste, *Nat. Chem.*, 2009, **1**, 482–486.
- 17 (a) M. Fañanás-Mastral and F. Aznar, *Organometallics*, 2009, **28**, 666–668; (b) L.-P. Liu, B. Xu, M. S. Mashuta and G. B. Hammond, *J. Am. Chem. Soc.*, 2008, **130**, 17642–17643.
- 18 G. Seidel, R. Mynott and A. Fürstner, *Angew. Chem., Int. Ed.*, 2009, **48**, 2510–2513.
- 19 (a) G. Seidel and A. Fürstner, *Angew. Chem., Int. Ed.*, 2014, **53**, 4807–4811; (b) C. Werlé, R. Goddard and A. Fürstner, *Angew. Chem., Int. Ed.*, 2015, **54**, 15452–15456.
- 20 C. García-Morales, X.-L. Pei, J. M. Sarria Toro and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2019, **58**, 3957–3961.
- 21 (a) P. Binger, P. Müller, R. Benn and R. Mynott, *Angew. Chem., Int. Ed.*, 1989, **28**, 610–611; (b) S. G. Feng, A. S. Gamble and J. L. Templeton, *Organometallics*, 1989, **8**, 2024–2031; (c) S. T. Nguyen, L. K. Johnson, R. H. Grubbs and J. W. Ziller, *J. Am. Chem. Soc.*, 1992, **114**, 3974–3975; (d) L. K. Johnson, R. H. Grubbs and J. W. Ziller, *J. Am. Chem. Soc.*, 1993, **115**, 8130–8145; (e) C. S. Yi, G. L. Geoffroy, C. A. White and A. L. Rheingold, *J. Am. Chem. Soc.*, 1993, **115**, 3806–3807; (f) A. Mayr, M. F. Asaro, T. J. Glines, D. Van Engen and G. M. Tripp, *J. Am. Chem. Soc.*, 1993, **115**, 8187–8196; (g) B. T. Flatt, R. H. Grubbs, R. L. Blanski, J. C. Calabrese and J. Feldman, *Organometallics*, 1994, **13**, 2728–2732; (h) M. A. Esteruelas, F. J. Lahoz, E. Oñate, L. A. Oro and B. Zeier, *Organometallics*, 1994, **13**, 1662–1668; (i) M. A. Esteruelas, F. J. Lahoz, E. Oñate, L. A. Oro, C. Valero and B. Zeier, *J. Am. Chem. Soc.*, 1995, **117**, 7935–7942; (j) C. Gauss, D. Veghini, O. Orama and H. Berke, *J. Organomet. Chem.*, 1997, **541**, 19–38; (k) J. Foerstner, A. Kakoschke, D. Stellfeldt, H. Butenschön and R. Wartchow, *Organometallics*, 1998, **17**, 893–896.
- 22 (a) S. G. Bratsch, *J. Chem. Educ.*, 1988, **65**, 34; (b) J. E. Huheey, E. A. Keiter and R. L. Keiter, *Inorganic Chemistry: Principles of Structure and Reactivity*, HarperCollins, New York, USA, 4th edn, 1993.
- 23 N. Kim and R. A. Widenhoefer, *Angew. Chem., Int. Ed.*, 2018, **57**, 4722–4726.
- 24 For examples of heteroatom-stabilized gold allenylidene complexes see: (a) M. M. Hansmann, F. Rominger and A. S. K. Hashmi, *Chem. Sci.*, 2013, **4**, 1552–1559; (b) X.-S. Xiao, W.-L. Kwong, X. Guan, C. Yang, W. Lu and C.-M. Che, *Chem.–Eur. J.*, 2013, **19**, 9457–9462; (c) L. Jin, M. Melaimi, A. Kostenko, M. Karni, Y. Apeloig, C. E. Moore, A. L. Rheingold and G. Bertrand, *Chem. Sci.*, 2016, **7**, 150–154; (d) X.-S. Xiao, C. Zou, X. Guan, C. Yang, W. Lu and C.-M. Che, *Chem. Commun.*, 2016, **52**, 4983–4986.
- 25 (a) M. W. Hussong, F. Rominger, P. Krämer and B. F. Straub, *Angew. Chem., Int. Ed.*, 2014, **53**, 9372–9375; (b) R. J. Harris and R. A. Widenhoefer, *Angew. Chem., Int. Ed.*, 2015, **54**, 6867–6869.
- 26 R. J. Harris and R. A. Widenhoefer, *Angew. Chem., Int. Ed.*, 2014, **53**, 9369–9371.
- 27 C. G. Frost, H. J. Edwards, S. D. Penrose and R. Gleave, *Synthesis*, 2010, 3243.
- 28 (a) D. V. Partyka, M. Zeller, A. D. Hunter and T. G. Gray, *Angew. Chem., Int. Ed.*, 2006, **45**, 8188–8191; (b) A. S. K. Hashmi, T. D. Ramamurthi and F. Rominger, *J. Organomet. Chem.*, 2009, **694**, 592–597.
- 29 For related homocoupling processes of gold carbene complexes see ref. 20 and J. M. Sarria Toro, C. García-Morales, M. Raducan, E. S. Smirnova and A. M. Echavarren, *Angew. Chem., Int. Ed.*, 2017, **56**, 1859–1863.
- 30 (a) H.-U. Siehl, F.-P. Kaufmann and K. Hori, *J. Am. Chem. Soc.*, 1992, **114**, 9343–9349; (b) R. Jost, J. Sommer, C. Engdahl and P. Ahlberg, *J. Am. Chem. Soc.*, 1980, **102**, 7663–7667.
- 31 (a) C. U. Pittman Jr and W. G. Miller, *J. Am. Chem. Soc.*, 1973, **95**, 2947–2956; (b) N. C. Deno, C. U. Pittman Jr and J. O. Turner, *J. Am. Chem. Soc.*, 1965, **87**, 2153–2157; (c) W. G. Miller and C. U. Pittman Jr, *J. Org. Chem.*, 1974, **39**, 1955–1956; (d) D. M. Dytneriski, K. Ranganaykublu, P. B. Singh and T. S. Sorensen, *Can. J. Chem.*, 1982, **60**, 2993–3004; (e) G. A. Olah, G. Asensio and H. Mayr, *J. Org. Chem.*, 1978, **43**, 1518–1520; (f) J. Wang, L. Zhang, Y. Jing, W. Huang and X. Zhou, *Tetrahedron Lett.*, 2009, **50**, 4978–4982.
- 32 H. Mayr, W. Förner and P. von Ragué Schleyer, *J. Am. Chem. Soc.*, 1979, **101**, 6032–6040.
- 33 (a) M. S. Hadfield, L. J. L. Häller, A.-L. Lee, S. A. Macgregor, J. A. T. O'Neill and A. M. Watson, *Org. Biomol. Chem.*, 2012, **10**, 4433–4440; (b) N. A. Rajabi, M. J. Atashgah, R. BabaAhmadi, C. Hyland and A. Ariaifard, *J. Org. Chem.*, 2013, **78**, 9553–9559.
- 34 T. Ohmura, Y. Yamamoto and N. Miyaura, *J. Am. Chem. Soc.*, 2000, **122**, 4990–4991.
- 35 E. C. Friedrich and D. B. Taggart, *J. Org. Chem.*, 1975, **40**, 720–723.
- 36 F. F. Mulks, P. W. Antoni, F. Rominger and A. S. K. Hashmi, *Adv. Synth. Catal.*, 2018, **360**, 1810–1821.
- 37 For the protodeauration of gold σ -allyl complexes through an S_E' pathway see ref. 8e and: (a) A. S. K. Hashmi, A. M. Schuster, S. Litters, F. Rominger and M. Pernpointner, *Chem.–Eur. J.*, 2011, **17**, 5661–5667; (b) R. BabaAhmadi, P. Ghanbari, N. A. Rajabi, A. S. K. Hashmi, B. F. Yates and A. Ariaifard, *Organometallics*, 2015, **34**, 3186–3195.



- 38 Analogous oxidations of gold carbene moieties have been observed under catalytic conditions³⁹ and from well defined gold carbene complexes.^{20,26}
- 39 C. A. Witham, P. Mauleón, N. D. Shapiro, B. D. Sherry and F. D. Toste, *J. Am. Chem. Soc.*, 2007, **129**, 5838–5839.
- 40 For an example of silyldeauration implicated in a catalytic transformation see: I. Nakamura, T. Sato, M. Terada and Y. Yamamoto, *Org. Lett.*, 2007, **9**, 4081–4083.
- 41 A similar conclusion was reached by Fürstner regarding the gold bis(anisyl) carbene complex $(\text{PCy}_3)\text{AuC}(p\text{-C}_6\text{H}_4\text{OMe})_2$.¹⁹

