

CORRECTION

View Article Online  
View Journal | View Issue



Cite this: *Chem. Sci.*, 2019, 10, 5849

## Correction: Identification of selective protein–protein interaction inhibitors using efficient *in silico* peptide-directed ligand design†

Andrew M. Beekman,<sup>\*a</sup> Marco M. D. Cominetti,<sup>a</sup> Samuel J. Walpole,<sup>a</sup>  
Saurabh Prabhu,<sup>a</sup> Maria A. O'Connell,<sup>a</sup> Jesus Angulo<sup>a</sup> and Mark Searcey<sup>\*ab</sup>

DOI: 10.1039/c9sc90088h

[www.rsc.org/chemicalscience](http://www.rsc.org/chemicalscience)

Correction for 'Identification of selective protein–protein interaction inhibitors using efficient *in silico* peptide-directed ligand design' by Andrew M. Beekman *et al.*, *Chem. Sci.*, 2019, DOI: 10.1039/c9sc00059c.

The authors regret that the structures of compounds 2, 3, 10 and 11 are incorrect in the original article. The correct structures are presented in the updated version of Table 1 below.

The original ESI was replaced by a correspondingly revised version on 1st May 2019 to reflect these changes.











<sup>a</sup>School of Pharmacy, University of East Anglia, Norwich Research Park, Norwich, Norfolk, NR47TJ, UK. E-mail: A.Beekman@uea.ac.uk; M.Searcey@uea.ac.uk

<sup>b</sup>School of Chemistry, University of East Anglia, Norwich Research Park, Norwich, Norfolk, NR47TJ, UK

† Electronic supplementary information (ESI) available: See DOI: 10.1039/c9sc00059c



**Table 1** IC<sub>50</sub> values for inhibition of the binding of FAM-p4 to *hDM2* and *hDMX* of small molecules.<sup>a</sup> A green background indicates the compound was designed for and was selective to the same protein. Yellow indicates the compound is a dual inhibitor. A red background indicates the compound was designed for one protein but was selective for the other protein

Structure	<i>hDMX</i> FA IC <sub>50</sub> (μM) [95% CI]	<i>hDM2</i> FA IC <sub>50</sub> (μM) [95% CI]	Structure	<i>hDMX</i> FA IC <sub>50</sub> (μM) [95% CI]	<i>hDM2</i> FA IC <sub>50</sub> (μM) [95% CI]
	0.013 [0.003, 0.046]	0.24 [0.16, 0.36]		>100	2.46 [1.01, 4.70]
	0.019 [0.005, 0.071]	0.10 [0.03, 0.76]		>100	5.77 [4.00, 8.37]
	1.28 [0.64, 2.52]	8.00 [6.12, 20.06]		>100	12.42 [8.61, 21.23]
	1.49 [1.06, 2.10]	>100		10.97 [8.28, 19.66]	22.44 [17.92, 29.60]
	0.97 [0.79, 1.44]	1.74 [0.89, 3.38]		1.17 [1.02, 4.20]	>100

<sup>a</sup> IC<sub>50</sub> values determined by non-linear regression of at least three independent experiments (see ESI, pg 10). Errors are 95% confidence intervals (CI). Fmoc, 9-fluorenylmethylcarbonyl.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

