ChemComm

COMMUNICATION

Cite this: *Chem. Commun.*, 2020, 56, 14199

Received 21st September 2020, Accepted 20th October 2020

DOI: 10.1039/d0cc06353c

rsc.li/chemcomm

A small-molecule probe for monitoring binding to prolyl hydroxylase domain 2 by fluorescence polarisation†

Zhihong Li,^a Shuai Zhen,^a Kaijun Su,^a Anthony Tumber,^b Quanwei Yu,^a Ying Dong,^a Michael McDonough, \mathbb{D}^{b} Christopher J. Schofield $\mathbb{D}^{*^{\text{b}}}$ and Xiaojin Zhang $\mathbb{D}^{*^{\text{ab}}}$

Inhibition of the dioxygen sensing hypoxia-inducible factor prolyl hydroxylases has potential therapeutic benefit for treatment of diseases, including anaemia. We describe the discovery of a smallmolecule probe useful for monitoring binding to human prolyl hydroxylase domain 2 (PHD2) via fluorescence polarisation. The assay is suitable for high-throughput screening of PHD inhibitors with both weak and strong affinities, as shown by work with clinically used inhibitors and naturally occurring PHD inhibitors.

Hypoxia in animals is associated with diseases including anaemia and cancer. The α , β -heterodimeric hypoxia-inducible transcription factors (HIFs) are of central importance in the chronic response to hypoxia.¹ HIF- α levels are regulated in a dioxygen availability limited manner by the C-4 hydroxylation of conserved prolyl-residues, a post-translational modification that signals for HIF- α degradation via the ubiquitin-proteasome machinery.² In normoxia, HIF- α levels rise, it dimerises with HIF- β , and HIF-mediated transcription is increased.² The Fe(II) and 2-oxoglutarate (2OG) dependent HIF-a prolyl hydroxylases (PHD1–3 in humans) (Fig. 1A) act as hypoxia sensors for the HIF system.³ Since HIF targets induce erythropoietin and vascular endothelial growth factor, PHD inhibition is of therapeutic interest; 4 PHD inhibitors that are 2OG competitors have recently been approved for the treatment of anaemia in chronic kidney disease and others are in clinical trials.⁵ Reduced activity of PHD and other 2OG oxygenases as a consequence of metabolic reprogramming in cancer cells, via PHD inhibition due to elevated succinate, fumarate, and/ or 2-hydroxyglutarate levels, is proposed to be important in disease progression.⁶ There is thus a need for efficient and

^a Laboratory of Drug Design and Discovery, Department of Chemistry, China

accurate assays for monitoring both the catalytic activity of the PHDs and the affinity of natural compounds to their active sites.

We and others have reported various assays for measuring PHD catalysis, $7-11$ including by monitoring 2OG consumption⁷ and by monitoring substrate depletion/product formation by mass spectrometry (MS)/nuclear magnetic resonance (NMR),^{8,9} or antibody based methods.¹⁰ Whilst these assays can be useful for identifying potent PHD inhibitors, they do not provide information on binding constants, in particular to the biologically relevant PHD–Fe complexes. This is important because substantial conformational changes occur during PHD catalysis and, likely, during the binding of some inhibitors.¹² Turnover COMMUNICATION

(a) Check for undates

Communications

Communi

Pharmaceutical University, Nanjing 211198, China. E-mail: zxj@cpu.edu.cn

 b Chemistry Research Laboratory, Department of Chemistry, University of Oxford, 12 Mansfield Road, Oxford OX1 3TA, UK.

E-mail: christopher.schofield@chem.ox.ac.uk

[†] Electronic supplementary information (ESI) available: Details of experimental procedures and additional experiments. See DOI: 10.1039/d0cc06353c

assays are also not ideal for monitoring the binding of compounds with weak affinities, e.g., fragments and natural metabolites, including tricarboxylic acid (TCA) cycle related compounds.¹³ Here we report the discovery of a fluorescence polarisation (FP) utilising affinity assay for PHD2; importantly, the assay employs the metal $Fe(\theta)$ cofactor and is suitable for screening metal-binding fragment-like inhibitors.

The design of the probe was based on the recently reported triazole containing PHD inhibitor 1 (Fig. 1B), which is a potent PHD2 inhibitor with an IC_{50} value of 78 nM as assessed by an FITC-labelled HIF-1a peptide probe-based FP binding assay, both against isolated PHD2 and in cells, and which manifests a promising safety profile.¹⁴ Modelling studies based on crystal structures of PHD2 in complex with other inhibitors, imply that 1 chelates the active site $Fe(II)$ in a bidentate manner and its glycine side chain occupies the region normally occupied by the methylene and C-5 carboxylate of 2OG (Fig. $S1$, ESI_1^+).¹⁵ Notably, these studies implied that the meta-position of the chlorophenyl group extends towards the outside of the active site (Fig. 1C), suggesting that functionalisation at this position with a fluorescent reporter group, such as fluorescein isothiocyanate (FITC) could be achieved without substantially perturbing active site binding (Fig. 1D). Communication

components with weak artificies. z_2 -fragments and taxental components are the components are the components are the components are the common points are the common points of the properties are realized a

Probe 12 was synthesised from phenol derivative 2 (Fig. 2), via 3, which was obtained by alkylation of 2 with 1,2-dibromoethane. Methyl ester hydrolysis of 3 yielded 4, which was condensed with N-Boc-1,6-hexanediamine (5) to yield amide 6, which was reacted with NaN_3 to give 7. Microwave promoted Huisgen click reaction between 7 and 8^{14} gave triazole 9, which was hydrolysed by base to give 10. Boc-deprotection (CF_3CO_2H) gave 11, which was derivatised with the reporter group (FITC), then subjected to ester hydrolysis, producing the designed functional PHD2 probe 12.

Fig. 2 Synthesis of PHD probe 12. Reagents and conditions: (a) 1,2 dibromoethane, K₂CO₃, CH₃CN, 60 °C, 9 h, 62%; (b) 30% KOH, 2 h, 91%; (c) EDCI, HOBT, TEA, CH₂Cl₂, rt, 5 h, 54%; (d) NaN₃, DMF, 100 °C, 6 h, 91%; (e) TBAF, DIPEA, CuI, MeOH, 80 °C, 4 h, 59%; (f) LiOH, H₂O, THF, 30 °C, 92%; (g) CF₃COOH, CH₂Cl₂, rt, overnight, 56%; (h) FITC, HBTU, DIPEA, DMF, rt; then LiOH, H₂O, THF, 30 °C, 45% (structure of probe 12 see Fig. 3A).

Fig. 3 (A) Structure of probe 12. (B) Dose–response inhibition curves of probe 12 and FG-4592 based on the SPE MS assay;¹⁶ (C) dose–response binding curve of probe 12 based on the titration FP assay (no additional iron ion and 2OG were added). See ESI† for details.

To investigate the effects of introducing the linker and bulky FITC group on the probe's ability to bind PHD2, the affinity of 10 with PHD2 was evaluated using our reported FP assay, which manifested an IC₅₀ of 310.6 \pm 9.7 nM (Fig. S2, ESI†). This observation suggests that the introduction of the linker and FITC group does not ablate the strong affinity of the active site binding elements of the probe, though its affinity is reduced somewhat. Encouraged by this result, we then investigated the biological applications of probe 12. First, a solid phase extraction (SPE)-MS coupled assay¹⁶ was performed to evaluate the potency of probe 12 for PHD2 inhibition. The resultant IC_{50} was 166 \pm 3 nM, which compares favourably with a PHD inhibitor approved for clinical use to treat anaemia associated with chronic kidney disease ${\rm (CKD)}^{17}$ (FG-4592, IC₅₀: 2587 \pm 20 nM) (Fig. 3B). The direct binding affinity of probe 12 to PHD2 was determined by monitoring the FP signals; the resulting EC_{50} value of probe 12 that binds to PHD2 is 27.4 \pm 1.2 nM (Fig. 3C).

The selectivity of probe 12 towards selected other human 2OG oxygenases, such as factor inhibiting HIF (FIH) and JmjC histone N^{ϵ} -methyl lysine demethylases (KDMs), was then evaluated using established assays (Table 1). The results indicated that probe 12 is highly selective for PHD2 over the tested 2OG dependent KDMs. Probe 12 was selective for PHD2 over FIH (which regulates the transcriptional activity of HIF), though to a lesser extent.

We then developed a small-molecule fluorescent probebased FP assay for PHD2. The influence of the polarity of the medium on the fluorescence intensity of probe 12 was

Table 1 IC_{50} values (nM) of probe 12 towards PHD2, and other 2OG oxygenases: FIH and typical JmjC KDMs

	Cpd $PHD2^a$ $FH1^a$	$KDM4E^a$	$KDM3A^{b}$ $KDM4A^{b}$	
12		166 ± 3 841 ± 16 40900 ± 350 >100000 >100000		

 a SPE MS assay.¹⁶ b AlphaScreen assay.¹⁰ Both assays are based on the hydroxylation of HIF-1a peptides. See ESI for details.

Fig. 4 Spectroscopic properties of probe 12. (A) Emission spectra of probe 12. (B) Excitation spectra of probe 12.

investigated by adding dioxane or DMSO in PBS buffer measured at a λ_{max} of 520 nm, respectively (Fig. S3, ESI†). The excitation and emission spectra of probe 12 are characterised by maxima λ = 505 nm (excitation) and 520 nm (emission) (Fig. 4). The results indicated that probe 12 has acceptable spectroscopic properties in the presence of dioxane or DMSO in buffer when measured at a λ_{max} of 520 nm (Fig. S3, ESI†). Thus, probe 12 was considered suitable for further development as a tool for studies on PHD2.

We optimised use of probe 12 for efficient affinity-based FP method for quantitative screening of PHD2 inhibitors. The developed method does not require the addition of 2OG and can employ Fe(π) that copurifies with PHD2,²⁰ though apo-protein and added metal ions can also be used (Fig. S4 and S5, ESI†). The assay employs low concentrations of probe 12 (30 nM) and PHD2 (20 nM) (Fig. 5A), the latter is a level much lower than most previously reported assays, e.g. employing matrix-assisted laser desorption/ionization time of flight mass spectrometry (MALDI-TOF MS) (50 μ M),⁸ NMR (10 μ M),⁹ or peptide-based FP assays (100 nM). 11 We investigated the dependency of the binding affinity on factors including pH (range 4–10), DMSO concentration, and incubation time (Fig. S6–S9, ESI†). The assay is robust with Z' values ≥ 0.80 (Fig. S9, ESI†). Once PHD2 and probe 12 had attained equilibrium (30 min), the complex was stable over 24 h (Fig. S7, ESI†). The binding affinity was reasonably stable in the presence of up to 16% (v/v) DMSO (Fig. S8, ESI†). The new assay complements

turnover based methods and is more efficient, simpler, and lower cost than previous methods (Fig. 5B).

We then validated the assay by evaluating the affinities of the clinical inhibitors FG-4592 (Roxadustat), AKB-6548 (Vadadustat), and 1 for PHD2. The IC_{50} values of FG-4592, AKB-6548, and 1 for inhibition of the PHD2/probe 12 interaction were 120.8 \pm 3.8 nM, 215.1 \pm 2.1 nM, and 21.5 \pm 2.3 nM, respectively. These values are in accord with the IC_{50} values determined by the HIF-1a peptide-based FP (FG-4592, 591.4 nM; AKB-6548, 608.7 nM; 1, 77.7 nM)¹¹ and SPE MS substrate turnover (FG-4592, 2.587 μ M; 1, 0.276 μ M) assays (Fig. 6). Though there are variations in the absolute IC_{50} values derived from the probe 12-based FP assay, the HIF-1 α peptidebased FP assay, or SPE MS substrate turnover assays, due to the different experimental conditions, 21 the trends of the inhibitory activity towards PHD2 are consistent for all of these.

Thus, we then used probe 12 to evaluate the binding affinities of naturally occurring 2OG analogues and PHD2 (Fig. 5B). The assay reveals that N-oxalylglycine (NOG), a plant metabolite and 2OG isostere,¹⁸ inhibits the PHD2/probe 12 interaction with an IC₅₀ of 2.52 \pm 0.6 µM. This value is similar to the IC₅₀ (3 \pm 1 μ M) determined by an NMR binding assay, $^{\mathrm{s}}$ though lower than that obtained by MS based turnover assay $(IC_{50}$ of 18.5 µM).⁸ It is proposed that in cells PHD activity may be regulated by TCA cycle intermediates and related compounds, $19,20,22$ thus we used our assay to investigate affinities of such compounds for PHD2 (Fig. 6 and Table S1, ESI†). Consistent with prior work using NMR-based and MSbased assays, ^{9,13} fumarate, and succinate bind weakly to PHD2 with IC₅₀ values of 77.94 \pm 1.8 µM and 64.0 \pm 1.4 µM, respectively; consistent with prior studies 2-hydroxylgutarate also bound weakly (Fig. 6). Chem Comm

Communication

Interestingly, the assay revealed that dicarboxylic aminoacids including aspartate (IC₅₀: 162.2 \pm 4.2 µM) and glutamate $(IC_{50}: 65.07 \pm 4.2 \mu M)$ bind PHD2, albeit weakly (Fig. 6). The concentrations of some amino acids, are high in the cytosol;

Fig. 5 (A) Optimisation of the concentration of probe 12 and PHD2 and inhibition by FG-4592 and NOG; (B) schematic representation of FP assays used to investigate the binding of probe 12 and PHD2 and the displacement of probe 12 by PHD2 inhibitors.

Fig. 6 IC₅₀ values of these representative compounds (^aHIF-1 α peptidebased FP assay; ^bNMR assay; ^cSPE MS assay).

Communication ChemComm

this is particularly true for glutamate where concentrations of \geq 10 mM are reported in some cells.²³ N-Acylated derivatives of Glu, are reported as PHD inhibitors, 24 but to our knowledge there are no reports of Glu itself binding to the PHDs. There is thus a possibility of crosstalk between amino acid metabolism and the HIF pathway/2OG oxygenase catalysis in cells. If indeed PHD activity in cells is mediated by glutamate, there are implications for intracellular inhibition of the PHDs by other small-molecules, both by drugs and TCA metabolites and the related oncometabolite 2-hydroxyglutarate.^{6,25}

The overall results demonstrate that the small-molecule probe 12-based FP assay is an efficient and cost-effective method for analysis of both strong and weakly binding fragment-like PHD2 inhibitors. The assay complements reported fluorescence based assays for $2OG$ oxygenases, 26 requires small amounts of reagents compared to turnover-assays $8-10$ and is suited to HTS application. Probe 12 binds well to FIH (Table 1); thus there is scope for its further development. Probe 12 did not manifest cytotoxicity in HEK293, Hep3B, and L02 cells (Fig. S10, ESI†), so may also be useful in exploring PHD biology and inhibition in cells.

In comparison to our reported affinity-based assay for PHD2 using a FITC-labelled HIF-1 α (556–574) peptide as the FP probe, 11 the probe 12-based FP assay has the advantage that it works with Fe(π); the HIF-1 α peptide-based FP assay employs $Mn(\pi)$ instead of Fe(π), because Fe(π)-complexed PHD2 hydroxylates the peptide probe causing it to lose affinity (Fig. S5B, $EST[†]$).¹¹ Additionally, given that the HIF peptide probe does not occupy key regions of the active site, e.g. the 2OG binding pocket, the HIF-1 α peptide-based FP assay fails to identify certain types of inhibitor, including fragment-like compounds, e.g. NOG (Fig. S11, ESI†), a problem overcome by the probe 12-based FP assay. The combined use of the two assays can provide information on whether inhibitors compete with HIF- α and/or 2OG. The assay was validated by work with clinically used inhibitors and naturally occurring fragment-like PHD inhibitors, such as TCA cycle intermediates and amino acids, the results of which are being followed up. We envision future applications of probe 12 in studying the biological roles of the PHDs and crosstalk between the HIF pathway, the TCA cycle, and amino acid metabolism. Communication Communication Communication Communication Communication 2020. Downloaded on 24. A pharmaceutic is are controlled under the properties are controlled under the same of the communication of the same of the sam

This work was supported by grants from the National Natural Science Foundation of China (grant code: 81973173), Jiangsu Province Funds for Excellent Young Scientists (grant code: BK20170088), the Six Talent Peaks Project (grand code: YY-023) of Jiangsu Province, the Wellcome Trust (grant code: 106244/Z/ 14/Z) and Cancer Research UK (grant code: C9047/A24759).

Conflicts of interest

The authors declare no conflict of interest.

Notes and references

1 R. Chowdhury, A. Hardy and C. J. Schofield, Chem. Soc. Rev., 2008, 37, 1308–1319.

- 2 M. H. Rabinowitz, J. Med. Chem., 2013, 56, 9369–9402.
- 3 C. J. Schofield and P. J. Ratcliffe, Nat. Rev. Mol. Cell Biol., 2004, 5, 343–354.
- 4 A. A. Joharapurkar, V. B. Pandya, V. J. Patel, R. C. Desai and M. R. Jain, J. Med. Chem., 2018, 61, 6964–6982.
- 5 Z. Li, Q. You and X. Zhang, J. Med. Chem., 2019, 62, 5725–5749.
- 6 H. Tarhonskaya, A. M. Rydzik, I. K. H. Leung, N. D. Loik, M. C. Chan, A. Kawamura, J. S. O. McCullagh, T. D. W. Claridge, E. Flashman and C. J. Schofield, Nat. Commun., 2014, 5, 4423.
- 7 L. A. McNeill, L. Bethge, K. S. Hewitson and C. J. Schofield, Anal. Biochem., 2005, 336, 125–131.
- 8 C. J. Stubbs, C. Loenarz, J. Mecinovic, K. K. Yeoh, N. Hindley, B. M. Lienard, F. Sobott, C. J. Schofield and E. Flashman, J. Med. Chem., 2009, 52, 2799–2805.
- 9 I. K. H. Leung, M. Demetriades, A. P. Hardy, C. Lejeune, T. J. Smart, A. Szollossi, A. Kawamura, C. J. Schofield and T. D. W. Claridge, J. Med. Chem., 2013, 56, 547–555.
- 10 R. Chowdhury, J. I. Candela-Lena, M. C. Chan, D. J. Greenald, K. K. Yeoh, Y. M. Tian, M. A. McDonough, A. Tumber, N. R. Rose, A. Conejo-Garcia, M. Demetriades, S. Mathavan, A. Kawamura, M. K. Lee, F. van Eeden, C. W. Pugh, P. J. Ratcliffe and C. J. Schofield, ACS Chem. Biol., 2013, 8, 1488–1496.
- 11 Y. Lei, T. Hu, X. Wu, Y. Wu, Q. Bao, L. Zhang, H. Xia, H. Sun, Q. You and X. Zhang, ACS Med. Chem. Lett., 2015, 6, 1236–1240.
- 12 C. Domene, C. Jorgensen and C. J. Schofield, J. Am. Chem. Soc., 2020, 142, 2253–2263.
- 13 K. S. Hewitson, B. M. Lienard, M. A. McDonough, I. J. Clifton, D. Butler, A. S. Soares, N. J. Oldham, L. A. McNeill and C. J. Schofield, J. Biol. Chem., 2007, 282, 3293–3301.
- 14 Y. Wu, Z. Jiang, Z. Li, J. Gu, Q. You and X. Zhang, J. Med. Chem., 2018, 61, 5332–5349.
- 15 M. A. McDonough, V. Li, E. Flashman, R. Chowdhury, C. Mohr, B. M. Lienard, J. Zondlo, N. J. Oldham, I. J. Clifton, J. Lewis, L. A. McNeill, R. J. Kurzeja, K. S. Hewitson, E. Yang, S. Jordan, R. S. Syed and C. J. Schofield, Proc. Natl. Acad. Sci. U. S. A., 2006, 103, 9814–9819.
- 16 J. P. Holt-Martyn, R. Chowdhury, A. Tumber, T.-L. Yeh, M. I. Abboud, K. Lippl, C. T. Lohans, G. W. Langley, W. Figg Jr., M. A. McDonough, C. W. Pugh, P. J. Ratcliffe and C. J. Schofield, ChemMedChem, 2020, 15, 270–273.
- 17 S. Dhillon, Drugs, 2019, 79, 563–572.
- 18 K. Al-Qahtani, B. Jabeen, R. Sekirnik, N. Riaz, T. D. W. Claridge, C. J. Schofield and J. S. O. McCullagh, Phytochemistry, 2015, 117, 456–461.
- 19 M. A. Selak, S. M. Armour, E. D. MacKenzie, H. Boulahbel, D. G. Watson, K. D. Mansfield, Y. Pan, M. C. Simon, C. B. Thompson and E. Gottlieb, Cancer Cell, 2005, 7, 77–85.
- 20 R. V. Duran, E. D. MacKenzie, H. Boulahbel, C. Frezza, L. Heiserich, S. Tardito, O. Bussolati, S. Rocha, M. N. Hall and E. Gottlieb, Oncogene, 2013, 32, 4549–4556.
- 21 T.-L. Yeh, T. M. Leissing, M. I. Abboud, C. C. Thinnes, O. Atasoylu, J. P. Holt-Martyn, D. Zhang, A. Tumber, K. Lippl, C. T. Lohans, I. K. H. Leung, H. Morcrette, I. J. Clifton, T. D. W. Claridge, A. Kawamura, E. Flashman, X. Lu, P. J. Ratcliffe, R. Chowdhury, C. W. Pugh and C. J. Schofield, Chem. Sci., 2017, 8, 7651–7668.
- 22 M. I. Abboud, T. E. McAllister, I. K. H. Leung, R. Chowdhury, C. Jorgensen, C. Domene, J. Mecinović, K. Lippl, R. L. Hancock, R. J. Hopkinson, A. Kawamura, T. D. W. Claridge and C. J. Schofield, Chem. Commun., 2018, 54, 3130–3133.
- 23 A. J. L. Cooper and T. M. Jeitner, Biomolecules, 2016, 6, 16.
- 24 D. R. Mole, I. Schlemminger, L. A. McNeill, K. S. Hewitson, C. W. Pugh, P. J. Ratcliffe and C. J. Schofield., Bioorg. Med. Chem. Lett., 2003, 13, 2677–2680.
- 25 P. Koivunen, S. Lee, C. G. Duncan, G. Lopez, G. Lu, S. Ramkissoon, J. A. Losman, P. Joensuu, U. Bergmann, S. Gross, J. Travins, S. Weiss, R. Looper, K. L. Ligon, R. G. W. Verhaak, H. Yan and W. G. Kaelin Jr, Nature, 2012, 483, 484–488.
- 26 X. Luo, Y. Liu, S. Kubicek, J. Myllyharju, A. Tumber, S. Ng, K. H. Che, J. Podoll, T. D. Heightman, U. Oppermann, S. L. Schreiber and X. Wang, J. Am. Chem. Soc., 2011, 133, 9451–9456.