## ChemComm



**View Article Online** 

## CORRECTION

Check for updates

Cite this: Chem. Commun., 2020, 56, 4130

## Correction: $O^2$ -(6-Oxocyclohex-1-en-1-yl)methyl diazen-1-ium-1,2-diolates: a new class of nitric oxide donors activatable by GSH/GST $\pi$ with both anti-proliferative and anti-metastatic activities against melanoma

Chengfeng Bai,†<sup>a</sup> Rongfang Xue,†<sup>a</sup> Jianbing Wu,<sup>a</sup> Tian Lv,<sup>a</sup> Xiaojun Luo,<sup>a</sup> Yun Huang,<sup>a</sup> Yan Gong,<sup>a</sup> Honghua Zhang,<sup>b</sup> Yihua Zhang\*<sup>a</sup> and Zhangjian Huang\*<sup>a</sup>

DOI: 10.1039/d0cc90145h

rsc.li/chemcomm

Correction for ' $O^2$ -(6-Oxocyclohex-1-en-1-yl)methyl diazen-1-ium-1,2-diolates: a new class of nitric oxide donors activatable by GSH/GST $\pi$  with both anti-proliferative and anti-metastatic activities against melanoma' by Chengfeng Bai *et al.*, *Chem. Commun.*, 2017, **53**, 5059–5062.

The authors regret that in the original article incorrect images were used for the "Control" panel in Fig. 2A and for the "COMC-6" panel in Fig. 2B. To completely correct the errors, the experiments presented in Fig. 2A and B were re-performed, and the original images in Fig. 2A and B should be replaced with the new images.


The correct Fig. 2 is presented here. The corrected figure does not affect any results presented in the originally-published version, nor the corresponding text description or the conclusion of the paper.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup> State Key Laboratory of Natural Medicines, Jiangsu Key Laboratory of Drug Discovery for Metabolic Diseases, Jiangsu Key Laboratory of Drug Screening, China. E-mail: zyhtgd@163.com, zhangjianhuang@cpu.edu.cn

<sup>b</sup> Foreign Languages Department, China Pharmaceutical University, Nanjing 210009, P. R. China

† These authors contributed equally.



**Fig. 2** (A) Effects of compounds on B16–BL6 cell migration. Data were expressed as the mean  $\pm$  SD of each group of cells. \*\*\**P* < 0.001. (B) Effects of compounds on B16–BL6 cell invasion. Data were expressed as the mean  $\pm$  SD of each group of cells. \*\*\**P* < 0.001. (C) Effects of compounds on B16–BL6 cell lateral migration. Data are expressed as the mean  $\pm$  SD of each group of cells. \*\*\**P* < 0.001. (C) Effects of compounds on B16–BL6 cell lateral migration. Data are expressed as the mean  $\pm$  SD of each group of cells. \*\*\**P* < 0.001. Scale bars: 200  $\mu$ m.