Cite this: *Phys. Chem. Chem. Phys.*,
2020, 22, 4840

DOI: 10.1039/d0cp90040k

rsc.li/pccp

Correction: Conformation of bis-nitroxide polarizing agents by multi-frequency EPR spectroscopy

Janne Soetbeer,^a Peter Gast,^b Joseph J. Walish,^c Yanchuan Zhao,^c
Christy George,^a Chen Yang,^a Timothy M. Swager,^c Robert G. Griffin^a and
Guinevere Mathies^{†*}

Correction for 'Conformation of bis-nitroxide polarizing agents by multi-frequency EPR spectroscopy' by Janne Soetbeer *et al.*, *Phys. Chem. Chem. Phys.*, 2018, **20**, 25506–25517.

We correct an error in eqn (4) in our published article. The correct expression for the dipolar coupling between two electron spins in the high-field limit is

$$\mathbf{D}^{\text{PAS,D}} = -\frac{\mu_0}{4\pi\hbar} \frac{g_1 g_2 \mu_B^2}{r_{12}^3} \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad (4)$$

Thus, a minus-sign was lacking previously. In our calculations the correct expression was used (the minus-sign was present) and, hence, our results and conclusions are unaltered.

We furthermore note that in eqn (1) the exchange interaction, J , was erroneously typeset in bold. Our assumption is that the exchange interaction is isotropic and can be represented by a scalar.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Francis Bitter Magnet Laboratory and Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^b Department of Physics, Huygens-Kamerlingh Onnes Laboratory, Leiden University, PO Box 9504, 2300 RA Leiden, The Netherlands

^c Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

[†] Current address: Laboratory of Physical Chemistry, Department of Chemistry and Applied Biosciences, ETH Zurich, Vladimir-Prelog-Weg 2, 8093 Zürich, Switzerland.

^{*} Current address: Department of Chemistry, University of Konstanz, 78464 Konstanz, Germany. E-mail: guinevere.mathies@uni-konstanz.de