

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)

Cite this: *Dalton Trans.*, 2020, **49**, 1334

Correction: Azamacrocycles and tertiary amines can be used to form size tuneable hollow structures or monodisperse oxide nanoparticles depending on the 'M' source

Graham E. Tilburey,^a Toby J. Blundell,^b S. V. Patwardhan,^a Stephen P. Argent^c and Carole C. Perry^{*a}

DOI: 10.1039/c9dt90289a
rsc.li/dalton

Correction for 'Azamacrocycles and tertiary amines can be used to form size tuneable hollow structures or monodisperse oxide nanoparticles depending on the 'M' source' by Graham E. Tilburey, *et al.*, *Dalton Trans.*, 2019, **48**, 15470–15479.

The authors would like to correct the author list, as S. V. Patwardhan is not included on the author list in the published article. The correct author list is shown above.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aBiomolecular and Materials Interface Research Group, Interdisciplinary Biomedical Research Centre, Nottingham Trent University, Clifton Lane, Nottingham NG11 8NS, UK
E-mail: carole.perry@ntu.ac.uk

^bDepartment of Chemistry and Forensics, Nottingham Trent University, Clifton Lane, Nottingham NG11 8NS, UK

^cSchool of Chemistry, University of Nottingham, University Park, Nottingham NG7 2RD, UK

