

PAPER

[View Article Online](#)
[View Journal](#) | [View Issue](#)
Cite this: *Food Funct.*, 2020, **11**, 617

Effect of semolina pudding prepared from *starch branching enzyme IIa* and *b* mutant wheat on glycaemic response *in vitro* and *in vivo*: a randomised controlled pilot study†

Marina Corrado,^a Anna Cherta-Murillo,^b Edward S. Chambers,^b Abigail J. Wood,^c Amy Plummer,^c Alison Lovegrove,^c Cathrina H. Edwards,^a Gary S. Frost^b and Brittany A. Hazard^{*,a,d}

Refined starchy foods are usually rapidly digested, leading to poor glycaemic control, but not all starchy foods are the same. Complex carbohydrates like resistant starch (RS) have been shown to reduce the metabolic risk factors for chronic diseases such as hyperglycaemia and overweight. The aim of the project was to develop a semolina-based food made from a *starch branching enzyme II (sbella/b-AB)* durum wheat mutant with a high RS content and to measure its glycaemic index using a double-blind randomised pilot study. We report here the amylose, RS and non-starch polysaccharide concentration of raw *sbella/b-AB* and wild-type control (WT) semolina. We measured RS after cooking to identify a model food for *in vivo* testing. Retrograded *sbella/b-AB* semolina showed a higher RS concentration than the WT control (RS = 4.87 ± 0.6 g per 100 g, 0.77 ± 0.34 g per 100 g starch DWB, respectively), so pudding was selected as the test food. Ten healthy participants consumed ~50 g of total starch from WT and *sbella/b-AB* pudding and a standard glucose drink. Capillary blood glucose concentrations were measured in the fasting and postprandial state (2 h): incremental area-under-the-curve (iAUC) and GI were calculated. We found no evidence of difference in GI between *sbella/b-AB* pudding and the WT control, but the starch digestibility was significantly lower in *sbella/b-AB* pudding compared to the WT control *in vitro* (C_{90} = 33.29% and 47.38%, respectively). Based on these results, novel *sbella/b-AB* wheat foods will be used in future *in vivo* studies to test the effect of different RS concentrations and different food matrices on glycaemia.

Received 21st October 2019,
Accepted 9th December 2019

DOI: 10.1039/c9fo02460c

rs.c.li/food-function

1. Introduction

Wheat represents one of the main staple crops globally. It is well-adapted and consumed world-wide because it is highly nutritious, providing up to 34% of the daily calorie intake in western diets, together with other cereals.¹ Like many staple foods, wheat is usually consumed after processing into a range of refined food-products; these foods are often perceived as unhealthy because they are rapidly digested and low in fibre.^{2,3} Starchy foods made from refined flour typically give rise to a

sharp increase in the postprandial blood glucose concentration and are therefore described as having a high glycaemic index.⁴ Starch that is digested more slowly is expected to result in a more gradual postprandial glycaemic response. In some cases, starch can resist or escape digestion in the small intestine ('resistant starch'), moving into the large intestine where it may be fermented by the resident microbiota into short-chain fatty acids (SCFAs). This is believed to have positive effects on glucose metabolism.^{5,6}

Starch is a ubiquitous storage polysaccharide in wheat endosperm and the main source of digestible carbohydrates; however, its digestibility depends on its native structure in the grain and physical form in foods. In the native state, starch exists as dense granules where the α -glucan chains are not readily available for amylase digestion.^{7,8} Most starch-based foods are cooked to improve digestibility as well as texture; heating starch in excess water results in water absorption and swelling of the starch granules leading to loss of native resistant starch (RS), a process known as gelatinisation. As the starch cools, starch polysaccharides tend to re-associate in a

^aFood Innovation and Health, Quadram Institute Bioscience, Norwich Research Park, UK. E-mail: brittany.hazard@quadram.ac.uk

^bSection for Nutrition Research, Department of Metabolism, Digestion and Reproduction, Faculty of Medicine, Imperial College London, London, UK

^cRothamsted Research, Harpenden, UK

^dDesigning Future Wheat and Molecules from Nature, John Innes Centre, Norwich Research Park, UK

†Electronic supplementary information (ESI) available. See DOI: 10.1039/c9fo02460c



more ordered and crystalline state (retrogradation) becoming more resistant to digestion.⁹ The physicochemical modifications that starch undergoes during processing, together with the native starch structure, can alter the resistance of starch to digestion in the small intestine.^{10,11} In high amylose starches, molecular interactions between amylose–amylose, amylose–amylopectin and amylopectin–amylopectin can lead to retrogradation. During gelatinisation, amylose leaches out from granules as a random coil which tends to form either single-helical complexes (amylose–lipid complexes) or double helices by self-association. The rate of self-association depends on chain-lengths, concentration and cooling rate; high amylose starches are more likely to form double helical structures which are thought to be resistant to α -amylase digestion.¹² Starch digestibility is also strongly influenced by the surrounding food matrix properties, which can hinder starch gelatinisation during processing conditions and/or limit starch exposure to digestive enzymes and fluids during gastrointestinal transit.^{13–15}

Another factor that may influence the digestibility of foods made from wheat semolina is the presence of non-starch polysaccharides (NSPs).¹⁶ NSPs derived from the grain cell walls completely resist enzymatic digestion in the small intestine.¹⁷ They are the main source of dietary fibre from durum wheat semolina (1.4 to 1.8% dry weight), with the major fraction being arabinoxylan AX (~70%) and (1→3,1→4)- β -D-glucan (mixed-linkage β -glucan or MLG) (~20%).¹⁸ AX has a backbone of β -D-xylopyranosyl residues linked by (1→4) glycosidic linkages with some residues substituted with α -L-arabinofuranosyl at positions 2 and/or 3.^{19,20} At high concentrations, AXs, particularly water-soluble AXs, can contribute to an increased meal viscosity and lower starch gelatinisation and hydrolysis, having implications for postprandial glycaemia.^{21,22}

In an attempt to increase the fibre content of refined wheat foods and lower their glycaemic index (GI), *sbeIIa/b*-AB durum wheat was developed by combining mutations in *sbeII* genes, encoding Starch Branching Enzyme II (SBEII) which catalyzes the addition of branch points during amylopectin biosynthesis.²³ The partial loss of function of SBEIIa and SBEIIb enzymes during starch formation leads to an increase in the amount of amylose and the proportion of starch that is resistant to digestion in raw semolina, producing pasta of acceptable quality with improved firmness after cooking.²⁴ Animal studies have shown a positive fermentation response to *sbeIIa/b*-AB wheat provided as pellets with an increase in the production of SCFA and lower colonic pH in rats fed with *sbeIIa/b*-AB flour compared to WT flour for 4 weeks.^{25,26}

In pig studies, other forms of RS were found to lower metabolizable energy intake, prolong satiety and decrease the kinetics of glucose absorption while altering gut hormone secretion.^{27,28} In an acute human study, decreased energy intake was observed after consumption of muffins containing high RS compared to a low RS breakfast.²⁹ Harrold J. *et al.* found that consumption of a new composite satiety ingredient made of whole-grain and high-amylose corn flour (30 g of RS on a dry weight basis) for breakfast led to increased satiety and

lower energy intake at dinner, compared to the maltodextrin control.³⁰

Therefore, increasing resistant starch in wheat seems to be a promising strategy for improving the dietary fibre content of wheat-based foods that could lead to steady glucose release in the gut and a gradual increase in postprandial glucose concentrations. Even though *sbeIIa/b*-AB wheat is known to have higher amylose and resistant starch concentration than its WT control in the raw state, very little is known about the physicochemical properties of starch and non-starch polysaccharides in *sbeIIa/b*-AB semolina after cooking and its impact on blood glucose response. In this study, we measured the RS concentration of two model foods, semolina porridge and semolina pudding. Semolina pudding was selected as a model food to measure starch digestibility (C_{90}) *in vitro* and to explore pudding GI *in vivo*, in a double-blind randomised controlled pilot study.

2. Materials & methods

2.1. Materials

Durum wheat [*T. turgidum* L. subsp. *Durum* (Desf.) Husn.] with high RS and amylose concentration was developed previously at the University of California Davis using the cultivar Kronos by combining mutations in *starch branching enzyme II* genes (*sbeIIa/b*-AB).³¹ For this study, the *sbeIIa/b*-AB mutant and WT control plants were grown at the University of California Experimental Field Station in Davis, CA, during the 2014–2015 growing season and harvested in bulk in June 2015.

2.2. Milling and particle size

Prior to milling, grains were conditioned using a two-stage water addition and de-branned using a Satake laboratory de-branner. Bulk durum wheat grains were milled to obtain semolina using a Bühler laboratory mill at the Allied Milling and Baking Technical Centre (Vanwall Road, Maidenhead, Berkshire, SL6 4UF). The particle size distributions of *sbeIIa/b*-AB and WT semolina suspended in water were determined by laser diffraction analysis on a Beckman Coulter LS 13 320.

2.3. Starch extraction from WT and *sbeIIa/b*-AB mutant grains

Approximately 50 wheat grains were left overnight at +4 °C in a Petri dish on moistened filter paper. They were then ground using a pestle and a mortar in approximately 5 mL of deionised water, after removing the husk and embryo manually. The extract was then filtered through a layer of Miracloth (475855, Calbiochem®) by washing with 30 mL of deionised water. The solution was centrifuged, at 2000g for 5 min and the recovered pellet was washed with 30 mL of 2% Sodium Dodecyl Sulfate (SDS). The pellet was washed and centrifuged twice in 5 mL of 2% SDS. The resulting starch pellet was washed a further three times with 5 mL of acetone and left to dry overnight in a fume hood to ensure complete evaporation of the solvent.³²



2.4. Food models: semolina processing for *in vitro* analyses

Mutant *sbeIIa/b*-AB semolina and WT control semolina were gelatinised by adding 1 mL of deionised water to ~100 mg of semolina followed by boiling for 10 min at 99 °C while stirring to ensure homogeneous gelatinisation of the starch (porridge model). The gelatinised samples were cooled on ice overnight with continuous stirring during which some retrogradation occurred (pudding model). The RS concentration of model food products (porridge and pudding) was analysed to identify one food product to be tested *in vivo* as described below.

2.5. Formulation and preparation of semolina pudding for *in vivo* analysis

Test meals consisted of semolina puddings made from *sbeIIa/b*-AB or WT control wheat semolina. Both pudding recipes were designed to provide approximately 50 g of total starch with a constant solid to liquid ratio consisting of 17.02% semolina, 82.75% water and 0.23% salt (w/w fresh weight basis). Dry ingredients (*sbeIIa/b*-AB or WT semolina and salt) were packed into coded containers at the Quadram Institute Bioscience (QIB) and shipped to the NIHR Imperial Clinical Research Facility, London (UK), where the individual portions were cooked prior to serving such that both participants and researchers preparing the pudding and assessing the outcomes were blinded. Wheat semolina puddings were prepared by simmering at 80 °C for 3 min while stirring to obtain homogeneous starch gelatinisation and cooled at room temperature for 10 min before storing at 4 °C overnight to allow the starch to retrograde. After preliminary analysis of the results, the test meals were unblinded to reveal the treatment and control for completing the statistical analysis.

Each test meal was prepared individually and served with 250 to 400 mL of water. The volume of water served with the test meal was adjusted to compensate for the water loss during the cooking of the semolina pudding and ensured a constant liquid–solid ratio across all meals.

2.6. Starch analysis: total starch, amylose, resistant starch and moisture content

Starch analyses are reported on a dry weight basis. The moisture of unprocessed semolina was determined using a modified version of the AACC (44-15A) air oven method, a one stage procedure, in which the amount of sample used was reduced to 100 mg due to the limited quantity of the available material. The total starch (TS) content of unprocessed semolina was determined using Megazyme's Total Starch Assay Kit (KOH format, AOAC 996.11). The resistant starch (RS) content was measured using Megazyme's Assay Kit (AOAC 2002.02) following the manufacturer's instructions.

2.7. Amylose determination

0.05% (w/v) suspension of starch (isolation as described above) in 6 mM iodine solution (0.229 g of re-sublimed iodine in 270 mL of DMSO and 30 mL of deionised water) was left to dissolve on a vertical multifunction rotator (PTR 35, Grant BIO)

providing end over end mixing to allow starch to dissolve overnight. The samples were then diluted 1:10 with deionised water and left for 30 min to allow the iodine–amylose complex to form a stable colour prior to reading the absorbance at 600 nm (Benchmark™ Plus Microplate Reader, Bio-Rad) against the reagent blank. The apparent amylose content was calculated using a standard curve of amylose standards (1–5 mg potato amylose type III, Sigma-Aldrich).^{33–35}

2.8. Non-starch polysaccharide analysis

2.8.1. Determination of total and water-extractable monosaccharides. Briefly, for 'total' monosaccharide analysis, 5 mg of semolina sample from WT and *sbeIIa/b*-AB mutant wheat lines were suspended in 10 mL of deionised water (0.5 mg mL⁻¹) and homogenised in a glass/Teflon homogeniser. Triplicate aliquots of 200 µL (equivalent to 100 µg of the original sample) were taken and freeze-dried overnight. For WE-monosaccharide analysis 5 mg of the semolina sample from WT and *sbeIIa/b*-AB wheat were resuspended in 1 mL of deionised water and incubated on a Denley spiromix for 30 minutes at 25 °C. The suspension was centrifuged at 2500g for 10 minutes and the supernatant was collected, aliquoted and freeze-dried overnight. Both TOT and WE samples were hydrolysed in 2 M trifluoroacetic acid (TFA) as described by Bromley *et al.*³⁶ Hydrolysates were diluted to 0.25 µg µL⁻¹ in deionised water. If samples were not to be analysed immediately, they were stored at –20 °C. Three technical replicate samples were aliquoted and further diluted in 10 µM 2-deoxygalactose solution (internal standard) to a concentration of 0.05 µg µL⁻¹.

Neutral monosaccharides were separated by HPAEC-PAD (using a Dionex ICS-5000+ equipped with an eluent generator with EGC 500 KOH cartridge; Thermo Scientific, USA) with a 23.5 min run on a CarboPac PA-20 column (guard column: 3 × 30 mm, analytical column: 3 × 150 mm; Thermo Scientific, USA) at 30 °C. Twenty-five µL of the diluted sample was injected into the column, using the following conditions: isocratic 4 mM KOH, 14.5 min; 100 mM KOH, 15–18 min; 4 mM KOH, 18.5–23.5 min; the flow rate was 0.5 mL min⁻¹. Authentic monosaccharide standards were subjected to the same acid hydrolysis as for the experimental samples to allow for any destruction of sugars during hydrolysis prior to the generation of calibration curves and were used for peak identification. All experimental samples were randomised prior to HPAEC analysis. Calibration standards were run at the beginning, middle and end of an analytical run cycle. All data were processed with Chromeleon analytical software (version 7.2SR4; Thermo Scientific, US) to calculate the amount of the analyte in an unknown sample using external standards/calibration curves. Total and WE-AX were calculated as the sum of xylose and arabinose minus an adjustment for arabinogalactan peptide^{37,38} and multiplied by 0.89 to convert monosaccharide values, for xylose and arabinose, to the polysaccharide arabinoxylan.

2.8.2. Enzyme fingerprinting. Enzymatic fingerprinting was used to measure arabinoxylan oligosaccharides (AXOS) and mixed-linked β-glucan oligosaccharides (GOS) released fol-



lowing enzymatic digestion of semolina and freeze-dried pudding samples from the WT and *sbeIIa/b*-AB mutant. The method used was as described by Ordaz-Ortiz *et al.* (2004);³⁹ except that recombinant enzymes were used for digestion. The method used for the identification of oligosaccharides was described by Ordaz-Ortiz & Saulnier (2005).⁴⁰ Semolina from WT and *sbeIIa/b*-AB mutant wheat lines were digested with *endo* 1,4 β -xylanase (E.C.3.2.1.8), a xylanase of the GH11 group from *Neocallimastix patriciarum* (PRO-E0062, Promix Limited, UK,) and *endo* 1,3(4) glucanase ('lichenase') (E.C.3.2.1.73) from *Clostridium thermocellum* NCIB 10682 (PRO-E0017, Promix Limited, UK) to digest AX and MLG, respectively. Following digestion, the samples were filtered, diluted in 10 μ M melibiose (internal standard), and analysed using a CarboPac PA1 analytical column (guard column: 2 \times 50 mm \times analytical column: 2 \times 250 mm) on a Dionex ICS-3000 (Thermo Scientific). The peak areas of the oligosaccharides released by enzyme digestion were expressed as percentages of the total peak areas of all AXOS. Two major GOS, G3 and G4, were released by enzymatic digestion. An estimate of 'total' MLG was therefore calculated as the sum of G3 + G4 peak areas, and the G3 to G4 ratio was calculated.

2.9. *In vitro* starch hydrolysis of pudding samples

To determine the effect of the combined mutations in *sbeIIa* and *sbeIIb* genes on the intrinsic susceptibility of starch to digestion by α -amylase, starch amylolysis was determined using an established *in vitro* digestion protocol.⁴¹ Puddings were prepared as described in section 2.5 (Formulation and preparation of semolina pudding). The samples were digested with porcine pancreatic α -amylase and the digestion products were quantified using PAHBAH assay for reducing sugar determination.⁴¹ Blank values taken for each assay prior to enzyme addition (Y_0) represent the baseline sugar value and were used to account for any endogenous sugar. Starch digestibility is expressed as C_{90} representing the percentage of *sbeIIa/b*-AB starch digested at 90 minutes compared to the WT control.⁴² Logarithm of slope (LOS) analysis was used to determine starch amylolysis in the first order equation

$$Ct = C_{\infty} (1 - e^{-kt}) \quad (1)$$

in the logarithmic form

$$\ln\left(\frac{dC}{dt}\right) = -kt + \ln(C_{\infty}k), \quad (2)$$

which gives a linear plot describing the relationship between LOS digestibility constants (k and C_{∞}), and time of amylolysis (t).⁴³ Here, Ct represents the concentration of the digestion product at time (t), C_{∞} is the product concentration at the end of the reaction time and k is the digestibility rate constant. C_{∞} can also be expressed as the percentage of starch digested over the total starch content of the sample. When endogenous sugar due to processing is present at the start of the reaction, the baseline sugar value (Y_0) is added to C_{∞} to estimate the percentage of starch digested in time (t).⁴⁴

2.10. *In vivo* glycaemic index pilot study

2.10.1. Subjects. Twelve healthy adults were recruited from a healthy volunteer database and deemed as healthy through a screening visit with a health questionnaire. Exclusion criteria were smoking, substance abuse, significant weight change, prediabetes or diabetes, familial history of diabetes, no food allergy or intolerance and not having participated in a research trial in the last 6 months. A total of 2 participants withdrew from the study for various reasons (unable to complete the meal within 15 min and unable to self-finger-prick). The characteristics of the 10 participants completing the study (8 females and 2 males, age 33 ± 2 years, BMI 23.53 ± 0.57 kg m⁻², mean \pm SE) can be found in Table 5. The study flow diagram based on the Consolidated Standards of Reporting Trials (CONSORT) can be found in ESI Fig. S1.† Screening and study visits took place at the National Institute for Health Research (NIHR)/Wellcome Trust Imperial Clinical Research Facility, London. The study was approved by the West London Research Ethics Committee 2 (13/LO/0696). This study was conducted from September to November 2017 according to the Declaration of Helsinki. All participants provided written informed consent before enrolment in the study.

2.10.2. Study design. The pilot study was conducted using a two-arm double-blind randomised cross-over design, based on the standardised GI screening methodology described by Brouns *et al.* 2005.⁴⁵ Twelve participants were randomised to four test meals: two glucose drinks (The Boots Company PLC, Nottingham, England), one WT semolina pudding and one *sbeIIa/b*-AB semolina pudding, given in four study visits separated by a wash-out period of 2–7 days.⁴⁵ Semolina pudding meals were accompanied by water to drink. Test foods and glucose drink were designed to provide 50 g of available carbohydrates. The nutritional composition of the puddings is reported in ESI Table S1.†

Participants arrived at the research facility at 08:30 am following an overnight fast. Two fasting blood samples (–10 and –5 min) were collected followed by administration of the test meal (*i.e.* *sbeIIa/b*-AB or WT semolina pudding and water drink) or glucose drink. Participants were asked to consume the test meal within 15 minutes at an even pace and capillary blood samples were collected at 15, 30, 45, 60, 90, and 120 min after consuming a test meal or glucose drink. Blood samples were self-obtained using disposable finger-pricks (Unistix needles) and glucose readings were measured using instant glucose analysers (Hemocue, Glucose 201+, Prospect Diagnostics Ltd, Dronfield, Derbyshire). Each study visit was of the same nature.

2.10.3. Statistical analysis. Statistical analyses, graphs and incremental area under the curve (iAUC) were obtained with R.⁴⁶ Datasets were curated using the reshape2 package and plots were obtained using the ggplot2 package.^{47,48} For each participant, glycaemic index (GI) was calculated from the blood glucose response values obtained from the iAUC calculated between 0 and 120 min as per Brouns *et al.* 2005.⁴⁹ GI was determined as the ratio of the test food (*sbeIIa/b*-AB or WT) and the reference glucose drink. The primary aim of the



study was to determine whether the GI of *sbeIIa/b-AB* pudding was different from the WT pudding control using a paired *t*-test. As a secondary study aim, we compared the iAUC 0–120 min of *sbeIIa/b-AB* and WT pudding using a paired *t*-test and the blood glucose concentrations after consuming *sbeIIa/b-AB* pudding and WT pudding by linear mixed-effects analysis (lme4 package).⁵⁰ The linear mixed-effects model included individual responses to the meal as the main effect, individual differences over time as random effects and changes over time due to the genotype as the fixed effect (with an interaction term). Visual inspection of the residual plots did not reveal any obvious deviations from homoscedasticity or normality and *p*-values were obtained by Satterthwaite approximation. RS, amylose and NSP concentration and composition of *sbeIIa/b-AB* were compared to those of the WT control using an independent two sample *t*-test. Further details of data analysis and statistical tests are described for each experiment in the Materials & methods sections. Data are presented as means \pm SEMs unless stated otherwise.

3. Results

3.1. Semolina non-starch polysaccharide characteristics

NSP content and composition were measured in the raw semolina from WT and *sbeIIa/b-AB* mutant to explore the contribution of other nutritional factors to digestibility changes and glycaemic response.

The total AX content of *sbeIIa/b-AB* was higher than that of the WT control in semolina (two sample *t*-test, $p = 0.0069$), (Table 1). However, there was no difference in the water-extractable AX (WE-AX) content between *sbeIIa/b-AB* semolina and WT (two sample *t*-test, $p = 0.32$).

There was no significant difference in total AXOS determined by enzymatic fingerprinting (sum of peak areas of all known AXOS peaks) in unprocessed *sbeIIa/b-AB* and WT semolina (two sample *t*-test, $p = 0.07$). Enzymatic fingerprinting revealed differences in the AXOS substitution pattern between *sbeIIa/b-AB* semolina and WT control (Table 2). Substituted AXOS were higher in *sbeIIa/b-AB* semolina compared to the WT control (two sample *t*-test, $p = 0.0016$, $p = 5.679 \times 10^{-11}$, respectively), but no significant differences in unsubstituted

Table 2 AXOS of *sbeIIa/b-AB* semolina compared to that of the wild-type control

Parameter (sum of peaks relative to the internal standard)	Semolina WT control	Semolina <i>sbeIIa/b-AB</i>
TOT-AXOS	10.02 \pm 0.74	12.64 \pm 0.97
AXOS unsubstituted	5.39 \pm 0.6	6.61 \pm 0.79
AXOS substituted	4.63 \pm 0.15	6.03 \pm 0.2
MLG (G3 + G4)	2.29 \pm 0.12	2.78 \pm 0.11

Means \pm SEMs of independently analysed semolina samples ($n = 4$) from bulked semolina.

AXOS were detected (two sample *t*-test, $p = 0.26$). To describe the structure of MLG and the distribution of β 1–3 and β 1–4 bonds, an estimate of total MLG was obtained by summing G3 and G4 GOS peak areas and the ratio of G3 to G4 GOS determined as the ratio of G3 to G4 GOS peak area.¹⁸ The mean of G3 + G4 of *sbeIIa/b-AB* semolina was higher than that of the WT control (two sample *t*-test, $p = 0.028$) and the ratio of G3 to G4 was 2.4 for *sbeIIa/b-AB* semolina vs. 2.2 for the WT control. NSP content and composition were also measured for freeze-dried pudding prepared alongside pudding portions served in the GI pilot study described below; these results can be found in the ESI (Tables S4 and S5†).

3.2. Starch characteristics before and after processing

The TS contents of bulked *sbeIIa/b-AB* and WT control semolina were 60.65 \pm 3.84 g per 100 g on a dry weight basis (DWB) and 67.90 \pm 1.44 g per 100 g, respectively (unpaired two-sample *t*-test, $p = 0.11$). The amylose content of starch isolated from *sbeIIa/b-AB* semolina was 33.05% \pm 0.69% (w/w dry matter basis) while the WT control was 23.11% \pm 1.5% (w/w dry weight basis) (unpaired two-sample *t*-test, $p = 0.004$).

Volume moment mean D [4, 3] particle size of *sbeIIa/b-AB* semolina was larger compared to that of the WT control (means \pm SEMs were 279 \pm 4.23 μ m and 258.42 \pm 5.54 μ m, respectively, $n = 3$ of bulked semolina, unpaired two sample *t*-test, $p = 0.004$). The particle size distribution is shown in ESI Fig. S2.†

The resistant starch content of raw *sbeIIa/b-AB* semolina (4.87 \pm 0.6 g per 100 g starch DWB) was higher than that of the WT control (0.77 \pm 0.34 g per 100 g starch DWB, two sample *t*-test, $p = 0.004$). Different amounts of resistant starch were detected after cooking ('gelatinisation') and cooling ('retrogradation') of *sbeIIa/b-AB* and WT semolina, as shown in Fig. 1. *sbeIIa/b-AB* semolina was characterised by a higher proportion of resistant starch than the WT control when gelatinised and retrograded.

3.3. In vitro starch digestibility of pudding

Mutant *sbeIIa/b-AB* pudding was less digested overall, and digestion occurred at a lower rate compared to the WT control. Amylolytic profiles of *sbeIIa/b-AB* and WT control are shown in Fig. 2a. *sbeIIa/b-AB* pudding starch was significantly less digested after 90 minutes, compared to the WT control

Table 1 Total (TOT) and water extractable (WE) arabinoxylans (AX) of *sbeIIa/b-AB* semolina compared to those of the wild-type controls (WT)

Parameter (mg g ⁻¹ semolina DWB)	Semolina WT control	Semolina <i>sbeIIa/b-AB</i>
WE-AX	1.17 \pm 0.11	1.51 \pm 0.51
TOT-AX	13.27 \pm 0.76	20.72 \pm 1.76

Means \pm sd of independently analysed semolina samples ($n = 3$) from bulked semolina. Total and WE-AX were calculated as the sum of xylose and arabinose minus an adjustment for arabinogalactan peptide as described in ref. 38, 51 and 52, and multiplied by 0.89 to convert monosaccharide values, for xylose and arabinose, to the polysaccharide arabinoxylan. Complete monosaccharide profiles are reported in ESI Table S3.†






Fig. 1 Proportion of resistant starch (expressed as the amount of RS per 100 g of semolina starch) of raw, cooked (gelatinised) and cooled (retrograded) semolina. Means \pm SEMs comparing *sbella/b-AB* to WT within each treatment. Dark blue and dark green datapoints in the figure represent means of *sbella/b-AB* and WT for each treatment. Light blue and light green datapoints represent $n = 3$ replicates of bulked semolina that were extracted and treated independently.




Fig. 2 (a) Starch digestibility of *sbella/b-AB* pudding (*sbella/b*) and wild-type pudding (WT), and starch amylolysis experimental data are shown by fitting a first-order equation based on the k and C_{∞} values obtained from the LOS analysis. Each experimental data point represents the mean value from the analysis of three pudding samples with error bars = \pm SEMs. (b) LOS plot obtained for WT pudding. (c) LOS plot for *sbella/b-AB* pudding. The linear phase is defined using eqn (2) from which k and C_{∞} values can be estimated.

(iAUC₉₀ mean difference = 11.923 $\mu\text{mol L}^{-1} \text{min}^{-1}$, [95CI 44 646.60, 71 756.06], paired t -test, $p = 0.0002839$). LOS analysis was used to determine the digestion rate at 90 minutes and both the starch amylolysis rate and extent were found to be lower for *sbella/b-AB* pudding compared to those for the WT control (C_{∞} and k). Linear plots with single rate constants were obtained from LOS analysis (Fig. 2b and c).

Table 3 Variable estimates from the digestibility curves and LOS analysis of *sbella/b-AB* pudding compared to those of WT control pudding

Pudding	C_{90} (%)	C_{∞} (%)	k (min^{-1})
WT control	47.38	50.9	0.0209
<i>sbella/b-AB</i>	33.29	39.76	0.0158

C_{90} is the proportion of starch digested by α -amylase after 90 min; this is obtained from the starch digestibility curve. C_{∞} is the total extent of starch amylolysis for each digestive phase accounting for Y_0 (endogenous baseline sugar) and k is the rate constant of the reaction; both parameters are obtained applying the LOS analysis to the digestibility data.

The total breakdown of starch represented by C_{∞} and the rate constant (k) of starch digestion were determined from the slope and y-intercept of the LOS plot and are reported in Table 3. Both C_{∞} and k were lower for *sbella/b-AB* pudding than the WT control. These kinetic parameters provided a good fit to experimental data as shown in Fig. 2a: however, the low R^2 values lower our confidence in the accuracy of the rate of digestion measured by the parameter k .

3.4. *In vivo* glycaemic index of pudding

Test meal characteristics are reported on a dry weight basis (DWB) in Table 4. The average pudding portion was 415.3 g \pm 12.9 g (mean \pm SEM of 10 puddings served) for *sbella/b-AB* with 85.1% moisture content and 384.95 g \pm 7.5 g (mean \pm SEM of 10 puddings served) for WT with 83.6% moisture content.

Mutant *sbella/b-AB* pudding had more resistant starch compared to WT pudding (Welch two sample t -test, $p = 0.0005$). The total starch content of WT pudding was found to be 2.3 g



Table 4 Test meals characteristics

Pudding	Semolina (g per serving DWB)	TS (g per serving, DWB)	RS (g per serving, DWB)
WT control	63.14	43.272	2.505
<i>sbeIIa/b</i> -AB	61.76	45.355	4.503

Semolina per pudding serving and starch characteristics of the pudding served on a DWB. TS and RS are calculated from the semolina starch content and adjusted for the measured moisture content of the pudding.

Table 5 Characteristics of the 10 participants completing the study

Males/females (<i>n</i>)	2/8
Age (y)	33 ± 2
Height (cm)	154 ± 17
Weight (kg)	69 ± 3
BMI (kg m ⁻²)	23.5 ± 0.6
Ethnicity	White Caucasian (7) Southwest Asian (3)
Fasting glucose (mmol L ⁻¹)	4.7 ± 0.07
Reference AUC (mmol min L ⁻¹)	198 ± 25
Time taken to eat the test meal (min)	11.8 ± 1.4

Means ± SEMs.

lower than that of the *sbeIIa/b*-AB pudding, possibly due to unexpected water losses during cooking (Welch two sample *t*-test, $p = 0.0347$). The test meals were consumed within approximately 12 minutes. A total of ten participants completed the study and their characteristics are shown in Table 5.

The mean GI and iAUCs of *sbeIIa/b*-AB and WT control puddings are reported in Table 6, and the capillary blood glucose concentrations of 10 participants who completed the study are reported in ESI Table S2.† We found no evidence of difference in GI (mean difference = −13.9 [95CI −17.00758, 44.80758], paired *t*-test, $p = 0.335$). The blood glucose profile after consuming *sbeIIa/b*-AB pudding (iAUC 0–120 min) did not significantly differ from the WT control (mean difference = −24.019 [95CI −23.46, 71.49], paired *t*-test, $p = 0.282$), Fig. 3b. No main effect of *sbeIIa/b*-AB pudding was found for reducing blood glucose concentration (linear mixed-effects model, $p = 0.37011$); however 60 minutes after consumption, *sbeIIa/b*-AB pudding resulted in a lower blood glucose concentration. This may suggest that the effect of the pudding genotype on blood glucose concentration is dependent on time, an interaction (time × genotype) which is significant at 60 minutes (linear




Fig. 3 (a) Postprandial changes in blood glucose concentration after *sbeIIa/b*-AB (*sbeIIa/b*) and wild-type (WT) control pudding meals. (b) iAUC blood glucose. Values are mean deviations from baseline ± SEMs ($n = 10$ individuals).

mixed-effects model, $p = 0.042$) but not overall (linear mixed-effects model, $p = 0.07$), Fig. 3a.

4. Discussion

Our study features a new approach that combines modified starch structures in raw materials with optimised food processing to design healthier starchy foods with steady glucose release. The *sbeIIa/b*-AB mutant wheat semolina used in this study was previously characterised by high-amylose and resistant starch concentrations. However, it was not known to what extent the desirable nutritional properties of *sbeIIa/b*-AB mutant wheat were preserved after processing the wheat into a food (in this case, semolina pudding), nor if these properties would impact *in vivo* glycaemic responses. In this study, we explored starch resistance to digestion *in vitro* after hydrothermal processing of semolina into a pudding and its GI in a pilot study *in vivo*. We also reported the NSP content and composition of *sbeIIa/b*-AB wheat in raw semolina for the first time.

In vitro digestion kinetic analysis of *sbeIIa/b*-AB pudding showed lower starch digestibility (iAUC₉₀), starch breakdown (C_{∞}) and lower rate of digestion (k) at 90 minutes compared to the WT control. A lower *in vitro* starch digestibility would be expected to coincide with a lower glucose response *in vivo*.⁴² Previous studies have shown a direct correlation between the

Table 6 Blood glucose responses after test meals

Meals	Glucose iAUC mmol L ⁻¹ min ⁻¹ (0–30 min)	Glucose iAUC mmol L ⁻¹ min ⁻¹ (0–60 min)	Glucose iAUC mmol L ⁻¹ min ⁻¹ (0–120 min)	Glycaemic index
WT control	18.4 ± 2.6	78.1 ± 7.2	135.0 ± 17.1	73.85 ± 12
<i>sbeIIa/b</i> -AB	19.2 ± 4.2	68.3 ± 10	111.0 ± 14	59.78 ± 7.6

Values are means ± SEM, $n = 10$ individuals.



digestibility index C_{90} obtained from starch digestibility curves and GI values measured *in vivo*. The GI difference between *sbeIIa/b-AB* and WT puddings was not statistically significant but it was consistent with the starch digestibility measured *in vitro* (C_{90}). Unexpected water losses during cooking may have affected the accuracy of the glycaemic load delivered, as the TS content of the pudding servings did not match as anticipated. Although the overall GI values were not statistically different, there was a tendency for lower blood glucose values at 60 minutes after consumption of *sbeIIa/b-AB* pudding. This suggests that while the cooking method may need to be optimised to exploit the potential of the higher RS content in lowering glucose response, consumption of *sbeIIa/b-AB* wheat-based food products may contribute to lower postprandial glycaemia in healthy subjects. Future studies on larger cohorts should focus on measuring the extent of the reduction in postprandial glycaemia.

It is worth noting that the RS content of unprocessed *sbeIIa/b-AB* semolina was lower than that reported in previous studies using the same genotypes.²⁴ The lower RS concentration in this batch of *sbeIIa/b-AB* semolina may have resulted from the environmental effects during the growth of the wheat.⁵⁷ However, it does suggest that the RS concentration in the pudding alone administered in this pilot study may not be sufficient to elicit a lower glycaemic response compared to the WT control. While the use of wheat with a higher content of RS may have been preferable, the authors worked with the available material which was considered to be satisfactory to develop novel durum wheat foods and to explore its effect on glycaemic response.

We observed that RS remained higher in *sbeIIa/b-AB* compared to the WT after cooking semolina into a pudding which was likely due to a combination of factors including starch retrogradation and starch native structure (amylose). *sbeIIa/b-AB* wheat starch may be more prone to gelatinisation than WT wheat starch. Li and colleagues characterised starch physicochemical properties of a range of wheat lines with different amylose concentrations by combining multiple alleles of *SBEII* genes. They reported lower gelatinization enthalpy (ΔH) and greater gelatinization range for high amylose mutant starches compared to the corresponding wild-type control. However, the bread wheat lines analysed by Li *et al.* had considerably a higher amylose content, ranging from 36% (WT) to 93% in line RS100.⁵³ Because of the relatively small difference in amylose concentration between *sbeIIa/b-AB* and WT wheat starch described in this study, a dramatic difference in the degree of gelatinisation seems unlikely.

Cooked and cooled ('retrograded') *sbeIIa/b-AB* semolina had ~2.5-fold more RS compared to the WT control. In this study, starch resistance to digestion was mainly determined by the structural and physical characteristics of raw semolina which influenced the properties of the surrounding food matrix.^{54,55} Milling of *sbeIIa/b-AB* and WT wheat grains resulted in *sbeIIa/b-AB* semolina with an ~8% larger particle size compared to WT semolina. Durum wheat is generally milled to semolina grade, a coarse particle size, to reduce the high starch damage

due to the kernel hardness.⁵⁶ Particle size has previously been shown to have influenced starch gelatinisation, bioaccessibility and postprandial glycaemic responses to starch-based foods.⁵⁷ However, the differences in the size of the fractions used in previous studies were three orders of magnitude larger than the subtle 21 μm difference in the size between mutant *sbeIIa/b-AB* and WT particles used in this study. In this case, particle size difference is unlikely to be a contributing mechanism to the difference in digestibility observed.¹⁵ The milling process was carried out under the same conditions for both WT and *sbeIIa/b-AB*. Because of the novelty of *sbeIIa/b-AB* wheat, we did not anticipate that milling *sbeIIa/b-AB* wheat grains would result in *sbeIIa/b-AB* semolina with a larger particle size. This is possibly one of the characteristics of *sbeIIa/b-AB* semolina which will need to be taken into account when using *sbeIIa/b-AB* durum wheat for food formulations.

sbeIIa/b-AB semolina used in this study was characterised by increased AX and MLG contents, derived from cell wall components. It has been suggested that under certain conditions (such as high molecular weight and high concentration) some NSP components may influence the viscosity of a food matrix and therefore alter the interaction between digestive enzymes and substrates.^{21,22} A higher ratio of G3 and G4 residues in *sbeIIa/b-AB* semolina compared to that in the WT control suggests a change in the MLG structure in *sbeIIa/b-AB* wheat. However, the increase in the MLG content (G3 + G4) and the change in the structure (G3 : G4) measured in raw *sbeIIa/b-AB* semolina are still within the range indicated by De Santis *et al.* for other durum wheat cultivars.⁵⁸ The total AX content of *sbeIIa/b-AB* semolina was higher than that of the WT control and we also observed an increased proportion of substituted AXOS. As AX solubility depends largely on their molecular weight and substitution degree,⁵⁹ a change in AX composition could correspond to a change in viscosity properties. However, due to the relatively small increase in the NSP content, it is unlikely that pudding viscosity was affected by these changes. Increases in NSP concentrations are of interest as they are known to exert health benefits *via* mechanisms linked to gut fermentation (prebiotic activity).⁶⁰ Overall, the difference in NSP concentrations in *sbeIIa/b-AB* wheat is an interesting observation and future studies should take into account their potential role in gut fermentation.

This study essentially provides new hypotheses regarding the applicability of high resistant starch semolina for food production and its potential effects on glucose metabolism. It is also important to consider that pilot studies are usually used to determine the feasibility of a study rather than test a hypothesis. For this reason we did not base our pilot study on a calculated sample size, instead we relied on a well standardised method for GI screening as described by Brouns *et al.* 2005⁴⁵ which suggests that the inclusion of 10 subjects provides a reasonable degree of power and precision to measure the GI. Furthermore, the use of the *sbeIIa/b-AB* semolina described in the study has helped the authors to fine-tune the RS concentration and formulation of wheat foods, which will



be tested in larger follow-up studies to measure the impact of *sbeIIa/b*-AB wheat on glucose response.

5. Conclusions

This is the first time that the glycaemic response to *sbeIIa/b*-AB wheat has been studied within a processed food matrix. In this pilot study, semolina puddings were prepared from WT control and *sbeIIa/b*-AB mutant wheat harvested in bulk and nutritionally relevant properties were examined both *in vitro* and *in vivo*. Using reverse genetic approaches to fine-tune the digestibility of complex carbohydrates in staple crops is becoming a widely accepted method to optimise energy release and improve the nutritional profile of staple foods. We have shown that RS and nutritional properties are influenced by processing. Despite the high RS concentration of *sbeIIa/b*-AB wheat, the GI of *sbeIIa/b*-AB pudding was not significantly different from that of WT control pudding. Nevertheless, there are likely advantages of *sbeIIa/b*-AB wheat such as lower glycaemic load, and increased RS and fibre intake. Future studies should focus on optimising the processing conditions of wheat-based foods with high RS and NSP concentrations. The present study highlights the importance of studying novel wheat with improved nutritional traits in the context of a food matrix to determine its potential impact on health. Further larger studies *in vivo* are needed to determine the physiological effect of *sbeIIa/b*-AB wheat foods and their potential to boost the dietary intake of RS.

Abbreviations

RS	Resistant starch
WT	Wild-type
DWB	Dry weight basis
GI	Glycaemic index
iAUC	Incremental area under the curve
NSPs	Non-starch polysaccharides
AX	Arabinoxylans
TOT-AX	Total arabinoxylans
WE-AX	Water extractable arabinoxylans
TOT-AXOS	Sum of arabinoxylan oligosaccharides
MLG	Mixed-linkage β -glucans

Conflicts of interest

None of the authors reported a conflict of interest related to the study.

Acknowledgements

We thank Dr George Savva (QIB) for statistical guidance, Prof. Peter Wilde (QIB), Dr Cristobal Uauy (JIC), Dr Frederick

Warren (QIB) and Dr Peter Ryden (QIB) for helpful discussions.

Infrastructure support was provided by the NIHR Imperial Biomedical Research Centre and the NIHR Imperial Clinical Research Facility, London (UK). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. The authors would like to thank all volunteers involved in the study.

The authors' responsibilities were as follows: MC, ACM, EC, AP and AW carried out the experiments. MC and EC processed the experimental data and performed the analysis. MC wrote the manuscript with support from CHE, BAH, AL and GSF. CHE and BAH supervised the project. All authors discussed the results and commented on the manuscript.

This work was supported by the UKRI Biotechnology and Biological Sciences Research Council Norwich Research Park Biosciences Doctoral Training Partnership grant number BB/M011216/1. The authors gratefully acknowledge the support of the BBSRC Institute Strategic Programme Food Innovation and Health BB/R012512/1 and its constituent projects BBS/E/F/000PR10343 (Theme 1, Food Innovation) and BBS/E/F/000PR10345 (Theme 2, Digestion in the Upper GI Tract) and previously by the Institute Strategic Programme Grant ('Food and Health'; Grant No. BB/J004545/1) to the Institute of Food Research. The authors would also like to acknowledge the support of the BBSRC Institute Strategic Programme Grants 'Molecules from Nature' – Crop Quality BBS/E/J/000PR9799 to the John Innes Centre and 'Designing Future Wheat' (BB/P016855/1) to Rothamsted Research and the John Innes Centre.

References

- 1 PHE, National Diet and Nutrition Survey 2014/2015 to 2015/2016, <https://www.gov.uk/government/statistics/ndns-results-from-years-7-and-8-combined>.
- 2 F. B. Hu, Are refined carbohydrates worse than saturated fat? 1–3, *Am. J. Clin. Nutr.*, 2010, **91**, 1541–1542.
- 3 F. B. Hu and W. C. Willett, Optimal Diets for Prevention of Coronary Heart Disease, *J. Am. Med. Assoc.*, 2002, **288**, 2569–2569.
- 4 G. Frost and A. Dornhorst, The relevance of the glycaemic index to our understanding of dietary carbohydrates, *Diabetic Med.*, 2000, **17**, 336–345.
- 5 J. H. Cummings, M. B. Roberfroid, H. Andersson, C. Barth, A. Ferro-Luzzi, Y. Ghos, M. Gibney, K. Hermansen, W. James, O. Korver, D. Lairon, G. Pascal and A. Voragen, A new look at dietary carbohydrate: chemistry, physiology and health JH Cummings, MB Roberfroid and members of the Paris Carbohydrate Group, *Eur. J. Clin. Nutr.*, 1997, **51**, 417–423.
- 6 C. S. Byrne, E. S. Chambers, D. J. Morrison and G. Frost, The role of short chain fatty acids in appetite regulation and energy homeostasis, *Int. J. Obes.*, 2015, **39**, 1331–1338.



- 7 A. Buléon, P. Colonna, V. Planchot and S. Ball, Starch granules: structure and biosynthesis, *Int. J. Biol. Macromol.*, 1998, **23**, 85–112.
- 8 F. J. Warren, B. Zhang, G. Waltzer, M. J. Gidley and S. Dhital, The interplay of α -amylase and amyloglucosidase activities on the digestion of starch in in vitro enzymic systems, *Carbohydr. Polym.*, 2015, **117**, 192–200.
- 9 H. Goesaert, K. Brijs, W. S. Veraverbeke, C. M. Courtin, K. Gebruers and J. A. Delcour, Wheat flour constituents: how they impact bread quality, and how to impact their functionality, *Trends Food Sci. Technol.*, 2005, **16**, 12–30.
- 10 R. C. Eerlingen, H. Jacobs and J. A. Delcour, Enzyme-Resistant Starch. V. Effect of Retrogradation of Waxy Maize Starch on Enzyme Susceptibility, *Cereal Chem.*, 1994, **71**, 351–355.
- 11 R. Cui and C. G. Oates, The effect of retrogradation on enzyme susceptibility of sago starch, *Carbohydr. Polym.*, 1997, **32**, 65–72.
- 12 H. Li, M. J. Gidley and S. Dhital, High-Amylose Starches to Bridge the “Fiber Gap”: Development, Structure, and Nutritional Functionality, *Compr. Rev. Food Sci. Food Saf.*, 2019, **18**, 362–379.
- 13 A. Htoon, A. K. Shrestha, B. M. Flanagan, A. Lopez-Rubio, A. R. Bird, E. P. Gilbert and M. J. Gidley, Effects of processing high amylose maize starches under controlled conditions on structural organisation and amylase digestibility, *Carbohydr. Polym.*, 2009, **75**, 236–245.
- 14 C. Hoebler, A. Karinthi, H. Chiron, M. Champ and J. L. Barry, Bioavailability of starch in bread rich in amylose: metabolic responses in healthy subjects and starch structure, *Eur. J. Clin. Nutr.*, 1999, **53**, 360–366.
- 15 C. H. Edwards, M. M. Grundy, T. Grassby, D. Vasilopoulou, G. S. Frost, P. J. Butterworth, S. E. Berry, J. Sanderson and P. R. Ellis, Manipulation of starch bioaccessibility in wheat endosperm to regulate starch digestion, postprandial glycemia, insulinemia, and gut hormone responses: a randomized controlled trial in healthy ileostomy participants 1,2, *Am. J. Clin. Nutr.*, 2015, **102**, 791–800.
- 16 J. M. Lattimer and M. D. Haub, Effects of Dietary Fiber and Its Components on Metabolic Health, *Nutrients*, 2010, **2**, 1266–1289.
- 17 H. N. Englyst and S. M. Kingman, Dietary fiber and resistant starch – A nutritional characterisation of plant polysaccharides, *Diet. Fiber*, 1987, 49–65.
- 18 M. A. De Santis, O. Kosik, D. Passmore, Z. Flagella, P. R. Shewry and A. Lovegrove, Comparison of the dietary fibre composition of old and modern durum wheat (*Triticum turgidum* spp. durum) genotypes, *Food Chem.*, 2018, **244**, 304–310.
- 19 P. R. Shewry, M. J. Hawkesford, V. Piironen, A.-M. Lampi, K. Gebruers, D. Boros, A. A. M. Andersson, P. Åman, M. Rakszegi, Z. Bedo and J. L. Ward, Natural Variation in Grain Composition of Wheat and Related Cereals, *J. Agric. Food Chem.*, 2013, **61**, 8295.
- 20 D. J. Mares and B. A. Stone, Studies on wheat endosperm i. Chemical composition and ultrastructure of the cell walls, *Aust. J. Biol. Sci.*, 1973, **26**, 793–812.
- 21 M. M.-L. Grundy, C. H. Edwards, A. R. Mackie, M. J. Gidley, P. J. Butterworth and P. R. Ellis, Re-evaluation of the mechanisms of dietary fibre and implications for macronutrient bioaccessibility, digestion and postprandial metabolism, *Br. J. Nutr.*, 2016, **116**, 816–833.
- 22 L. Saulnier, N. Peneau and J. F. Thibault, Variability in grain extract viscosity and water-soluble arabinoxylan content in wheat, *J. Cereal Sci.*, 1995, **22**, 259–264.
- 23 I. J. Tetlow and M. J. Emes, A review of starch-branching enzymes and their role in amylopectin biosynthesis, *IUBMB Life*, 2014, **66**, 546–558.
- 24 B. Hazard, X. Zhang, P. Colasuonno, C. Uauy, D. M. Beckles and J. Dubcovsky, Induced mutations in the starch branching enzyme II (SBEII) genes increase amylose and resistant starch content in durum wheat, *Crop Sci.*, 2012, **52**, 1754–1766.
- 25 B. Hazard, X. Zhang, M. Naemeh, M. K. Hamilton, B. Rust, H. E. Raybould, J. W. Newman, R. Martin and J. Dubcovsky, Mutations in Durum Wheat SBEII Genes affect Grain Yield Components, Quality, and Fermentation Responses in Rats, *Crop Sci.*, 2015, **6**, 2813–2825.
- 26 A. R. Bird and A. Regina, High amylose wheat: A platform for delivering human health benefits, *J. Cereal Sci.*, 2018, **82**, 99–105.
- 27 C. Souza da Silva, D. Haenen, S. J. Koopmans, G. J. E. J. Hooiveld, G. Bosch, J. E. Bolhuis, B. Kemp, M. Müller and W. J. J. Gerrits, Effects of resistant starch on behaviour, satiety-related hormones and metabolites in growing pigs, *Animal*, 2014, **8**, 1402–1411.
- 28 P. R. Regmi, T. A. T. G. van Kempen, J. J. Matte and R. T. Zijlstra, Starch with High Amylose and Low in Vitro Digestibility Increases Short-Chain Fatty Acid Absorption, Reduces Peak Insulin Secretion, and Modulates Incretin Secretion in Pigs, *J. Nutr.*, 2011, **141**, 398–405.
- 29 H. J. Willis, A. L. Eldridge, J. Beiseigel, W. Thomas and J. L. Slavin, Greater satiety response with resistant starch and corn bran in human subjects, *Nutr. Res.*, 2009, **29**, 100–105.
- 30 J. Harrold, L. Breslin, J. Walsh, J. Halford and C. Pelkman, Satiety effects of a whole-grain fibre composite ingredient: Reduced food intake and appetite ratings, *Food Funct.*, 2014, **5**, 2574–2581.
- 31 B. Hazard, X. Zhang, M. Naemeh and J. Dubcovsky, Registration of Durum Wheat Germplasm Lines with Combined Mutations in SBEIIa and SBEIIb Genes Conferring Increased Amylose and Resistant Starch, *J. Plant Regist.*, 2014, **8**, 334–338.
- 32 T. Verhoeven, B. Fahy, M. Leggett, G. Moates and K. Denyer, Isolation and characterisation of novel starch mutants of oats, *J. Cereal Sci.*, 2004, **40**, 69–79.
- 33 C. A. Knutson, Evaluation of variations in amylose–iodine absorbance spectra, *Carbohydr. Polym.*, 2000, **42**, 65–72.
- 34 C. A. Knutson and M. J. Grove, Rapid Method for Estimation of Amylose in Maize Starches, *Cereal Chem.*, 0094, **71**, 469–471.
- 35 C. A. Knutson, A Simplified Colorimetric Procedure for Determination of Amylose in Maize Starches, *Cereal Chem.*, 1986, **63**, 89–92.



- 36 J. R. Bromley, M. Busse-Wicher, T. Tryfona, J. C. Mortimer, Z. Zhang, D. M. Brown and P. Dupree, GUX1 and GUX2 glucuronyltransferases decorate distinct domains of glucuronoxylan with different substitution patterns, *Plant J.*, 2013, **74**, 423–434.
- 37 N. Anders, M. D. Wilkinson, A. Lovegrove, J. Freeman, T. Tryfona, T. K. Pellny, T. Weimar, J. C. Mortimer, K. Stott, J. M. Baker, M. Defoin-Platel, P. R. Shewry, P. Dupree and R. A. C. Mitchell, Glycosyl transferases in family 61 mediate arabinofuranosyl transfer onto xylan in grasses, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 989–993.
- 38 N. Anders, M. D. Wilkinson, A. Lovegrove, J. Freeman, T. Tryfona, T. K. Pellny, T. Weimar, J. C. Mortimer, K. Stott, J. M. Baker, M. Defoin-Platel, P. R. Shewry, P. Dupree and R. A. C. Mitchell, Glycosyl transferases in family 61 mediate arabinofuranosyl transfer onto xylan in grasses - Supporting Information, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, 1–10, DOI: 10.1021/jo101696z.
- 39 J. J. Ordaz-Ortiz, F. Guillon, O. Tranquet, G. Dervilly-Pinel, V. Tran and L. Saulnier, Specificity of monoclonal antibodies generated against arabinoxylans of cereal grains, *Carbohydr. Polym.*, 2004, **57**, 425–433.
- 40 J. J. Ordaz-Ortiz, M. F. Devaux and L. Saulnier, Classification of wheat varieties based on structural features of arabinoxylans as revealed by endoxylanase treatment of flour and grain, *J. Agric. Food Chem.*, 2005, **53**, 8349–8356.
- 41 C. H. Edwards, M. Maillot, R. Parker and F. J. Warren, A comparison of the kinetics of in vitro starch digestion in smooth and wrinkled peas by porcine pancreatic α -amylase, *Food Chem.*, 2018, **244**, 386–393.
- 42 C. H. Edwards, N. Cochetel, L. Setterfield, N. Perez-Moral and F. J. Warren, A single-enzyme system for starch digestibility screening and its relevance to understanding and predicting the glycaemic index of food products, *Food Funct.*, 2019, **10**, 4751–4760.
- 43 C. H. Edwards, F. J. Warren, P. J. Milligan, P. J. Butterworth and P. R. Ellis, A novel method for classifying starch digestion by modelling the amylolysis of plant foods using first-order enzyme kinetic principles, *Food Funct.*, 2014, **5**, 2751–2758.
- 44 P. J. Butterworth, F. J. Warren, T. Grassby, H. Patel and P. R. Ellis, Analysis of starch amylolysis using plots for first-order kinetics, *Carbohydr. Polym.*, 2012, **87**, 2189–2197.
- 45 F. Brouns, I. Bjorck, K. N. Frayn, A. L. Gibbs, V. Lang, G. Slama and T. M. S. Wolever, Glycaemic index methodology, *Nutr. Res. Rev.*, 2005, **18**, 145–171.
- 46 R Core Team, R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria, 2014, <http://www.R-project.org/>.
- 47 H. Wickham, *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag, New York, 2016.
- 48 H. Wickham, Reshaping Data with the reshape Package, *J. Stat. Software*, 2007, **21**, 1–20.
- 49 F. Brouns, I. Bjorck, K. N. Frayn, A. L. Gibbs, V. Lang, G. Slama and T. M. S. Wolever, Glycaemic index methodology, *Nutr. Res. Rev.*, 2005, **18**, 145–145.
- 50 D. Bates, M. Mächler, E. Zurich, B. M. Bolker and S. C. Walker, Fitting linear mixed-effects models using lme4, *J. Stat. Software*, 2014, **67**, 1–48.
- 51 N. Anders, K. Stott, P. R. Shewry, J. C. Mortimer, P. Dupree, T. Weimar, A. Lovegrove, R. A. C. Mitchell, J. Freeman, M. D. Wilkinson, T. Tryfona, M. Defoin-Platel, J. M. Baker and T. K. Pellny, Glycosyl transferases in family 61 mediate arabinofuranosyl transfer onto xylan in grasses, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 989–993.
- 52 H. N. Englyst, M. E. Quigley and G. J. Hudson, Determination of dietary fibre as non-starch polysaccharides with gas-liquid chromatographic, high-performance liquid chromatographic or spectrophotometric measurement of constituent sugars, *Analyst*, 1994, **119**, 1497–1509.
- 53 H. Li, S. Dhital, A. J. Slade, W. Yu, R. G. Gilbert and M. J. Gidley, Altering starch branching enzymes in wheat generates high-amylose starch with novel molecular structure and functional properties, *Food Hydrocolloids*, 2019, **92**, 51–59.
- 54 S. Dhital, F. J. Warren, P. J. Butterworth, P. R. Ellis and M. J. Gidley, Mechanisms of starch digestion by α -amylase—Structural basis for kinetic properties, *Crit. Rev. Food Sci. Nutr.*, 2017, **57**, 875–892.
- 55 H. Patel, P. G. Royall, S. Gaisford, G. R. Williams, C. H. Edwards, F. J. Warren, B. M. Flanagan, P. R. Ellis and P. J. Butterworth, Structural and enzyme kinetic studies of retrograded starch: Inhibition of α -amylase and consequences for intestinal digestion of starch, *Carbohydr. Polym.*, 2017, **164**, 154–161.
- 56 J. A. Delcour, C. Bruneel, L. J. Derde, S. V. Gomand, B. Pareyt, J. A. Putseys, E. Wilderjans and L. Lamberts, Fate of Starch in Food Processing: From Raw Materials to Final Food Products, *Annu. Rev. Food Sci. Technol.*, 2010, **1**, 87–111.
- 57 C. H. Edwards, F. J. Warren, G. M. Campbell, S. Gaisford, P. G. Royall, P. J. Butterworth and P. R. Ellis, A study of starch gelatinisation behaviour in hydrothermally-processed plant food tissues and implications for in vitro digestibility, *Food Funct.*, 2015, **6**, 3634–3641.
- 58 M. A. De Santis, O. Kosik, D. Passmore, Z. Flagella, P. R. Shewry and A. Lovegrove, Data set of enzyme fingerprinting of dietary fibre components (arabinoxylan and β -glucan) in old and modern Italian durum wheat genotypes, *Data Brief*, 2018, **16**, 1062–1068.
- 59 C. M. Courtin and J. A. Delcour, Arabinoxylans and Endoxylanases in Wheat Flour Bread-making, *J. Cereal Sci.*, 2002, **35**, 225–243.
- 60 J. A. Delcour, X. Rouau, C. M. Courtin, K. Poutanen and R. Ranieri, Technologies for enhanced exploitation of the health-promoting potential of cereals, *Trends Food Sci. Technol.*, 2012, **25**, 78–86.

