


Cite this: *Nanoscale*, 2020, 12, 7995

Correction: A robust Mn@FeNi-S/graphene oxide nanocomposite as a high-efficiency catalyst for the non-enzymatic electrochemical detection of hydrogen peroxide

Shaktivel Manavalan,^a Jaysiva Ganesamurthi,^a Shen-Ming Chen,^{*a} Pitchaimani Veerakumar^{b,c} and Keerthi Murugan^a

DOI: 10.1039/d0nr90068k

rsc.li/nanoscale

Correction for 'A robust Mn@FeNi-S/graphene oxide nanocomposite as a high-efficiency catalyst for the non-enzymatic electrochemical detection of hydrogen peroxide' by Shaktivel Manavalan et al., *Nanoscale*, 2020, 12, 5961–5972.

The authors have noticed that the original article contains an incorrect version of Fig. 6(b). Therefore, a corrected version of Fig. 6 is provided below:




Fig. 6 (a) CV curves of bare GCE, GO, FeNi-S, Mn@FeNi-S, Mn@FeNi-S/GO-modified GCEs containing 25 μM of H₂O₂ at a scan rate of 50 mV s⁻¹. (b) CV curves of Mn@FeNi-S/GO/GCE with and without addition of 25 μM H₂O₂. (c) CV curves of Mn@FeNi-S/GO-modified GCE at different scan rates ranging from 20–300 mV s⁻¹ in 25 μM of H₂O₂, and (d) the corresponding plot of peak current versus square root of the scan rate. All experiments were conducted in N₂-saturated 0.1 M PB (pH 7.0).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aDepartment of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei 10608, Taiwan, Republic of China.
E-mail: smchen78@ms15.hinet.net; Fax: +886-2-27025238

^bDepartment of Chemistry, National Taiwan University, Taipei 10617, Taiwan, Republic of China

^cInstitute of Atomic and Molecular Sciences, Academia Sinica, Taipei 10617, Taiwan, Republic of China

