Organic & Biomolecular Chemistry



View Article Online

PAPER

Check for updates

Cite this: Org. Biomol. Chem., 2020, **18**, 1760

Ruthenium-catalysed cyclisation reactions of 1,11dien-6-ynes leading to biindenes[†]

Takanori Matsuda 🕩 * and Naoto Yonekubo

Received 28th January 2020, Accepted 15th February 2020 DOI: 10.1039/d0ob00179a

rsc.li/obc

1,2-Bis(2-allylphenyl)ethynes undergo cycloisomerisation reactions in the presence of Cp*Ru(II) catalysts to produce 2,2'-dimethyl-3H,3'H-1,1'-biindenes. On the other hand, tandem ring-closing metathesis of 1,2-bis(2-allylphenyl)ethynes using the Hoveyda–Grubbs 2nd generation catalyst led to the formation of 2,2'-unsubstituted biindenes. Various symmetrical and unsymmetrical bicyclic dienes were prepared by these ruthenium-based cyclisation methods.

Introduction

3H,3'H-1,1'-Biindenes have previously been prepared by the oxidative homocouplings of (1H-inden-1-yl)lithiums to yield diastereomeric mixtures of 1H,1'H-1,1'-biindenes, followed by base-promoted double bond isomerisation.¹ There are fewer than fifty known biindenes, and some of them have been used as ligands for transition metals,² while a biindene-derived diol has been used as a chiral ligand in the titanium(rv)-catalysed enantioselective additions of diethylzinc to aldehydes.³

Transition-metal-catalysed cycloisomerisation reactions of enynes are powerful tools for the synthesis of various carboand heterocyclic compounds.⁴ This method allows for the rapid atom-economical construction of a complex cyclic structure from a linear substrate. The ring-closing metathesis (RCM) of dienes and enynes revolutionised the way in which cycloalkenes are assembled, and has been extremely useful in modern organic synthesis.⁵ Herein, we report that 1,11-dien-6ynes can undergo both cycloisomerisation and tandem RCM reactions catalysed by ruthenium complexes. Notably, these reactions are used to prepare 1,1'-biindenes from 1,2-bis(2allylphenyl)ethynes.

Results and discussion

When 1,2-bis(2-allylphenyl)ethyne $(1a)^6$ was heated at 60 °C in EtOH in the presence of 5 mol% CpRuCl(PPh₃)₂ for 24 h, it cycloisomerised to afford 2,2'-dimethyl-3*H*,3'*H*-1,1'-biindene (2a) in 21% yield (Table 1, entry 1). The use of Cp*RuCl(PPh₃)₂ improved the yield of 2a to 45% (entry 2), while the reaction in the presence of Cp*RuCl(cod) afforded **2a** in 34% yield (entry 3). The use of a cationic ruthenium catalyst generated *in situ* from Cp*RuCl(cod) and NaPF₆ gave **2a** in 55% yield (entry 4); indeed, preformed cationic [Cp*Ru(MeCN)₃]PF₆ exhibited comparable activity (entry 5). The effect of the phosphine ligand was next examined; **2a** was formed in 38% yield when the Cp*RuCl(cod)–BINAP catalyst system was used (entry 6); however, use of P(C₆F₅)₃ increased the yield of **2a** to 61%, together with a 24% yield of the [2 + 2 + 2] cycloadduct **3a** (entry 7).^{7,8} The reaction performed in MeOH in the presence of Cp*RuCl(cod)–P(C₆F₅)₃ furnished **2a** in 85% isolated yield without any noticeable amount of **3a** (entry 8). A similar result was obtained when the reaction was performed at 40 °C (entry 9). Interestingly, the reaction delivered cycloadduct **3a** as the major product when performed in *i*-PrOH (entries 10 and 11).⁹

With the optimised reaction conditions in hand, various diallyl diphenylacetylenes 1b-l bearing substituents on their benzene rings were subjected to the ruthenium-catalysed cycloisomerisation conditions (Table 2). The reaction of 1,2-bis (2-allyl-4-methylphenyl)ethyne (1b) afforded tetramethylbiindene 2b in 67% yield (entry 1), whereas symmetrical dienvnes 1c and 1d bearing methyl or methoxy groups the 5 positions of their benzene rings afforded 2c and 2d, respectively, in good yields (entries 2 and 3). In contrast, the reactions of chloro- and trifluoromethyl-substituted dienynes 1e and 1f formed the [2 + 2 + 2] cycloadducts 3 as major products under the standard conditions (1e: 2e 24% + 3e 41%; 1f: 3f 87%). The cycloisomerisation products from 1e and 1f were obtained as the major products in yields of 44% and 29%, respectively, when the reaction was performed with $[Cp*Ru(MeCN)_3]PF_6$ (entries 4 and 5). The naphthalene derivative 1g was also converted into the corresponding product 2g (entry 6), while unsymmetrically substituted biindenes 2h-l were similarly prepared by cycloisomerising dienynes 1h-l (entries 7-11).

The cycloisomerisation conditions were successfully applied to dienyne **1m** devoid of *o*-phenylene tethers, which

Department of Applied Chemistry, Tokyo University of Science, 1-3 Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan. E-mail: mtd@rs.tus.ac.jp

[†]Electronic supplementary information (ESI) available: Experimental procedures and characterisation data for new compounds. See DOI: 10.1039/d0ob00179a

 Table 1
 Ruthenium-catalysed cycloisomerisation of 1,2-bis(2-allylphenyl)ethyne (1a)^a



Entry	Ru catalyst	Ligand (mol%)	Additive	Solvent	Temp. (°C)	Time (h)	$\operatorname{Yield}^{b}(\%)$ of $2\mathbf{a}$	$\operatorname{Yield}^{b}(\%)$ of $3a$
1	$CpRuCl(PPh_3)_2$		_	EtOH	60	24	21	
2	$Cp*RuCl(PPh_3)_2$	_	_	EtOH	60	4	45	
3	Cp*RuCl(cod)	_	_	EtOH	60	24	34	
4	Cp*RuCl(cod)	_	NaPF ₆	EtOH	60	24	55	
5	[Cp*Ru(MeCN) ₃]PF ₆	_	_	EtOH	60	12	56	
6	Cp*RuCl(cod)	rac-BINAP (5)	_	EtOH	60	24	38	
7	Cp*RuCl(cod)	$P(C_6F_5)_3(10)$	_	EtOH	60	24	61	24
8	Cp*RuCl(cod)	$P(C_6F_5)_3(10)$	_	MeOH	60	24	85	
9	Cp*RuCl(cod)	$P(C_6F_5)_3(10)$	_	MeOH	40	24	87	
10	Cp*RuCl(cod)	$P(C_6F_5)_3(10)$	_	<i>i</i> -PrOH	60	24	11	55
11	Cp*RuCl(cod)	$P(C_6F_5)_3(10)$		<i>i</i> -PrOH	40	24	14	75
^a Reacti	on conditions: 1a (0.050	mmol), ruthenium o	eatalyst (2.5 μ	mol, 5 mol%	6), ligand (Ru : I	P = 1:2, solve	nt (0.5 mL, 0.1 M). ^b	Isolated yield.

led to the formation of 1,1'-bicyclopentene **2m** in 82% yield in the presence of $[Cp*Ru(MeCN)_3]PF_6$ (Scheme 1, (a)). The alternative cycloisomerisation product **4m** was obtained in 55% yield when **1m** was reacted at 0 °C (Scheme 1, (b)).^{10,11} Heating **4m** in the presence of the ruthenium catalyst gave **2m** in 49% yield, but no isomerisation was observed in the absence of the ruthenium catalyst. Based on these results as well as previous studies, we conclude that 2-methylene-1,1'-bi-(cyclopentylidene) **4m** is the initial cycloisomerisation product, and that **4m** is also catalytically isomerised to **2m** by the ruthenium catalyst.

Two possible reaction pathways can be proposed for the ruthenium-catalysed cycloisomerisation of 1,11-dien-6-yne 1 (Scheme 2). Path (a) involves the formation of a hydroruthenium species from the catalyst and MeOH,¹² a Markovnikov hydroruthenation to the C=C bond of 1 to form A, consecutive carboruthenation (through **B** to **C**), β -hydride elimination that releases 4, and the final double bond isomerisation of 4 to afford product 2. On the other hand, in path (b), dienyne 1 first undergoes oxidative cyclisation on ruthenium to generate the ruthenacyclopentene species D. The unreacted alkene moiety in **D** then inserts into the Ru-C(sp²) bond to give the ruthenacycloheptene intermediate E. Subsequent β -hydride elimination (to form **F**) followed by reductive elimination yields 4, which then isomerises to 2 catalysed by a hydroruthenium species. Alternatively, β -hydride elimination from D generates alkenylruthenium hydride G, which also leads to 4 through intramolecular carboruthenation (to \mathbf{F}) or hydroruthenation (to H). Reductive elimination from intermediate **E** is possible, which gives rise to the [2 + 2 + 2] cycloadduct 3.

Dienyne **1n** or **1o**, in which one allyl group is replaced with a crotyl or a methallyl group, was found to be unreac-

tive toward cycloisomerisation, which reveals that the reaction is limited to dienynes with unsubstituted C=C double bonds (Chart 1). Moreover, 1,2-bis[2-(vinyloxy)phenyl] ethyne (1p) also failed to react, and a complex mixture of products was obtained when unsymmetrical dienyne 1q, bearing malonate and *o*-phenylene tethers, was reacted.¹³

We have been interested in the catalytic syntheses of silole derivatives¹⁴ and the cycloisomerisation of bis-silicon-bridged **1r** was envisaged as a method for the synthesis of a bi(1-silaindene).¹⁵ However, the reaction of **1r** under conditions similar to those described above led to a totally different outcome: 1,1',2,2'-tetrahydro-4,4'-bi(1-silanaphthalene) **5** was obtained in 43% yield as the sole product after full conversion of **1r** (Scheme 3). The silanaphthalene **5** may have formed through a stitching reaction mediated by a hydroruthenium species in a manner analogous to the path (a) in Scheme 2, but with initial anti-Markovnikov hydroruthenation.

Tandem ring-closing metathesis (RCM) of 1,11-dien-6ynes that form 1,1'-bicyclopentene derivatives has previously been studied,¹⁶ but those of 1,2-bis(2-allylphenyl) ethynes have, to the best of our knowledge, never been examined. If allowed, this reaction provides a route to 3,3'H-1,1'-biindenes that lack substituents at their 2 and 2' positions, which is complementary to the cycloisomerisation of 1. Tandem RCM of **1a** in the presence of the Hoveyda– Grubbs 2nd generation catalyst at 100 °C in toluene (0.1 M) afforded the desired biindene **6a** in 60% yield (Table 3, entry 1). A lower concentration of **1a** resulted in an improved yield of **6a**, and 0.02 M was found to be optimal for the present reaction (entries 2 and 3). Other Grubbs catalysts were not suitable for this transformation (entries 4 and 5), while the reaction with 3 mol% catalyst gave a

Organic & Biomolecular Chemistry









Scheme 2 Possible reaction pathways for the cycloisomerisation of 1m (X = C(CO₂Me)₂).



Chart 1 Dienynes that failed to undergo ruthenium-catalysed cycloisomerisation.

^{*a*} Isolated yield (average of two runs). ^{*b*} 5 mol% [Cp*Ru(MeCN)₃]PF₆ was used as catalyst. ^{*c*} Reaction was performed at 60 °C. ^{*d*} The crude reaction mixtures contained byproducts such as 3. ^{*e*} Reaction was performed at 80 °C in MeOH (0.05 M).

similar result (entry 6). As for the reaction temperature, $100 \, ^{\circ}$ C was found to be the best among those examined for the RCM of 1 (entries 6–8).

Various diallyl diphenylacetylenes **1b-l**, which were successfully cycloisomerised (*vide supra*), were examined under the RCM conditions (Table 4). Symmetrical (**1b-g**) and unsymme-



Scheme 3 Cycloisomerisation of 1r.

Table 3 Tandem RCM of 1a

	Grubbs metathe toluen	is catalyst			
Entry	Grubbs catalyst (mol%)	Conc. (M)	Temp. (°C)	Time (h)	Yield ^a (%)
1	Hoveyda–Grubbs 2nd cat. (5)	0.1	100	0.5	60
2	Hoveyda–Grubbs 2nd cat. (5)	0.04	100	1	67
3	Hoveyda–Grubbs 2nd cat. (5)	0.02	100	3	77
4	Grubbs 2nd cat. (5)	0.02	100	6	44
5	Stewart–Grubbs cat. (5)	0.02	100	6	26
6	Hoveyda–Grubbs cat. 2nd (3)	0.02	100	6	82
7	Hoveyda–Grubbs cat. 2nd (3)	0.02	90	6	67
0	Hovevda-Grubbs cat 2nd (3)	0.02	110	6	74

trical (1h-l) dienynes were converted through tandem RCM into biindenes 6b-l in yields ranging from 63% to 96%. Furthermore, dienynes 10-q, which failed to cycloisomerise, also reacted to afford the corresponding metathesis products 40-q, respectively, in good yields. However, the attempted tandem RCM of the bis-silicon-bridged 1r resulted in no conversion under various metathesis conditions.

Conclusions

In conclusion, we developed ruthenium-catalysed cycloisomerisation and tandem-RCM methods for the synthesis of bicyclic conjugated dienes, in which two rings (cycloalkenes) are constructed. Cycloisomerisation of 1,11-dien-6-ynes afforded 2,2'-dimethyl-[1,1'-bi(cyclopentene)] derivatives catalysed by Cp*Ru. On the other hand, 2,2'-unsubsituted bicyclopentenes were prepared through the tandem RCM of 1,11-dien-6-ynes with the Hoveyda–Grubbs catalyst.¹⁷

able 4	Tandom	DCM	of diam	mor 1
able 4	Tandem	RCM	of alenv	nes I

1

8

9

10

11



Table 4 (Contd.)



^{*a*} Isolated yield (average of two runs).

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by JSPS, Japan (Grant-in-Aid for Scientific Research (C) No. 16K05783).

Notes and references

- 1 P. Nicolet, J.-Y. Sanchez, A. Benaboura and M. J. M. Abadie, *Synthesis*, 1987, 202–203.
- 2 (a) T. R. Kelly and P. Meghani, J. Org. Chem., 1990, 55, 3684–3688; (b) R. C. Kerber and B. Waldbaum, Organometallics, 1995, 14, 4742–4754; (c) B. R. Waldbaum and R. C. Kerber, *Inorg. Chim. Acta*, 1999, 291, 109–126; (d) D. Tews and P. E. Gaede, Organometallics, 2001, 20, 3869–3875.
- 3 X.-w. Yang, J.-h. Shen, C.-s. Da, H.-s. Wang, W. Su, D.-x. Liu, R. Wang, M. C. K. Choi and A. S. C. Chan, *Tetrahedron Lett.*, 2001, **42**, 6573–6575.
- 4 For recent reviews on transition-metal-catalysed cycloisomerisations, see: (*a*) C. Aubert, L. Fensterbank, P. Garcia,

Organic & Biomolecular Chemistry

M. Malacria and A. Simonneau, Chem. Rev., 2011, 111, 1954-1993; (b) I. D. G. Watson and F. D. Toste, Chem. Sci., 2012, 3, 2899-2919; (c) Y. Yamamoto, Chem. Rev., 2012, 112, 4736–4769; (d) A. Marinetti, H. Jullien and A. Voituriez, Chem. Soc. Rev., 2012, 41, 4884-4908; (e) D.-H. Zhang, Z. Zhang and M. Shi, Chem. Commun., 2012, 48, 10271-10279; (f) D. P. Day and P. W. H. Chan, Adv. Svnth. Catal., 2016, 358, 1368-1384: (g) C. I. Stathakizis and A. L. Zografos, Nat. Prod. Rep., 2016, 33, 1093-1117; (h) Y. Hu, M. Bai, Y. Yang and Q. Zhou, Org. Chem. Front., 2017, 4, 2256-2275.

- 5 *Handbook of Metathesis*, ed. R. H. Grubbs, Wiley-VCH, Weinheim, 2003, vol. 2.
- 6 Dienynes **1** were easily prepared by the Stille coupling of 1,2-bis(2-bromophenyl)ethynes with allyltributylstannane.
- 7 For ruthenium-catalysed [2 + 2 + 2] cycloaddition of 1,11-dien-6-ynes, see: (a) D. Tanaka, Y. Sato and M. Mori, *J. Am. Chem. Soc.*, 2007, 129, 7730–7731. For the reaction catalysed by rhodium complexes, see: (b) H. Sagae, K. Noguchi, M. Hirano and K. Tanaka, *Chem. Commun.*, 2008, 3804–3806.
- 8 Results with other phosphine ligands (2a/1a + 3a): PPh₃ (12%/30%); P(*c*-Hex)₃ (8%/38%); P(OPh)₃ (18%/69%); DPPE (6%/30%); DPPBZ (25%/62%).
- 9 The formation of a ruthenium hydride might be suppressed in *i*-PrOH in our case; hence the reaction is directed to proceed via the path (b) in Scheme 2. The results in other solvents are as follows: *t*-BuOH (2a not detected, 3a 11%); no reaction in (CF₃)₂CHOH.
- 10 A similar cycloisomerisation catalysed by a rhodium(1) complex has been reported. D. S. Perekalin, N. V. Shvydkiy, Y. V. Nelyubina and A. R. Kudinov, *Chem. Eur. J.*, 2015, 21, 16344–16348. See alsoA. A. Suleymanov, D. V. Vasilyev, V. V. Novikov, Y. V. Nelyubina and D. S. Perekalin, *Beilstein J. Org. Chem.*, 2017, 13, 639–643.
- 11 The corresponding double bond isomers were also detected when 1k and 1l were reacted in the presence of [Cp*Ru(MeCN)₃]PF₆.
- 12 Y. Yamamoto, Y. Nakagai, N. Ohkoshi and K. Itoh, J. Am. Chem. Soc., 2001, **123**, 6372–6380.
- 13 See the ESI[†] for other unsuccessful cycloisomerisation substrates.
- 14 (a) T. Matsuda, Y. Suda and Y. Fujisaki, *Synlett*, 2011, 813–816; (b) T. Matsuda, Y. Yamaguchi, M. Shigeno, S. Sato and M. Murakami, *Chem. Commun.*, 2011, 47, 8697–8699; (c) T. Matsuda and Y. Ichioka, *Org. Biomol. Chem.*, 2012, 10, 3175–3177.
- 15 S. Yamaguchi, C. Xu, H. Yamada and A. Wakamiya, *J. Organomet. Chem.*, 2005, **690**, 5365–5377.
- 16 (a) S.-H. Kim, W. J. Zuercher, N. B. Bowden and R. H. Grubbs, *J. Org. Chem.*, 1996, 61, 1073–1081;
 (b) H. Clavier, A. Correa, E. C. Escudero-Adán, J. Benet-Buchholz, L. Cavallo and S. P. Nolan, *Chem. – Eur. J.*, 2009, 15, 10244–10254.
- 17 For a related study that employs dienes, see: D. Sémeril,C. Bruneau and P. H. Dixneuf, *Helv. Chim. Acta*, 2001, 84, 3335–3341.

8