



Cite this: *Org. Biomol. Chem.*, 2020, **18**, 6086

Influence of the reducing-end anomeric configuration of the Man₉ epitope on DC-SIGN recognition†

Noelia de la Cruz,^a Javier Ramos-Soriano,*^a José J. Reina,^a José L. de Paz,^a Michel Thépaut,^b Franck Fieschi,^b Ana Sousa-Herves^a and Javier Rojo *^a

High-mannose (Man₉GlcNAc₂) is the main carbohydrate unit present in viral envelope glycoproteins such as gp120 of HIV and the GP1 of Ebola virus. This oligosaccharide comprises the Man₉ epitope conjugated to two terminal *N*-acetylglucosamines by otherwise rarely-encountered β-mannose glycosidic bond. Formation of this challenging linkage is the bottleneck of the few synthetic approaches described to prepare high mannose. Herein, we report the synthesis of the Man₉ epitope with both alpha and beta configurations at the reducing end, and subsequent evaluation of the impact of this configuration on binding to natural receptor of high-mannose, DC-SIGN. Using fluorescence polarization assays, we demonstrate that both anomers bind to DC-SIGN with comparable affinity. These relevant results therefore indicate that the more synthetically-accessible Man₉ alpha epitope may be deployed as ligand for DC-SIGN in both *in vitro* and *in vivo* biological assays.

Received 6th July 2020,
Accepted 27th July 2020

DOI: 10.1039/d0ob01380c

rsc.li/obc

Introduction

Carbohydrates are ubiquitous in nature and participate in many biological events relevant to health and disease, such as cell growth and differentiation, fertilization, inflammation, tumour progression and metastasis, viral infection, and many others.¹ The function of carbohydrates in these processes results from interactions with specific receptors, mainly proteins known as lectins.² Carbohydrate-protein interactions are characterised by a high selectivity and a low affinity, the latter of which is compensated by multivalent interactions resulting from the presentation of multiple copies of the carbohydrate epitopes and receptors in nature.³ Understanding carbohydrate-mediated processes therefore requires the accessibility of multivalent carbohydrate platforms capable of intervening with these processes.^{4–6}

High-mannose (Man₉GlcNAc₂) is a complex relevant *N*-glycan present in several viral envelope glycoproteins (Fig. 1). This sugar plays a crucial role during the attachment of pathogens to cells in the first stages of the infection process through the interaction with the DC-SIGN (Dendritic Cell-

Specific ICAM-3 Grabbing Non-integrin) receptor, found in lipid patches at the dendritic cell (DC) surface.⁷ In particular, gp120 of HIV-1 has an average of 24 *N*-linked glycans clustered on the protein surface, of which 53–76% are high-mannose derivatives.⁸ This cluster presentation is fundamental to the efficient interaction with DC-SIGN.




Fig. 1 Chemical structure of high-mannose and its constitutive parts, the beta Man₉ and the GlcNAc₂ epitopes.

^aGlycosystems Laboratory, Instituto de Investigaciones Químicas (IIQ), CSIC – Universidad de Sevilla, Av. Américo Vespucio 49, Seville 41092, Spain. E-mail: javiramossoriano@gmail.com, javier.rojo@iiq.csic.es

^bUniv. Grenoble Alpes, CNRS, CEA, Institut de Biologie Structurale, 38000 Grenoble, France

† Electronic supplementary information (ESI) available: ¹H and ¹³C NMR spectra and selected HSQC and MS spectra. See DOI: 10.1039/D0OB01380C



Towards better understanding the recognition process between high-mannose and DC-SIGN, several multivalent systems have been envisaged. However, the structural complexity of the constituent sugars and the difficulty of obtaining sufficient quantities of pure material from natural sources renders the preparation of multivalent systems carrying the natural beta epitope rather challenging. In fact, very few examples are reported in the literature describing the synthesis of Man₉ and high-mannose multivalent systems.^{9–11} In this context, different approaches have been developed for the preparation of simpler compounds that mimic the multivalent presentation of the sugar found in nature. Small fragments of the Man₉ epitope have been synthesised and evaluated in different glycomimetic systems in pursuit of sugars simpler than Man₉ but with reasonable binding affinities for DC-SIGN.^{12–15} Recently, we have developed a straightforward synthetic strategy for the preparation of the Man₉ epitope with the natural beta configuration at the reducing end.¹⁶ Although the synthesis that we have described is competitive with those published previously by other laboratories, the preparation of the β-mannose unit remains the limiting step. The synthesis of the appropriately-functionalised beta-mannose monosaccharide building block, required to perform the necessary glycosylation steps towards the preparation of the nonasaccharide, demands eleven synthetic steps with a global yield of 43% from *S*-tolyl-α-D-mannopyranoside.¹⁶ As a possible solution, we considered whether the more synthetically-accessible α-mannose analogue would provide a similar affinity to DC-SIGN than the beta anomer, therefore circumventing the requirement for the complicated β-mannose building block, in the preparation of the corresponding carbohydrate multivalent systems.

To evaluate our proposed strategy, we have synthesised the alpha and beta anomers of the Man₉ epitope and the corresponding trivalent glycocluster to examine the differences in binding affinities to the DC-SIGN receptor using a fluorescence polarization assay.

Results and discussion

Preparation of Man₉ epitopes and glycoclusters

The synthesis of beta Man₉ using a convergent straightforward synthetic strategy was previously described by our laboratory.¹⁶ In this work, we used the same strategy to prepare the corresponding alpha anomer.

The di-, penta-, and trisaccharides building blocks were common intermediates for both anomers, (Fig. 2) being the mannose of the reducing end with the alpha configuration in the anomeric position prepared as described in Scheme 1.

The synthesis of the appropriately-protected reducing end mannose was achieved as depicted in Scheme 1, starting from the previously described mannosyl derivative 5.¹⁷ Mannose 5 was orthogonally protected using a consecutive approach, performing purification only at the final step. Positions 4 and 6 were protected *via* the formation of the benzylidene acetal by





Fig. 2 Retrosynthetic scheme for the preparation of αMan₉ and βMan₉.




Scheme 1 Consecutive synthesis of mannose derivative 6.

subjecting 5 to benzaldehyde dimethyl acetal and camphorsulfonic acid (CSA). Subsequently, position 3 was protected as the *p*-methoxybenzyl ether, *via* the initial formation of a tin acetal between position 2 and 3, followed by reaction with *p*-methoxybenzyl chloride (PMBCl) and tetrabutyl ammonium iodide (TBAI). The free OH in position 2 was then protected by benzylation with benzoyl anhydride in the presence of triethyl amine and a catalytic amount of dimethylamino pyridine (DMAP). Finally, the position 3 was selectively deprotected with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) to yield the target mannose derivative 6 in 26% overall yield from mannose 5 following final chromatographic purification.

With the monosaccharide 6 in hand, glycosylation with the previously synthesised trisaccharide 7 and pentasaccharide 8,¹⁶ enabled the formation of the protected Man₉ nonasaccharide 11 (Scheme 2).

The glycosylation between trisaccharide 7 and mannose derivative 6 was performed using *N*-iodo succinimide and trifluoromethanesulfonic acid as the glycosylation promotor to obtain tetrasaccharide 9 in 88% yield. Next, the benzylidene acetal was removed in acidic media using *p*-TsoH to afford the tetrasaccharide 10 (bearing unprotected hydroxyl groups at positions 4 and 6 of the terminal mannose at the reducing end) in 78% yield. The final glycosylation between tetrasaccharide 10 and pentasaccharide 8 took place only at the position 6 hydroxyl group of the acceptor, since this is more reactive than the sterically hindered hydroxyl group at position 4. This glycosylation was carried out using the same conditions




Scheme 2 Synthesis of the α Man₉ 12.

as above to yield the protected nonasaccharide **11** in 73% yield. The nonasaccharide **11** was characterised by ^1H and ^{13}C NMR and ESI-MS. Finally, global deprotection step with NaOMe and 2 M NaOH in MeOH and toluene cleaved all *O*-benzoyl groups affording the α Man₉ **12** in excellent yield (Scheme 2). In this way, the synthesis of the alpha anomer of Man₉ was achieved in an expedient manner through a convergent strategy. Both Man₉ epitopes (alpha and beta) were prepared with a short spacer at the anomeric position functionalised with a terminal azido group. This group permits conjugation to multivalent scaffolds using the Cu(I) azide–alkyne cycloaddition (CuAAC) reaction.^{20,21}

To examine the effect that the multivalent presentation has on the binding DC-SIGN to both anomers, we prepared the trivalent glycoclusters with the alpha and beta Man₉ epitopes (Scheme 3).

For this purpose, we employed a trialkynylated pentaerythritol scaffold, frequently used for the preparation of carbohydrate multivalent systems by our group. This scaffold **14** was prepared in two steps from pentaerythritol as previously reported.¹⁸ The coupling of α Man₉ and β Man₉ ligands was then carried out *via* a click chemistry¹⁹ CuAAC reaction promoted by CuSO₄, sodium ascorbate and tris[(1-benzyl-1*H*-1,2,3-triazol-4-yl)methyl]amine (TBTA) under mild conditions to obtain the corresponding trivalent glycoclusters. These multivalent systems were purified by treatment with Quadrasil MP resin to remove the copper catalyst and by G50 Sephadex chromatography to afford the glycoclusters **15** and **16** in good yields (Scheme 3). The final step was the substitution of the chlorine atom for an azido group at the end of the linker at the

Scheme 3 Synthesis of the trivalent glycoclusters **17** and **18**.

focal position of the scaffold. This reaction was performed using an excess of NaN₃ in DMF at 70 °C for two days furnish azido-functionalised glycoclusters **17** and **18** in excellent yields.

Finally, it was necessary to introduce a chromophore for the fluorescence polarization assays. An alkynyl derivative of fluorescein, the commercially available FAM-alkyne 6-isomer **19**, was therefore conjugated to the α Man₉ **12**, the β Man₉ **13**, and their corresponding trivalent glycoclusters **17** and **18**, respectively, by a CuAAC click reaction promoted by CuBr and TBTA in DMSO at room temperature (Scheme 4).


The compounds were treated with Quadrasil MP resin and then, submitted to a LH20 Sephadex chromatography to deliver the fluorescently-labelled compounds **20**, **21**, **22** and **23** in excellent yields.

With the fluorescent tool compounds prepared, we evaluated their capacity to interact with DC-SIGN, the natural receptor of this Man₉ ligand.

Fluorescence polarization assays

Fluorescence polarization is widely used to study binding events owing to several advantages of this technique.^{22,23} The assay requires only small quantities of ligands and receptors, is well suited for high-throughput screening and does not require the attachment of any of the partners to a surface, allowing ligands and receptors diffuse freely in solution. In particular, fluorescence polarization experiments have been successfully employed in the analysis of carbohydrate–protein interactions.^{24–27} Thus, we selected this assay to evaluate the binding affinity of our alpha and beta epimers of Man₉ (**20** and **21**) and their corresponding multivalent systems (**22** and **23**) for DC-SIGN using the tetraivalent Extracellular Domain (ECD) of the lectin. For this, we have used a microtiter plate where different concentrations of the protein (DC-SIGN ECD), ranging from 75 nM to 28 μM in Tris buffer, were added to the





Scheme 4 Synthesis of fluorescence labelled compounds **20**, **21**, **22** and **23**.

wells containing a fixed final 10 nM concentration of appropriate fluorescent ligands (**20–23**). We observed an increase in the fluorescence polarization value with increasing protein concentrations, demonstrating that the fluorescence ligands bound to DC-SIGN ECD. The resulting Langmuir isotherm curves are represented in Fig. 3. From these curves, the K_D of the binding process for each ligand was calculated. For the α and β anomers of Man_9 , similar values of K_D , (5.2 ± 1.2 and $4.6 \pm 1.3 \mu\text{M}$, respectively) were found demonstrating that the configuration of the anomer in the mannosyl unit at the reducing end of the oligosaccharide does not play a critical role in the binding process.




Fig. 3 Langmuir isothermal curves of binding between DC-SIGN ECD and fluorescence compounds: (a) αMan_9 (**20**); (b) βMan_9 (**21**); (c) trivalent glycocluster **22**; and (d) trivalent glycocluster **23**.

In order to evaluate the influence of the anomer configuration when the Man_9 ligand is presented in a multivalent scaffold, fluorophore labelled glycoclusters **22** and **23** were tested in the fluorescence polarization assays (Fig. 3). In this case, the K_D were one order of magnitude lower than the data found for the monovalent systems, indicating a clear multivalent effect (0.53 ± 0.09 and $0.37 \pm 0.04 \mu\text{M}$ for the trivalent αMan_9 and βMan_9 , respectively). Again, no significant differences between the dissociation constants of the two anomers were found as in the case of the monovalent ligands.

Conclusions

The stereochemical configuration of sugars is fundamental to their selective recognition by lectins. Indeed, just a simple difference in the configuration of one position can drive the recognition by one specific lectin, *e.g.* glucose (with all OH groups in equatorial disposition) *versus* galactose (which bears an axial OH at C4). Furthermore, the configuration of the glycosidic bonds between the monosaccharidic units that form an oligosaccharide are relevant for their recognition and function. In the case of high mannose, all the mannoses of the Man_9 epitope display the alpha configuration of the glycosidic bonds, as usual for this type of sugar. However, the linkage between the Man_9 epitope and the GlcNAc disaccharide exhibits a beta configuration. This glycosidic bond configuration is rare for mannoses and presents a significant hurdle towards addressing this bond configuration *via* chemical synthesis. All *N*-glycans have this moiety as connection with the corresponding Asn in the *N*-glycan proteins. This specific arrangement is critical for the proper orientation of the mannosyl cluster, providing a rigid structure on the protein surface to optimize the binding to DC-SIGN. However, when new glyco-multivalent systems are designed, the glycan epitope is normally conjugated to the multivalent scaffold with a flexible



linker and therefore the specific orientation provided by the beta-mannose linkage is likely not necessary. To demonstrate this hypothesis, we have synthesised the Man₉ epitope of high mannose with alpha and beta configuration in the terminal mannose at the reducing end (20 and 21) and the corresponding trivalent glycoclusters (22 and 23). These ligands were labelled with a fluorescein derivative and their binding to DC-SIGN evaluated by fluorescence polarization assays. Our findings have shown that the configuration, alpha or beta, of the anomeric position of the terminal mannose does not influence in the binding affinity for the natural receptor DC-SIGN and therefore they can be used interchangeably, meaning the more synthetically-accessible alpha-linked epimers can be deployed instead of the more challenging natural beta analogues. We anticipate this knowledge is of key relevance in the design of new glycosystems of high mannose, since notably reduces the complexity, the time and the cost of the glycan synthesis by circumventing the synthetic bottleneck of the beta mannose preparation.

Experimental

Materials and methods

Solvents were HPLC grade and used as received unless otherwise stated. Size exclusion chromatography was performed with Sephadex LH20, G-25 and G-50 (GE Healthcare). Thin-layer chromatography (TLC) was performed on silica plates (Merck). Reagents and QuadraSil® MP were purchased from Sigma Aldrich. 2-Azidoethyl α -D-mannopyranoside¹⁷ and trialkynylated scaffold 14¹⁸ were synthesised as previously described. NMR experiments were performed using a Bruker Advance DRX 400 instrument. NMR chemical shifts are reported in ppm (δ units) downfield from the CDCl₃ signal or the HOD peak (D₂O). 2D experiments (COSY and HSQC) were performed when necessary. NMR spectra were analysed with MestreNova software.

Synthesis of ligands

2-Azidoethyl 2-O-benzoyl-4,6-O-benzylidene- α -D-mannopyranoside (6). To a solution of 2-azidoethoxy- α -D-mannopyranose (5) (800 mg, 3.21 mmol) in CH₃CN (8 mL), CSA (186 mg, 0.8 mmol) and benzaldehyde dimethyl acetal (0.5 mL, 3.55 mmol) were added at rt. The reaction mixture was stirred overnight and the mixing became solidified during addition of the reagent, which indicates the progress of the reaction. The reaction mixture was quenched with Et₃N (250 μ L) and dissolved in excess of EtOAc and water. The reaction mixture was extracted twice with EtOAc and the combined organic layers were dried over anh. MgSO₄, filtered and concentrated under vacuum. After being dissolved in anh. toluene (29 mL), dibutyltin oxide (879 mg, 3.55 mmol) was added to the resulting mixture containing the 4,6-O-benzylidene derivative. The reaction mixture was kept under reflux at 110 °C for 4 h, cooled to rt and PMBCl (0.5 mL, 3.55 mmol) and TBAI (370 mg, 3.55 mmol) were added to it under Ar atmosphere. The reac-

tion mixture was kept under reflux at 110 °C for 1 h. After removal the solvent, the crude was diluted with CH₂Cl₂ and washed with water. The organic layer was dried over anh. MgSO₄, filtered and concentrated under vacuum. To the resulting mixture containing 4,6-O-benzylidene-3-O-*p*-methoxybenzyl derivative in CH₂Cl₂ (15 mL), Bz₂O (1.5 g, 6.42 mmol), Et₃N (1.3 mL, 9.63 mmol) and a catalytic amount of DMAP were subsequently added and the reaction was stirred at rt for 1 h. The reaction mixture was washed with NaHCO₃ sat. aq. soln. The organic phase was dried over anh. MgSO₄, filtered and concentrated under vacuum. Finally, to a stirred solution of the α -D-mannoside derivative in CH₂Cl₂ (70 mL) and water (2.7 mL), DDQ (2.23 g, 9.63 mmol) was added at rt. After 1 h, NaHCO₃ sat. aq. soln. was added, and the mixture was extracted with CH₂Cl₂. The extract was washed several times with NaHCO₃ sat. aq. soln. and then dried over anh. MgSO₄, filtered and concentrated under vacuum. The crude was purified by column chromatography on silica gel (EtOAc/*n*-hexane 1 : 3) to give the compound 6 (320 mg, 0.72 mmol, 26%) as a colourless oil. $[\alpha]_D = -37$ (*c* 1.00, CHCl₃). ¹H-NMR (400 MHz, CDCl₃) δ : 8.11 (m, 2H, H-Ar), 7.60 (m, 1H, H-Ar), 7.56–7.45 (m, 4H, H-Ar), 7.43–7.35 (m, 3H, H-Ar), 5.67 (s, 1H, H_{acetal}), 5.52 (dd, $J_{2,3} = 3.6$, $J_{2,1} = 1.6$ Hz, 1H, H-2), 5.01 (d, $J_{1,2} = 1.6$ Hz, 1H, H-1), 4.41 (dt, $J_{3,4} = 9.6$, $J_{3,2} = 3.9$ Hz, 1H, H-3), 4.33 (dd, $J_{6a,6b} = 9.9$, $J_{6a,5} = 4.4$ Hz, 1H, H-6a), 4.06 (t, $J_{4,3} = J_{4,5} = 9.5$ Hz, 1H, H-4), 4.03–3.84 (m, 3H, H-5, H-6b, H-1'a), 3.73–3.63 (m, 1H, H-1'b), 3.55–3.40 (m, 2H, H-2'), 2.32 (d, $J_{OH,3} = 4.2$ Hz, 1H, OH). ¹³C-NMR (100 MHz, CDCl₃) δ : 166.0, 137.1, 133.5, 129.9, 129.5, 129.3, 128.5, 128.3, 126.3, 102.2, 98.7, 79.2, 72.5, 68.7, 67.1, 66.9, 63.8, 50.4. ESI-MS *m/z* calcd for C₂₂H₂₃N₃O₇: 441.2; found: 464.2 [M + Na]⁺.

Tetrasaccharide 9. To a solution of acceptor 6 (49 mg, 0.11 mmol), donor 7 (275 mg, 0.17 mmol) and 4 Å molecular sieves in anh. CH₂Cl₂ (5.8 mL) were added and the mixture was stirred at –20 °C for 30 min. Then, NIS (39 mg, 0.17 mmol) and TfOH (2.9 μ L, 0.03 mmol) were added and the reaction was stirred at –20 °C for 15 min. The reaction was quenched with NaHCO₃ sat. aq. soln. The reaction mixture was filtered through Celite and washed several times with CH₂Cl₂. The organic layer was washed with Na₂S₂O₃ sat. aq. soln. and then dried over anh. MgSO₄, filtered and concentrated under vacuum. The crude was purified by column chromatography on silica gel (EtOAc/*n*-hexane 1 : 2 to 1 : 1.25) to give the tetrasaccharide 9 (193 mg, 0.10 mmol, 88%) as a white solid. $[\alpha]_D = -22$ (*c* 1.00, CHCl₃). ¹H-NMR (400 MHz, CDCl₃) δ : 8.22 (m, 1H, H-Ar), 8.19–8.12 (m, 2H, H-Ar), 8.09–7.92 (m, 7H, H-Ar), 7.92–7.81 (m, 7H, H-Ar), 7.76–7.68 (m, 2H, H-Ar), 7.64–7.19 (m, 36H, H-Ar), 7.14–6.95 (m, 4H), 6.78 (m, 1H), 6.05–5.81 (m, 5H), 5.79–5.69 (m, 2H), 5.64 (dd, $J = 9.2$, $J = 3.1$ Hz, 1H), 5.57 (s, 1H, H-1), 5.49 (br s, 1H, H_{acetal}), 5.35 (s, 1H, H-1), 5.05 (d, $J = 1.5$ Hz, 1H, H-1), 4.91 (s, 1H, H-1), 4.72–4.50 (m, 4H), 4.47 (t, $J = 2.8$ Hz, 1H), 4.38 (m, 1H), 4.35–4.13 (m, 4H), 4.07–3.94 (m, 3H), 3.94–3.79 (m, 3H), 3.65 (m, 1H), 3.35 (m, 2H). ¹³C-NMR (100 MHz, CDCl₃) δ : 166.4, 166.0, 165.9, 165.8, 165.5, 165.3, 165.3, 165.2, 165.2, 165.0, 164.8, 137.0, 133.7, 133.5, 133.3, 133.2, 133.1, 133.1, 133.1, 133.0, 132.8,



130.1–129.6, 129.3, 129.2, 129.2, 129.1, 128.9–128.3 101.9, 99.8, 90.1, 79.1, 71.9, 71.4, 70.4, 70.1, 69.8, 69.7, 69.5, 68.8, 67.6, 67.2, 67.0, 66.3, 64.1, 63.6, 63.5, 62.9, 62.5, 50.4. ESI-MS m/z calcd for $C_{110}H_{93}N_3O_{32}$: 1967.6; found: 1990.4 $[M + Na]^+$. ESI-HRMS m/z calcd for $C_{110}H_{93}N_3O_{32}Na$ $[M + Na]^+$: 1990.5634; found: 1990.5609

Tetrasaccharide 10. *p*-Toluenesulfonic acid monohydrate (25 mg, 0.13 mmol) was added to a stirred solution of tetrasaccharide **9** (173 mg, 0.09 mmol) in CH_3CN (4.4 mL) at rt. After 24 h, the reaction mixture was quenched with Et_3N (30 μ L) and concentrated under vacuum. The residue was purified by column chromatography on silica gel (acetone/toluene 1 : 5) to give the tetrasaccharide **10** (130 mg, 0.07 mmol, 78%) as a white amorphous solid. $[\alpha]_D = -15$ (c 1.00, $CHCl_3$). 1H -NMR (400 MHz, $CDCl_3$) δ : 8.14 (m, 2H, H-Bz), 8.07–7.93 (m, 13H, H-Bz), 7.90–7.82 (m, 4H, H-Bz), 7.70 (m, 2H, H-Bz), 7.61–7.18 (m, 34H, H-Bz), 6.13 (t, $J = 10.2$ Hz, 1H), 6.03–5.90 (m, 3H), 5.84–5.72 (m, 2H), 5.67–5.57 (m, 3H, 1H-1), 5.52 (d, $J = 1.6$ Hz, 1H, H-1), 5.09 (br s, 1H, H-1), 5.00 (d, $J = 1.7$ Hz, 1H, H-1), 4.73–4.31 (m, 12H), 4.31–4.15 (m, 2H), 4.01 (m, 2H), 3.95–3.78 (m, 3H), 3.59 (dt, $J = 10.4, 4.8$ Hz, 1H), 2.34 (m, 2H). ^{13}C -NMR (100 MHz, $CDCl_3$) δ : 166.4, 166.3, 166.0, 165.9, 165.7, 165.5, 165.2, 164.9, 133.6, 133.5, 133.4, 133.3, 133.2, 133.0, 130.2–129.8, 129.6, 129.5, 129.4, 129.3, 129.1, 129.0, 128.9, 128.6–128.4, 100.8, 99.4, 98.1, 76.4, 73.3, 72.4, 70.5, 70.3, 69.9, 69.7–69.6, 69.4, 68.2, 67.8, 67.0, 66.6, 63.9–63.8, 62.6, 62.3, 50.4 (C-2). ESI-MS m/z calcd for $C_{103}H_{89}N_3O_{32}$: 1879.5; found: 962.5 $[M + Na]^+$. ESI-HRMS m/z calcd for $C_{103}H_{89}N_3O_{32}Na$ $[M + Na]^+$: 1902.5321; found: 1902.5307.

α Man₉ protected (11). To a solution of acceptor **10** (110 mg, 0.06 mmol), donor **8** (228 mg, 0.09 mmol) and 4 Å molecular sieves in anh. CH_2Cl_2 (4.1 mL) were added and the mixture was stirred at -20 °C for 30 min. Then, NIS (42 mg, 0.09 mmol) and TfOH (3.1 μ L, 17.55 μ mol) were added and the reaction was stirred at -20 °C for 15 min. The reaction was quenched with $NaHCO_3$ sat. aq. soln. The reaction mixture was filtered through Celite and washed several times with CH_2Cl_2 . The organic layer was washed with $Na_2S_2O_3$ sat. aq. soln. and then dried over anh. $MgSO_4$, filtered and concentrated under vacuum. The crude was purified by column chromatography on silica gel ($EtOAc/n$ -hexane 4 : 5 \rightarrow 1 : 1) to give the nonasaccharide **11** (185 mg, 0.04 mmol, 73%) as a white solid. $[\alpha]_D = -19$ (c 1.00, $CHCl_3$). 1H -NMR (400 MHz, $CDCl_3$) δ : 8.30–7.67 (m, 52H, HBz), 7.56–7.18 (m, 80H, HBz), 6.99 (m, 2H, HBz), 6.81 (t, $J = 7.5$, 1H, HBz), 6.22–5.80 (m, 16H), 5.55 (s, 1H, H-1), 5.46 (m, 2H, 2H-1), 5.32 (s, 1H, H-1), 5.18 (s, 1H, H-1), 5.05 (m, 2H, 2H-1), 4.74–4.02 (m, 36H, H-1, H-1), 3.84–3.72 (m, 2H), 3.49–3.32 (m, 3H). ^{13}C -NMR (100 MHz, $CDCl_3$) δ : 166.5, 166.4, 166.4, 166.3, 166.1, 166.0, 165.8, 165.6, 165.6, 165.4, 165.4, 165.3, 165.0, 165.0, 164.8, 164.8, 164.6, 133.5, 133.4, 133.4, 133.2, 133.2, 133.1, 130.2–129.5, 129.5, 129.5, 129.3, 129.2, 129.2, 129.0, 129.0, 128.9, 128.9, 128.7–128.3, 101.1, 100.9, 100.3, 99.9, 99.5, 98.5, 97.7, 97.5, 78.2, 77.9, 75.6, 72.4, 72.2, 71.9, 71.1, 70.8, 70.5, 70.1–69.4, 68.9, 68.1, 68.0, 67.7, 67.3–67.0, 66.6, 66.4, 66.1, 63.7, 63.5, 63.1, 62.6, 62.4, 50.4. ESI-MS m/z calcd for $C_{245}H_{203}N_3O_{73}$: 4354.2; found: 2200.1

$[M + 2Na]^{2+}$ and 1473.5 $[M + 3Na]^{3+}$. ESI-HRMS m/z calcd for $C_{245}H_{203}N_3O_{73}Na$ $[M + 2Na]^{2+}$: 2189.1115; found: 2189.1088.

α Man₉ (12). To a solution of compound **11** (166 mg, 0.04 mmol) in MeOH/toluene (4 : 1, 1.7 mL), NaOMe (11 mg, 0.20 mmol) and a NaOH 2 M solution (0.7 mL) were added and the reaction was stirred at 50 °C for 5 h. After neutralization with Amberlite IR-120H⁺, the solution was filtered and concentrated. The crude was purified by size-exclusion chromatography (Sephadex G-25, $H_2O/MeOH$ 9/1), giving Man₉ (**12**) (55 mg, 0.04 mmol, 93%) as a white amorphous solid. 1H -NMR (400 MHz, D_2O) δ : 5.40 (s, 1H, H-1), 5.33 (s, 1H, H-1), 5.30 (s, 1H, H-1), 5.15 (s, 1H, H-1), 5.08–5.01 (m, 3H, 3 \times H-1), 4.88 (s, 1H, H-1), under D_2O (1H, H-1), 4.19–3.60 (m, 58H). ^{13}C -NMR (100 MHz, D_2O) δ : 102.2, 100.8, 100.6, 100.0, 99.5, 98.0, 78.8–78.5, 73.2, 73.2, 72.7, 71.1, 71.1, 70.3–70.0, 69.6, 66.9, 66.8, 66.6, 65.7, 65.5, 65.2, 61.2–61.0, 50.2. ESI-MS m/z calcd for $C_{56}H_{95}N_3O_{46}$: 1545.5; found: 1568.2 $[M + Na]^+$, 795.5 $[M + 2Na]^{2+}$.

D3- α Man₉-Cl (15). Nonasaccharide **12** (22.0 mg, 14.23 μ mol) and [2-[2-(2-chloroethoxy)ethoxy]ethoxy]ethoxymethyl trikis(2-propynylloxymethyl)-methane (**14**) (1.6 mg, 3.59 μ mol) were dissolved in $H_2O/DMSO$ (1 : 1, 0.5 mL). Fresh solutions of $CuSO_4 \cdot 5H_2O$ (1.80 μ mol), tris[(1-benzyl-1*H*-1,2,3-triazol-4-yl)methyl]amine (TBTA) (3.59 μ mol) and sodium ascorbate (5.39 μ mol) were added to a sealed microwave vial. The solution was heated at 60 °C in a microwave oven for 30 min. A metal scavenger resin, QuadrasilMP, was added to the reaction solution and stirred for 20 min at rt. After that, the mixture was filtered and the resulting solution was purified by size-exclusion chromatography (Sephadex G-50, $H_2O/MeOH$ 9 : 1) yielding the glycocluster **15** (12 mg, 2.37 μ mol, 66%) as a white amorphous solid. 1H -NMR (400 MHz, D_2O) δ : 8.06 (s, 3H, $H_{triazole}$), 5.41 (br s, 3H, 3H-1), 5.37–5.27 (m, 6H, 6H-1), 5.15 (br s, 3H, 3H-1), 5.09–5.02 (m, 9H, 9H-1), under D_2O (6H, 6H-1), 4.67 (m, 3H, O CH_2CHHN), 4.57 (br s, 3H, $OCHHC_{triazole}$), 4.14–3.62 (m, 198H). ^{13}C -NMR (100 MHz, D_2O) δ : 144.2, 125.4, 102.2, 100.7, 99.8, 99.5, 98.0, 78.8–78.4, 73.3–73.2, 72.7, 72.0, 71.1–71.0, 70.8, 70.3, 70.0, 70.0, 69.7, 69.6, 69.5, 66.9, 66.8, 65.9, 65.8, 65.6, 65.4, 65.2, 64.8, 63.4, 62.5, 61.0, 50.1, 43.2. ESI-MS m/z calcd for $C_{191}H_{320}N_9O_{145}Cl$: 5080.7; found: 2562.9 $[M + 2Na]^{2+}$, 1715.5 $[M + 3Na]^{3+}$, and 1295.7 $[M + 4Na]^{4+}$.

D3- α Man₉-N₃ (17). To a solution of glycocluster **15** (9.0 mg, 1.77 μ mol) in DMF (1 mL) sodium azide (1.2 mg, 17.71 μ mol) was added. The mixture was stirred at 70 °C for 2 days. After that time, the solvent was evaporated under vacuum and the crude was purified using Amicon Ultra-15 centrifugal filters (MWCO 3 kDa), yielding the glycocluster **17** (9 mg, 1.77 μ mol, quant.) as a white amorphous solid. 1H -NMR (400 MHz, D_2O) δ : 8.06 (s, 3H, $H_{triazole}$), 5.41 (br s, 3H, 3H-1), 5.35–5.29 (m, 6H, 6H-1), 5.15 (br s, 3H, 3H-1), 5.07–5.03 (m, 9H, 9H-1), under D_2O (6H, 6H-1), 4.67 (m, 3H, $OCHHC_{triazole}$), 4.58 (br s, 3H, $OCHHC_{triazole}$), 4.14–3.60 (m, 198H). ^{13}C -NMR (100 MHz, D_2O) δ : 144.2, 125.5, 102.3, 100.7, 99.8, 99.5, 98.0, 78.8–78.4, 73.3–73.2, 72.0, 71.1–71.0, 70.3–69.9, 69.7–69.5, 66.9–66.8, 65.9, 65.6, 65.4, 65.2, 63.4, 62.5, 61.1–61.0, 50.1, 50.1, 44.6.



ESI-MS m/z calcd for $C_{191}H_{320}N_{12}O_{145}Cl$: 5087.2; found: 2567.0 $[M + 2Na]^{2+}$, 1781.1 $[M + 3Na]^{3+}$, and 1294.6 $[M + 4Na]^{4+}$.

D3- β Man₉-Cl (16). Nonasaccharide **13** (22.0 mg, 14.23 μ mol) and **14** (1.6 mg, 3.59 μ mol) were dissolved in H₂O/DMSO (1 : 1, 0.5 mL). Fresh solutions of CuSO₄·5H₂O (1.80 μ mol), TBTA (3.59 μ mol) and sodium ascorbate (5.39 μ mol) were added to a sealed microwave vial. The solution was heated at 60 °C in a microwave oven for 30 min. A metal scavenger resin, QuadrasilMP, was added to the reaction solution and stirred for 20 min at rt. After that, the mixture was filtered and the resulting solution was purified by size-exclusion chromatography (Sephadex G-50, H₂O/MeOH 9 : 1) yielding the glyco-cluster **16** (14.8 mg, 2.91 μ mol, 81%) as a white amorphous solid. ¹H-NMR (400 MHz, D₂O) δ : 8.05 (s, 3H, 3H_{triazole}), 5.42 (br s, 1H, 3H-1), 5.34–5.28 (m, 6H, 6H-1), 5.15 (br s, 3H, 3H-1), 5.09–5.03 (m, 9H, 9H-1), 4.87 (br s, 3H, 3H-1), 4.67 (m, 3H, OCH₂CHN), 4.61–4.56 (br s, 6H, H-1, OCHHC_{triazole}), 4.18–3.56 (m, 198H). ¹³C-NMR (100 MHz, D₂O) δ : 144.1, 125.5, 102.3, 102.2, 100.8, 100.7, 100.6, 99.9, 99.5, 98.0, 80.9, 78.9–78.5, 74.0, 73.3–73.2, 72.7, 71.1, 70.8, 70.3–69.9, 69.7–69.5, 68.3–67.7, 66.9–66.8, 65.6–65.2, 63.5, 62.5, 61.1–61.0, 50.3, 44.7, 43.2. ESI-MS m/z calcd for $C_{191}H_{320}N_9O_{145}Cl$: 5080.7; found: 2561.1 $[M + 2Na]^{2+}$, 1715.2 $[M + 3Na]^{3+}$, and 1293.5 $[M + 4Na]^{4+}$.

D3- β Man₉-N₃ (18). To a solution of glycocluster **16** (20.0 mg, 3.94 μ mol) in DMF (1 mL) sodium azide (2.6 mg, 39.36 μ mol) was added. The mixture was stirred at 70 °C for 2 days. After that time, the solvent was evaporated under vacuum and the crude was purified using Amicon Ultra-15 centrifugal filters (MWCO 3 kDa), yielding the glycocluster **18** (20 mg, 3.94 μ mol, quant.) as a white amorphous solid. ¹H-NMR (400 MHz, D₂O) δ : 8.05 (s, 3H, 3H_{triazole}), 5.41 (br s, 1H, 3H-1), 5.34–5.29 (m, 6H, 6H-1), 5.15 (br s, 3H, 3H-1), 5.07–5.03 (m, 9H, 9H-1), under D₂O (3H, 3H-1), 4.68 (m, 3H, OCHHC_{triazole}), 4.60–4.55 (m, 6H, H-1, OCH₂CHN), 4.12–3.46 (m, 198H). ¹³C-NMR (100 MHz, D₂O) δ : 144.1, 125.5, 102.3, 102.2, 100.8, 100.7, 100.6, 99.9, 99.5, 98.0, 80.9, 78.9–78.4, 74.0, 73.3–73.2, 72.7, 72.0, 71.1, 70.3–69.9, 69.7–69.5, 69.2, 68.3, 67.7, 66.9–66.8, 65.6–65.2, 63.5, 62.4, 61.1–61.0, 50.2, 50.1, 44.7. ESI-MS m/z calcd for $C_{191}H_{320}N_{12}O_{145}$: 5087.2; found: 2566.5 $[M + 2Na]^{2+}$, 1717.9 $[M + 3Na]^{3+}$, and 1293.9 $[M + 4Na]^{4+}$.

α Man₉-fluorescein (20). To a solution of the FAM-alkyne 6-isomer **19** (2.67 mg, 6.48 μ mol) and the epitope α Man₉ **12** (5 mg, 3.24 μ mol) in DMSO (300 μ L), another solution of CuBr (3.24 μ mol) and TBTA (6.48 μ mol) in the same solvent (150 μ L) was added. The reaction mixture was stirred on at rt. Thereafter, a metal scavenger resin, QuadrasilMP, was added to the reaction solution and stirred for 20 min at rt. After that, the mixture was filtered and the resulting solution was purified by size-exclusion chromatography (Sephadex LH-20, H₂O/MeOH 9 : 1), yielding the fluorescent probe **20** (6.3 mg, quant.) as an orange amorphous solid. ¹H-NMR (400 MHz, D₂O) δ : 8.22–7.87 (m, 3H), 7.68 (br s, 1H), 7.30–7.07 (m, 2H), 6.87–6.54 (m, 4H), 5.40–5.24 (m, 3H, 3H-1), 5.11 (br s, 1H, 1H-1), 5.04 (br s, 3H, 3H-1), under D₂O (2H, 2H-1), 4.64 (m, 2H), 4.12–3.57 (m, 58H). ¹³C-NMR (100 MHz, selected data obtained from

HSQC, D₂O) δ : 131.2, 128.2, 128.2, 127.9, 121.7, 103.0, 102.2, 100.8, 100.6, 100.0, 99.5, 98.0, 78.8–78.5, 73.2, 73.2, 72.7, 71.1, 71.1, 70.3–70.0, 69.6, 66.9, 66.8, 66.6, 65.7, 65.5, 65.2, 61.2–61.0, 50.2. ESI-MS: m/z calcd for $C_{80}H_{110}N_4O_{52}$: 1958.6, found: 978.1 $[M - 2H]^{2-}$.

β Man₉-fluorescein (21). To a solution of the FAM-alkyne 6-isomer **19** (6.6 mg, 10.36 μ mol) and the epitope β -Man₉ **13** (8 mg, 5.18 μ mol) in DMSO (300 μ L), another solution of CuBr (5.18 μ mol) and TBTA (10.36 μ mol) in the same solvent (150 μ L) was added. The reaction mixture was stirred on at rt. Thereafter, a metal scavenger resin, QuadrasilMP, was added to the reaction solution and stirred for 20 min at rt. After that, the mixture was filtered and the resulting solution was purified by size-exclusion chromatography (Sephadex LH-20, H₂O/MeOH 9 : 1), yielding the fluorescent probe **21** (10.1 mg, quant.) as an orange amorphous solid. ¹H-NMR (400 MHz, D₂O) δ : 8.05–7.98 (m, 2H), 7.94 (m, 1H), 7.57 (br s, 1H), 7.15–7.07 (m, 2H), 6.69–6.63 (m, 4H), 5.37 (br s, 1H, H-1), 5.31–5.26 (m, 2H, 2H-1), 5.10 (br s, 1H, H-1), 5.06–5.00 (m, 4H, 4H-1), 4.65 (m, 2H, CH₂C_{triazole}), 4.59 (m, 2H, CH₂CH₂N), 4.49 (br s, 1H, H-1), 4.21 (m, 1H), 4.12–4.05 (m, 6H), 4.03–3.92 (m, 6H), 3.90–3.60 (m, 42H), 3.41 (m, 1H). ¹³C-NMR (100 MHz, selected data obtained from HSQC, D₂O) δ : 131.0, 128.2, 128.2, 127.9, 121.7, 103.0, 102.3, 102.1, 100.7, 100.5, 99.9, 99.5, 97.9, 81.0, 78.9, 78.6, 78.5, 78.4, 74.0, 73.2–73.1, 72.6, 71.1, 70.3–69.9, 69.4, 68.4, 66.9–66.8, 65.6–65.4, 61.1–60.9, 50.3. ESI-MS: m/z calcd for $C_{80}H_{110}N_4O_{52}$: 1958.6, found: 1982.3 $[M - H]^-$, and 1002.1 $[M - 2H]^{2-}$.

D3- α Man₉-fluorescein (22). To a solution of the FAM-alkyne 6-isomer **19** (1.3 mg, 3.14 μ mol) and the trivalent glycocluster of α Man₉ **17** (8 mg, 1.57 μ mol) in DMSO (400 μ L), another solution of CuBr (1.57 μ mol) and TBTA (3.14 μ mol) in the same solvent (200 μ L) was added. The reaction mixture was stirred on at rt. Thereafter, a metal scavenger resin, QuadrasilMP, was added to the reaction solution and stirred for 20 min at rt. After that, the mixture was filtered and the resulting solution was purified by size-exclusion chromatography (Sephadex LH-20, H₂O/MeOH 9 : 1), yielding the fluorescent probe **22** (8.7 mg, quant.) as an orange amorphous solid. ¹H-NMR (400 MHz, D₂O) δ : 8.06–7.78 (m, 6H), 7.57 (br s, 1H), 7.07–6.88 (m, 2H), 6.62–6.42 (m, 4H), 5.35–5.16 (m, 9H, 9H-1), 5.06 (br s, 3H, 3H-1), 4.96 (br s, 9H, 9H-1), under D₂O (6H, 6H-1), 4.55–4.29 (m, 6H), 4.07–3.11 (m, 213H). ¹³C-NMR (100 MHz, selected data obtained from HSQC, D₂O) δ : 126.8, 122.9, 102.9, 102.3, 100.7, 99.8, 99.5, 98.0, 78.8–78.4, 73.3–73.2, 72.0, 71.1–71.0, 70.3–69.9, 69.7–69.5, 66.9–66.8, 65.9, 65.6, 65.4, 65.2, 63.4, 62.5, 61.1–61.0, 50.1. ESI-MS: m/z calcd for $C_{215}H_{335}N_{13}O_{151}$: 5514.9, found: 2749.4 $[M - 2H]^{2-}$, and 1833.1 $[M - 3H]^{3-}$.

D3- β Man₉-fluorescein (23). To a solution of the FAM-alkyne 6-isomer **19** (2.3 mg, 5.5 μ mol) and the trivalent glycoclusters of β Man₉ **18** (14 mg, 2.75 μ mol) in DMSO (180 μ L), another solution of CuBr (2.75 μ mol) and TBTA (5.5 μ mol) in the same solvent (90 μ L) was added. The reaction mixture was stirred on at rt. Thereafter, a metal scavenger resin, QuadrasilMP, was added to the reaction solution and stirred for 20 min at rt.



After that, the mixture was filtered and the resulting solution was purified by size-exclusion chromatography (Sephadex LH-20, H₂O/MeOH 9:1), yielding the fluorescent probe 23 (15.0 mg, quant.) as an orange amorphous solid. ¹H-NMR (400 MHz, D₂O) δ : 8.02–7.80 (m, 6H), 7.54 (br s, 1H), 7.01–6.87 (m, 2H), 6.58–6.46 (m, 4H), 5.40–5.30 (m, 6H, 6H-1), 5.21 (m, 6H, 6H-1), 5.06 (br s, 3H, 3H-1), 4.96 (m, 9H, 9H-1), 4.53–4.41 (m, 12H, 3H-1), 4.29 (br s, 6H), 4.07–3.05 (m, 204H). ¹³C-NMR (100 MHz, selected data obtained from HSQC, D₂O) δ : 131.0, 129.6, 128.5, 124.0, 122.4, 103.8, 102.3, 102.2, 100.8, 100.7, 100.6, 99.9, 99.5, 98.0, 80.9, 78.9–78.4, 74.0, 73.3–73.2, 72.7, 72.0, 71.1, 70.3–69.9, 69.7–69.5, 69.2, 68.3, 67.7, 66.9–66.8, 65.6–65.2, 63.5, 62.4, 61.1–61.0, 50.1 (OCH₂CH₂N). ESI-MS: m/z calcd for C₂₁₅H₃₃₅N₁₃O₁₅₁: 5514.9, found: 2748.5 [M – 2H]²⁻, and 1830.8 [M – 3H]³⁻.

DC-SIGN ECD expression

DC-SIGN ECD expression and production was performed as previously described. Briefly, the DC-SIGN ECD protein is expressed as inclusion bodies in *E. coli*, refolded and purified using a mannose affinity chromatography step followed by a size exclusion.^{28,29} Both steps ensure respectively the selection of correctly folded and oligomerized protein.

DC-SIGN binding assays by fluorescence polarization

The fluorescence polarization measurements were performed in 384-well microplates (black polystyrene, non-treated, Corning), using a TRIAD multimode microplate reader (from Dynex) with excitation and emission wavelengths of 485 and 535 nm, respectively. Fluorescent compounds 20–23 and DC-SIGN-ECD were dissolved in Tris buffer (25 mM, pH 8, 150 mM NaCl, 4 mM CaCl₂). 15 μ L of a 20 nM fluorescent ligand solution were transferred to each well. Then, 15 μ L of protein solutions, with concentrations ranging from 75 nM to 28 μ M, were added to the microplate wells. Therefore, the total sample volume in each well was 30 μ L. The microplate was shaken in the dark for 5 min, before reading. Blank wells contained 15 μ L of the DC-SIGN-ECD solution and 15 μ L of Tris buffer, and their measurements were subtracted from all values. All experiments on samples were performed in replicates of three.

Wells containing 15 μ L of the 20 nM fluorescence compound solution and 15 μ L of Tris buffer afforded the background polarization of the fluorescent molecule, in the absence of protein. This value was subtracted from the polarization values of all the samples, giving the increment in the fluorescence polarization (ΔP). The average ΔP values of three replicate wells were plotted against the concentration of DC-SIGN-ECD, and the resulting curve was fitted to the equation for a one-site binding model: $y = \Delta P_{\max}x/[K_D + x]$ where ΔP_{\max} is the maximal value of ΔP and K_D is the dissociation constant of the interaction.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by MINECO (CTQ2017-86265-P, PGC2018-099497-B-100 and IJCI-2015-23272), and ISCIII RETICS ARADyAL (RD16/0006/0011). Grants were co-funded by the European Regional Development Fund (ERDF). A. S.-H. and N. d. l. C thank MINECO for their Juan de la Cierva-Incorporacion and FPI contracts, respectively. This work used the Multistep Protein Purification Platform for human CLR production of the Grenoble Instruct-ERIC Center (ISBG; UMS 3518 CNRS-CEA-UGA-EMBL) within the Grenoble Partnership for Structural Biology (PSB), supported by FRISBI (ANR-10-INBS-05-02) and GRAL, financed within the University Grenoble Alpes graduate school (Ecoles Universitaires de Recherche) CBH-EUR-GS (ANR-17-EURE-0003). F. F. also acknowledge the French Agence Nationale de la Recherche (ANR) PIA for Glyco@Alps (ANR-15-IDEX-02). We acknowledge Dr Michael P. O'Hagan for the revision of the English version of the manuscript.

References

- 1 A. Varki, *Glycobiology*, 2017, **27**, 3–49.
- 2 H. Kaltner, J. Abad-Rodríguez, A. P. Corfield, J. Kopitz and H.-J. Gabius, *Biochem. J.*, 2019, **476**, 2623–2655.
- 3 J. J. Lundquist and E. J. Toone, *Chem. Rev.*, 2002, **102**, 555–578.
- 4 N. Jayaraman, *Chem. Soc. Rev.*, 2009, **38**, 3463–3483.
- 5 V. Wittmann, *Curr. Opin. Chem. Biol.*, 2013, **7**, 982–989.
- 6 B. Lepenies, J. Lee and S. Sonkaria, *Adv. Drug Delivery Rev.*, 2013, **65**, 1271–1281.
- 7 Y. van Kooyk and T. B. H. Geijtenbeek, *Nat. Rev. Immunol.*, 2003, **3**, 697–709.
- 8 L. Mathys and J. Balzarini, *PLoS One*, 2015, **10**, e0130621, DOI: 10.1371/journal.pone.0130621.
- 9 V. Y. Dudkin, M. Orlova, X. Geng, M. Mandal, W. C. Olson and S. J. Danishefsky, *J. Am. Chem. Soc.*, 2004, **126**, 9560–9562.
- 10 L. X. Wang, J. Ni, S. Singh and H. Li, *Chem. Biol.*, 2004, **11**, 127–134.
- 11 S.-K. Wang, P.-H. Liang, R. D. Astronomo, T.-L. Hsu, S.-L. Hsieh, D. R. Burton and C.-H. Wong, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 3690–3695.
- 12 F. Chiodo, P. M. Enríquez-Navas, J. Angulo, M. Marradi and S. Penadés, *Carbohydr. Res.*, 2015, **405**, 102–109.
- 13 H.-C. Wen, C.-H. Lin, J.-S. Huang, C.-L. Tsai, T.-F. Chen and S.-K. Wang, *Chem. Commun.*, 2019, **55**, 9124–9127.
- 14 H. Feinberg, R. Castelli, K. Drickamer, P. H. Seeberger and W. I. Weis, *J. Biol. Chem.*, 2007, **282**, 4202–4209.
- 15 J. Ramos-Soriano, J. J. Reina, B. M. Illescas, N. de la Cruz, L. Rodríguez-Pérez, F. Lasala, J. Rojo, R. Delgado and N. Martín, *J. Am. Chem. Soc.*, 2019, **141**, 15403–15412.
- 16 J. Ramos-Soriano, M. C. de la Fuente, N. de la Cruz, R. C. Figueiredo, J. Rojo and J. J. Reina, *Org. Biomol. Chem.*, 2017, **15**, 8877–8882.



- 17 J. J. Reina and J. Rojo, *Tetrahedron Lett.*, 2006, **47**, 2475–2578.
- 18 R. Ribeiro-Viana, M. Sánchez-Navarro, J. Luczkowiak, J. R. Koeppe, R. Delgado, J. Rojo and B. G. Davis, *Nat. Commun.*, 2012, **3**, 1303.
- 19 H. C. Kolb, M. G. Finn and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2001, **40**, 2004–2021.
- 20 C. W. Tornøe, C. Christensen and M. Meldal, *J. Org. Chem.*, 2002, **67**, 3057–3062.
- 21 V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, *Angew. Chem., Int. Ed.*, 2002, **41**, 2596–2599.
- 22 W. A. Lea and A. Simeonov, *Expert Opin. Drug Discovery*, 2011, **6**, 17–32.
- 23 K. Kakehi, Y. Oda and M. Kinoshita, *Anal. Biochem.*, 2001, **297**, 111–116.
- 24 P. Sörme, B. Kahl-Knutsson, M. Huflejt, U. J. Nilsson and H. Leffler, *Anal. Biochem.*, 2004, **334**, 36–47.
- 25 S. Maza, M. Mar Kayser, G. Macchione, J. Lopez-Prados, J. Angulo, J. L. de Paz and P. M. Nieto, *Org. Biomol. Chem.*, 2013, **11**, 3510–3525.
- 26 C. Solera, G. Macchione, S. Maza, M. M. Kayser, F. Corzana, J. L. de Paz and P. M. Nieto, *Chem. – Eur. J.*, 2016, **22**, 2356–2369.
- 27 I. Joachim, S. Rikker, D. Hauck, D. Ponader, S. Boden, R. Sommer, L. Hartmann and A. Titz, *Org. Biomol. Chem.*, 2016, **14**, 7933–7948.
- 28 J. J. Reina, I. Díaz, P. M. Nieto, N. E. Campillo, J. A. Páez, G. Tabarani, F. Fieschi and J. Rojo, *Org. Biomol. Chem.*, 2008, **6**, 2743–2754.
- 29 G. Tabarani, M. Thépaut, D. Stroebel, C. Ebel, C. Vivès, P. Vachette, D. Durand and F. Fieschi, *J. Biol. Chem.*, 2009, **284**, 21229–21240.

