


RESEARCH ARTICLE

View Article Online
View Journal | View IssueCite this: *Org. Chem. Front.*, 2020, 7, 3633Received 11th August 2020,
Accepted 26th September 2020

DOI: 10.1039/d0qo00967a

rsc.li/frontiers-organic

C–O/C–S difunctionalized benzene derivatives via multicomponent coupling of tetraynes†


Liangliang Yao, Qiong Hu, Yu Lei, Li Bao and Yimin Hu *

C–O/C–S difunctionalization of fused highly substituted benzene derivatives was conducted *via* the multicomponent coupling reaction of tetraynes, sulfoxides, and cyclopropanones. This reaction is associated with several bond cleavage and formation reactions in one pot, and also features exquisite regioselectivity and excellent yields. Preliminary mechanistic studies reveal that the reaction possibly proceeds in a sequential [2 + 2] cycloaddition, ring-opening of the cyclopropanone, protonation and dealkylation pathway.

Carbon–heteroatom bond formation is one of the most important tools used in pharmaceutical chemistry, medicinal chemistry, and materials sciences.¹ One-step synthesis of C_{sp²}–O and C_{sp²}–S bonds is an important and challenging task for chemists. Xiao and Chen disclosed the first case of aryne insertion into the S=O bond of sulfoxide for the direct synthesis of 1,2-O,S-disubstituted arenes.² Wang's group developed a synthesis technique for *ortho*-aryloxydiaryl sulfides in the absence of an external electrophile.³ Li and co-workers reported tandem aryne S=O bond insertion/C–H functionalization, featuring arene 1,2,3-trisubstitution from a relatively simple aryl allyl sulfoxide.⁴ Peng and co-workers synthesized *o*-aryloxytriaryl sulfonium salts instead of thioethers *via* aryne insertion into diaryl sulfoxides.⁵ Studer and his group designed a new protocol for the direct synthesis of *o*-arylsulfinyl aryl vinyl ethers by using aryne and aryl vinyl sulfoxides *via* aryne insertion into the S–O σ -bond and concomitant stereospecific S–O vinyl migration.⁶ Gogoi's group designed an interesting three component reaction for 2-[(*o*-methylthio)aryloxy] substituted dialkyl maleate derivatives by using aryne, dimethyl sulfoxide and activated alkynes.⁷ Many *o*-oxygen substituted aryl sulfides have been synthesized *via* aryne insertion into the S=O bond of sulfoxides. Two or more substituted *o*-alkylthio aryl esters are the important structural units in biologically active molecules and medicines, such as cox-2 inhibitors,⁸ boldine,⁹ and antimetabolites¹⁰ (Scheme 1a). Therefore, an efficient strategy

is desirable to be developed for the C–O/C–S difunctionalization of fused highly substituted benzene derivatives.

Demonstrated by Hoye's group in 2012,¹¹ the hexadehydro-Diels–Alder (HDDA) reaction, a new strategy to produce benzyne intermediates, can assemble three consecutive functional groups on a benzene ring in a “one-pot” process, and is compatible with various reagents and functional groups.¹² These benzyne intermediates are suitable for versatile transformations, thereby greatly expanding the fields of aryne chemistry and drug synthesis.¹³ As part of our ongoing research on the efficient construction of polysubstituted arenes *via* HDDA-derived benzyne chemistry,¹⁴ we envisioned that different tetrayne substrates can be used as HDDA-derived benzyne precursors to form the C–O/C–S difunctionalized and concomitant *o*-alkylthio aryl ester core structure benzene derivatives by inserting benzyne into the S=O bond and opening the ring of cyclopropanone (Scheme 1b). To the best of our knowledge, this hypothesis has not been verified by using benzyne intermediates. Thus, exploring the resultant intermediate could provide a novel platform to incorporate sulfide and ester groups into valuable aromatic compounds.



Scheme 1 Bioactivity with the *ortho*-alkylthio aryl ester core structure and our work.

Laboratory of Functional Molecular Solids, Ministry of Education, Anhui Key Laboratory of Molecular-Based Materials, School of Chemistry and Materials Science, Anhui Normal University, Wuhu, Anhui 241000, China.
E-mail: yiminhu@ahnu.edu.cn

† Electronic supplementary information (ESI) available: Experimental procedures and characterization of all new compounds. CCDC 1571269–1571270. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d0qo00967a



A multicomponent coupling reaction of HDDA-derived benzynes, cyclopropenones, and sulfoxides was performed to produce various C–O/C–S difunctionalized benzene derivatives in good to excellent yields. The process of introducing sulfide and ester moieties into an aromatic ring can be achieved in the absence of catalysts, bases, or oxidants and therefore could provide a broad range of construction methods for fully functionalized arenes. Synthesis chemists can achieve the rapid synthesis of drug molecules and other high-value compounds.

Results and discussion


Inspired by the preliminary result, tetrayne **1a** was used as a HDDA-derived benzyne precursor and 2,3-diphenylcycloprop-2-enone **2a** in DMSO at 90 °C to optimize the reaction conditions. Screening results revealed that the optimal yield could be obtained at 120 °C in DMSO at the 1 : 1.1 ratio of tetrayne to cyclopropenone in an 8 h reaction. It was verified that DMSO played a crucial role in this multicomponent coupling reaction. The optimal conditions were as follows: 1.0 equiv. of tetraynes reacted with 1.1 equiv. of cyclopropenones in DMSO at 120 °C for 8 h.

The scope of tetrayne substrates was investigated under the optimized conditions. As depicted in Table 1, various products (**4a–4q**) were readily isolated with good to excellent yields (ranging from 69% to 89%) from the reactions of tetraynes **1** with 2,3-diphenylcycloprop-2-enone **2a** in DMSO. The effect of different tetraynes on the product yield was also examined. When tolerated, tetrayne substrates bearing different esters (OMe, OEt, and OⁱPr) gradually increased in yields (**4a** (82%), **4d** (85%), and **4j** (89%)). By contrast, the substituent groups in the aryl rings of tetraynes bearing electron-donating groups, including *p*-Me, *m*-Me, and *p*-Et (**4k**, **4l**, and **4m**), exhibited higher yields than the electron-withdrawing groups, such as *p*-F and *p*-Cl (**4n** and **4o**) possibly because the electron-withdrawing groups decreased the reactivity. Compound **4j** had the highest isolated yield (89%) among the examined products.

Furthermore, 2,3-di-*p*-tolylcycloprop-2-enone and 2,3-bis(4-fluorophenyl)cycloprop-2-enone can participate in the reaction to afford good yields (**4p** (85%) and **4q** (74%)). The structures of **4a** and **4m** were confirmed through X-ray diffraction.¹⁵


As shown in Table 2, 68%–82% yields were obtained from the reactions of tetraynes **1** with 2,3-diphenylcycloprop-2-enone **2a** in *n*-butyl sulfoxide. The carboxylates of the tetraynes with OⁱPr, OEt and OMe provided minimal impact, providing **4r** (79%), **4s** (80%), and **4w** (82%), respectively. The yields of benzene ring-containing tetraynes with alkyl groups as electron donors were slightly higher than those of tetraynes directly bearing halogen atoms as electron-withdrawing groups. For example, the yields of **4t** and **4u** were 74% and 76%, respectively, and that of **4v** was only 68%. These results revealed the potential of the multicomponent coupling reaction of current HDDA-derived benzynes with cyclopropenones and sulfoxides for the synthesis of C–O/C–S difunctionalized benzene derivatives.

Table 1 Preparation of fully functionalized *o*-(methylthio)phenyl acrylates^{a,b}




^a Conditions: tetraynes **1** (0.5 mmol), cyclopropenones **2** (1.1 equiv), H₂O (1.0 equiv), DMSO (2 mL), stirred at 120 °C for 8 h. ^b Yield of the isolated product after flash column chromatography.




Table 2 Preparation of polysubstituted *o*-(butylthio)phenyl acrylates^{a,b}

^a Conditions: tetraynes **1** (0.5 mmol), cyclopropanones **2** (1.1 equiv), H₂O (1.0 equiv), and *n*-butyl sulfoxide (2 mL), stirred at 120 °C for 8 h.
^b Yield of the isolated product after flash column chromatography.

Labeling and control experiments were performed to gain insight into the reaction mechanism (Scheme 2). DMSO-*d*₆ was used instead of DMSO for the synthesis of **4d** under the



Scheme 2 Deuterium-labeling and control experiments.



Scheme 3 Proposed reaction mechanism.

optimized reaction conditions (Scheme 2, eqn (1)). The deuterium-labeled product **d-4d** was isolated in 83% yield. ¹H NMR analysis confirmed the complete deuterium incorporation at the “CH₃” moiety, indicating that DMSO serves as the “SMe” source. Another independent experiment was performed in which D₂O was used instead of H₂O with **1d**, and **2a** in DMSO-*d*₆ to further investigate the reaction mechanism (Scheme 2, eqn (2)). The expected product **d-4d'** was formed in 84% yield with complete deuterium incorporation at the vinylic position. These results suggest that our reaction proceeded via the [2 + 2] cycloaddition of a HDDA-derived benzyne with DMSO solvent, and that H₂O served as the proton source (details are in the Experimental section and ESI†).

A possible mechanism underlying the formation of C–O/C–S difunctionalized benzene derivatives was proposed on the basis of the above results and previous literature^{2,7,16} (Scheme 3). First, the HDDA-derived benzyne intermediate **A** underwent [2 + 2] cycloaddition insertion into the S=O bond of dimethyl sulfoxide to form a four-membered ring intermediate **B**. Owing to the ring strain, this intermediate exhibited ring opening to afford an *o*-quinone intermediate **C**. Intermediate **C** is in resonance with zwitterion **D**. Nucleophilic attack of cyclopropanone **2a** on **D** leads to zwitterion **E**, which was further protonated in the presence of residual water to form intermediate **F**. Finally, the resultant hydroxide anion from H₂O abstracted the methyl cation of intermediate **F** to generate the product **d-4d'**.

Conclusions

In conclusion, we have developed a multicomponent coupling reaction of HDDA-derived benzynes, cyclopropanones and sulfoxides for the direct synthesis of C–O/C–S difunctionalized benzene derivatives. This reaction strategy has been accomplished through several bond cleavage and formation reactions in a one-pot procedure. Compared with previous work, this strategy introduces sulfide and ester moieties at the adjacent positions of an arene ring with the C–O and C–S bonds and derivatives thereof. The reaction does not require a transition



metal catalyst but exhibits excellent regioselectivity and produces all types of highly substituted fused benzene derivatives under mild conditions with good to excellent yields. Further work on the applications and scope extension of this protocol is on-going in our laboratory.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors thank the National Natural Science Foundation of China (22071001, 21572002) and the Department of Human Resources of Anhui Province for financial support.

Notes and references

- (a) *Catalyzed Carbon-Heteroatom Bond Formation*, ed. A. K. Yudin, Wiley-VCH, Weinheim, 2011; (b) J. Yin, in *Applications of Transition Metal Catalysis in Drug Discovery and Development*, ed. M. L. Crawley and B. M. Trost, John Wiley and Sons, Hoboken, 2012, pp. 97–163; (c) A. Bhunia, S. R. Yetra and A. T. Biju, Recent advances in transition-metal-free carbon-carbon and carbon-heteroatom bond-forming reactions using arynes, *Chem. Soc. Rev.*, 2012, **41**, 3140–3152; (d) X. Li, X. Li, M. N. Banis, B. Wang, A. Lushington, X. Cui, R. Li, T.-K. Sham and X. Sun, Tailoring interactions of carbon and sulfur in Li-S battery cathodes: significant effects of carbon-heteroatom bonds, *J. Mater. Chem. A*, 2014, **2**, 12866–12872; (e) H.-T. Kim, H. Shin, I.-Y. Jeon, M. Yousaf, J. Baik, H.-W. Cheong, N. Park, J.-B. Baek and T.-H. Kwon, Carbon-heteroatom bond formation by an ultrasonic chemical reaction for energy storage systems, *Adv. Mater.*, 2017, **29**, 1702747.
- F.-L. Liu, J.-R. Chen, Y.-Q. Zou, Q. Wei and W.-J. Xiao, Three-component coupling reaction triggered by insertion of arynes into the S=O bond of DMSO, *Org. Lett.*, 2014, **16**, 3768–3771.
- H.-Y. Li, L.-J. Xing, M.-M. Lou, H. Wang, R.-H. Liu and B. Wang, Reaction of arynes with sulfoxides, *Org. Lett.*, 2015, **17**, 1098–1101.
- Y. Li, D. Qiu, R. Gu, J. Wang, J. Shi and Y. Li, Aryne 1,2,3-trifunctionalization with aryl allyl sulfoxides, *J. Am. Chem. Soc.*, 2016, **138**, 10814–10817.
- X. Li, Y. Sun, X. Huang, L. Zhang, L. Kong and B. Peng, Synthesis of *o*-aryloxy triarylsulfonium salts via aryne insertion into diaryl sulfoxides, *Org. Lett.*, 2017, **19**, 838–841.
- Y. Li and A. Studer, Reaction of arynes with vinyl sulfoxides: Highly stereospecific synthesis of ortho-sulfinylaryl vinyl ethers, *Org. Lett.*, 2017, **19**, 666–669.
- H. Hazarika, K. Neog, A. Sharma, B. Das and P. Gogoi, Three-component coupling reactions of aryne, DMSO, and activated alkyne: stereoselective synthesis of 2-[(*o*-methylthio)aryloxy]-substituted dialkyl maleates, *J. Org. Chem.*, 2019, **84**, 5846–5854.
- (a) A. S. Kalgutkar, B. C. Crews, S. W. Rowlinson, C. Garner, K. Seibert and L. J. Marnett, Aspirin-like molecules that covalently inactivate cyclooxygenase-2, *Science*, 1998, **280**, 1268–1270; (b) A. S. Kalgutkar, K. R. Kozak, B. C. Crews, G. P. Hochgesang Jr. and L. J. Marnett, Covalent modification of cyclooxygenase-2 (COX-2) by 2-acetoxyphenyl alkyl sulfides, a new class of selective COX-2 inactivators, *J. Med. Chem.*, 1998, **41**, 4800–4818; (c) C. F. Pereira, J. T. M. L. Paridaen, K. Rutten, M. C. D. G. Huigen, M. van de Bovenkamp, J. Middel, N. Beerens, B. Berkhout, R. Schuurman, L. J. Marnett, J. Verhoef and H. S. L. M. Nottet, Aspirin-like molecules that inhibit human immunodeficiency virus 1 replication, *Antiviral Res.*, 2003, **58**, 253–263; (d) H. K. Jain, V. K. Mouryab and R. K. Agrawal, Inhibitory mode of 2-acetoxyphenyl alkyl sulfides against COX-1 and COX-2: QSAR analyses, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 5280–5284.
- F. A. Thometa, P. Pinyolb, J. Villenab, L. J. Espinozaa and P. G. Reveco, Cytotoxic thiocarbamate derivatives of boldine, *Nat. Prod. Commun.*, 2010, **5**, 1587–1590.
- T. H. Porter, C. M. Bowman and K. Folkers, Antimetabolites of coenzyme Q. 16. New alkylmercaptoquinones having antimalarial curative activity, *J. Med. Chem.*, 1973, **16**, 115–118.
- T. R. Hoye, B. Baire, D. Niu, P. H. Willoughby and B. P. Woods, The hexadecylo-Diels-Alder reaction, *Nature*, 2012, **490**, 208–212.
- For selected reviews and examples, see: (a) T. R. Hoye, B. Baire and T. Wang, Tactics for probing aryne reactivity: mechanistic studies of silicon-oxygen bond cleavage during the trapping of (HDDA-generated) benzyne by silyl ethers, *Chem. Sci.*, 2014, **5**, 545–550; (b) R. Karmakar and D. Lee, Reactions of arynes promoted by silver ions, *Chem. Soc. Rev.*, 2016, **45**, 4459–4470; (c) X. Xiao, T. Wang, F. Xu and T. R. Hoye, CuI-mediated bromoalkynylation and hydroalkynylation reactions of unsymmetrical benzyne: Complementary modes of addition, *Angew. Chem., Int. Ed.*, 2018, **57**, 16564–16568; (d) S. Gupta, Y. Lin, Y. Xia, D. J. Winka and D. Lee, Alder-ene reactions driven by high steric strain and bond angle distortion to form benzocyclobutenes, *Chem. Sci.*, 2019, **10**, 2212–2217; (e) S. Arora, J. Zhang, V. Pogula and T. R. Hoye, Reactions of thermally generated benzyne with six-membered N-heteroaromatics: pathway and product diversity, *Chem. Sci.*, 2019, **10**, 9069–9076; (f) S. Ghorai, Y. Lin, Y. Xia, D. J. Wink and D. Lee, Silver-catalyzed annulation of arynes with nitriles for synthesis of structurally diverse quinazolines, *Org. Lett.*, 2020, **22**, 626–630.
- For selected examples, see: (a) D. Niu and T. R. Hoye, The aromatic ene reaction, *Nat. Chem.*, 2013, **6**, 34–40; (b) T. Wang and T. R. Hoye, Hexadecylo-Diels-Alder (HDDA)-enabled carbazolyne chemistry: Single step, de novo construction of the pyranocarbazole core of alkaloids of the *Murraya koenigii* (Curry tree) family, *J. Am. Chem.*



- Soc.*, 2016, **138**, 13870–13873; (c) R. Karmakar and D. Lee, Total synthesis of Selaginpulvilin C and D relying on in situ formation of arynes and their hydrogenation, *Org. Lett.*, 2016, **18**, 6105–6107; (d) S. P. Ross and T. R. Hoye, Reactions of hexadehydro-Diels–Alder benzynes with structurally complex multifunctional natural products, *Nat. Chem.*, 2017, **9**, 523–530; (e) X. Xiao and T. R. Hoye, The domino hexadehydro-Diels–Alder reaction transforms polyynes to benzynes to naphthynes to anthracynes to tetraynes (and beyond?), *Nat. Chem.*, 2018, **10**, 838–844; (f) X. Xiao and T. R. Hoye, One-pot, three-aryne cascade strategy for naphthalene formation from 1,3-diyne and 1,2-benzdiyne equivalents, *J. Am. Chem. Soc.*, 2019, **141**, 9813–9818.
- 14 (a) Y. Hu, J. Ma, L. Li, Q. Hu, S. Lv, B. Liu and S. Wang, Fused multifunctionalized dibenzoselenophenes from tetraynes, *Chem. Commun.*, 2017, **53**, 1542–1545; (b) B. Liu, C. Mao, Q. Hu, L. Yao and Y. Hu, Direct methylation and carbonylation of in situ generated arynes via HDDA-Wittig coupling, *Org. Chem. Front.*, 2019, **6**, 2788–2791; (c) B. Liu, Q. Hu, F. Yang, X. Zheng and Y. Hu, Fused multifunctionalized bridge aromatic hydrocarbons from in situ-generated arynes and anthracene derivatives, *Chin. Chem. Lett.*, 2020, **31**, 1305–1308; (d) X. Zheng, B. Liu, F. Yang, Q. Hu, L. Yao and Y. Hu, Access to benzoxazepines and fully substituted indoles via HDDA Coupling, *Org. Lett.*, 2020, **22**, 956–959.
- 15 CCDC 1571270 (**4a**) and 1571269 (**4m**) contain the supplementary crystallographic data for this paper.†
- 16 H. Hazarika and P. Gogoi, Direct synthesis of ortho-methylthio allyl and vinyl ethers via three component reaction of aryne, activated alkene and DMSO, *Org. Biomol. Chem.*, 2020, **18**, 2727–2738.

