


Cite this: *RSC Adv.*, 2020, 10, 1995

Cu-metal organic frameworks (Cu-MOF) as an environment-friendly and economical catalyst for one pot synthesis of tacrine derivatives†

Hoda Mollabagher,^a Salman Taheri,^a Mohammad majid Mojtahedi^a and SeyedAmirhossein Seyedmousavi^b

The present work describes the catalytic activity of Cu-MOF for the one-pot synthesis of tacrine derivatives via a four-component reaction of 2-hydroxynaphthalene-1,4-dione, aldehydes, malononitrile and cycloketones in the presence of AlCl_3 . The structure of the synthesized compound is confirmed by ^1H NMR, ^{13}C NMR, IR, and MASS. The catalyst prepared under pressure is characterized by powder X-ray diffraction and SEM. The noteworthy advantages of this procedure include its broad substrate scope, high yields up to 93%, atom economy, using readily available starting materials and a powerful recyclable nano catalyst. Additionally, there is no need to use column chromatography for purifying products so, it has the potential for large-scale applications in pharmaceutical industries. Another advantage of this method is the ability to recycle the catalyst up to 3 times and reuse it.

Received 3rd December 2019

Accepted 2nd January 2020


DOI: 10.1039/c9ra10111j

rsc.li/rsc-advances

1. Introduction

According to the World Health Organization, over 30 million people worldwide suffer from Alzheimer's disease (AD), and research has shown that this number is rising substantially.¹ AD is actually a type of brain dysfunction that gradually degrades the mental abilities of the patient and leads to memory impairment and dementia. Acetylcholine is one of the substances that

transmits the message between the two nerve cells and then decomposes with cholinesterase. In AD reducing the acetylcholine disrupts the transmission of messages between the two nerve cells.² Therefore, many studies have been done to strengthen the cholinesterase system includes acetylcholine release agents such as 4-aminopyridine (1), and anticholinesterase drugs, such as physostigmine (2) and acetylcholinesterase inhibitors like tacrine (3) and 4-aminoquinoline (4).



^aChemistry and Chemical Engineering Research Center of Iran, PO Box 14115-186, Tehran, Iran. E-mail: taheri@ccerci.ac.ir

^bProcess Engineering Department, Faculty of Chemical Engineering, Tarbiat Modares University, Tehran, Iran

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra10111j

Among the listed compounds, tacrine is one of the drugs that boost acetylcholine by inhibiting cholinesterase enzymes³ and is known as an important reference with amazing pharmacological properties. It was first identified in 1993 as a drug for the treatment of AD.⁴ Studying of tacrine analogues is still of interest to researchers investigating AD. As a result, several




Heterogeneous catalysts have also many other advantages such as easily isolation from the mixture of the reaction, be recyclable and less contamination in final product.¹⁸ Metal organic frameworks (MOFs) attracted the attention of the scientific community around 1990 (ref. 19) and then it has been effectively used as heterogeneous catalysis improving efficiency

Our work has presented a novel procedure to the synthesis of tacrine derivatives, starting from aldehyde, malononitrile, 2-hydroxynaphthalene-1,4-dione and cycloketone derivative as one-pot reaction without the requirement for separating the pyranic intermediate. Applying the Cu-MOF as heterogeneous catalyst assistances the formation of pyranic intermediates, followed by the addition of aluminum chloride to the Friedländer quinoline reaction, without interfering with the two catalysts in the reaction process. The presence of Cu active sites in Cu-MOF has made it a suitable candidate for the synthesis of pyrene compounds. We describe convenient synthesis methods for tacrine derivatives from the simple starting material.

2.1. Chemicals and apparatus

All starting materials were purchased from Sigma-Aldrich and Merck companies and used without additional purification. Reactions were monitored by aluminum TLC plate, silica gel coated with fluorescent indicator F₂₅₄. Melting points were measured using a Buchi B-545 apparatus through the capillary tube method and reported without any correction. IR spectra were recorded from KBr disks using FT-IR Bruker Vector-22 infrared spectrometer in the range of 400–4000 cm⁻¹. The ¹H NMR and ¹³C NMR were run on an FT-NMR Bruker



This journal is © The Royal Society of Chemistry 2020

UltraShield™ (500 MHz) in DMSO-*d*₆ and CDCl₃ as a solvent and the chemical shifts are expressed as δ units with tetramethylsilane (TMS) as internal standard. Mass spectra were obtained on an Agilent Technologies apparatus at ionization potential of 70 eV. The concentration of copper in the prepared catalysts was determined using inductively coupled plasma optical emission spectroscopy by SPECTRO ARCOS ICP-OES spectrometer.

2.2. General procedure for the preparation of compounds 6a–k

A mixture of 2-hydroxy-1,4-naphthoquinone **1** (0.174 g, 1.0 mmol), aromatic aldehyde **2** (1.0 mmol), malononitrile **3** (0.06 g, 1.0 mmol), and catalytically amount of Cu-MOF (0.02 g) in dry 1,2-dichloroethane (DCE) (5.0 mL) was stirred under reflux condition until completion reaction and disappearance of the starting materials on TLC (EtOAc : *n*-hexane 1 : 1). Then, AlCl₃ (0.1995 g, 1.5 mmol) and cycloketone **5** (1.5 mmol) were added into the mixture of the reaction. The mixture was continuously stirred under argon atmosphere under reflux. After completion, the mixture was centrifuged to isolated catalyst and then the solvent was evaporated under reduced pressure, H₂O and THF (1 : 1) were added and the mixture was basified with 10% sodium hydroxide solution to pH = 8–9. After stirring for 30 min, the mixture was diluted using CH₂Cl₂ and the organic layer was dried over MgSO₄. In the end, the crude is crystallization by ethyl acetate to obtain the pure product.

2.3. Preparation of nano catalyst Cu-MOF

The nano catalyst was synthesized with modification according to the method described in previous articles.²³ 0.24 gram (1 mmol) of copper(II) nitrate trihydrate and 0.24 gram (1 mmol) 1,4-benzenedioic acid gradually dissolve in 20 mL *N,N*-dimethylformamide (DMF). The resulting mixture was then transferred into a 25 mL autoclave reactor and placed under 5 barr at 80 °C for 4 hours. After cooling to the room temperature, the precipitate was centrifuged and Cu-MOF turquoise powder was obtained. To activate the catalyst and remove the DMF from the cavities, Cu-MOF was put into Soxhlet extractor for 1 day with dichloromethane as solvent.

The point to consider is that the use of subterranean radiation, as well as reacting in a reactor under pressure, can reduce the synthesis time of MOF from 24 hours to 4 hours, as well as lower the temperature from 100 °C to 80 °C than that of other papers. The ICP-OES showed the Cu content in Cu-BDC structure was about 38.95%.

3. Results and discussion

3.1. Structural analysis of Cu-MOF nano catalyst

FT-IR is a main technique for determination of the functional groups in the structure of compounds. Therefore, the study of FT-IR spectra is an accepted method among chemists to identify materials.^{24,25} In Fig. 1, the FT-IR spectrum of Cu-MOF shows clearly the removing DMF from Cu-MOF by washing with dichloromethane. Comparison of the diagrams (A) and (B)

denotes the deletion of the peak associated with the DMF methyl groups in region 2936. Remove peak in region 3486, indicates a withdrawal of water and carboxylic acid present in the cavity, after the activation of the catalyst.

According to the reported literature,²⁶ the Cu-MOF structure releases DMF molecules during the activation process, and has active copper sites (Scheme 2).

The powder XRD pattern of the Cu-MOF compound clearly indicates the presence of copper in its structure and the sharp peak appearing in less than 15 indicates a highly crystalline structure (Fig. 2). According to the XRD spectra, the average size of the crystals (*D*) was 52 nm using the Scherrer equation (1) by replacing λ as the wavelength of the X-ray beam, $\beta_{1/2}$ as the line broadening at half the maximum intensity and θ as the Bragg's angle.

$$D = \frac{0.9\lambda}{\beta_{1/2} \cos \theta} \quad (1)$$

SEM techniques can be used to obtain the surface, shape, and appearance of particles. As it clear in Fig. 3, the cubic Cu-MOF structure that was synthesis under pressure is well-defined and it has good agreement with other published articles.²⁷

3.2. Catalytic behaviors of nano-Cu-MOF for the synthesis tacrine derivatives

The multicomponent reaction of benzaldehyde, malononitrile, 2-hydroxy-1,4-naphthoquinone, and cyclohexanone under different conditions were investigated to determine the optimal conditions for tacrine synthesis. Various solvents such as ethanol, H₂O, dichloromethane, tetrachloroethylene (C₂Cl₄) and 1,2-dichloroethane (EtCl₂) were screened in the presence of the various catalyst under an air atmosphere. As mentioned before the most of the reported papers on synthesis of tacrine derivatives from the pyranic intermediates, include two or more stages which cause unfavorable higher reaction time and the undesirable less efficiency,²⁸ it is also clearly visible in Table 1 (Entry 1 and 2).

In order to optimize the reaction conditions and increase the yield, one step reaction was investigated by using triethylamine and aluminum chloride which does not go further because of undesired interaction between reagents (Entry 3). To overcome this problem, we have used a heterogeneous catalyst, MOF nano catalyst, in dichloroethane as solvent. Our research has revealed that 4 mg catalyst per mmol of reagent has the best efficiency. The optimum condition was achieved by the domino reaction benzaldehyde, malononitrile, 2-hydroxy-1,4-naphthoquinone and cyclohexanone in presence of 4 mg per mmol of Cu-MOF and 1.2 mmol of aluminum chloride.

The elemental analyses, ¹H, ¹³C NMR, FT-IR spectra and MS of the product clearly indicated the formation of **6a**. The IR spectra of **6a** exhibited ν_{\max} at 1631 cm^{−1} for the carbonyl function and 3372 cm^{−1} and 3452 cm^{−1} for NH₂. The ¹H NMR spectra of **6a** showed the proton attached to spiro carbon and proton attached to the nitrogen were resonated as a singlet at






Fig. 1 FT-IR spectra of Cu-MOF (A) after activation, (B) before activation that present omitting DMF.

δ 5.26 and 4.20 respectively. The ^1H -decoupled ^{13}C NMR spectrum of **6a** showed 26 distinct resonances in agreement with the proposed structure **6a**, carbonyl groups carbon displayed ^{13}C resonance signal at δ 182.2 and 176.7 ppm, respectively.

After the optimization of reaction conditions, it was generalized by different types of aromatic aldehydes and cycloketones. The domino one-pot synthesis of tacrine derivatives was achieved from the reaction of various aromatic aldehydes, malononitrile, and 2-hydroxy-1,4-naphthoquinone in the presence of Cu-MOF nano catalyst followed by the increase of various ketones and aluminum chloride in short reaction time with high yield as shown in Table 2.

One advantage of using heterogeneous catalysts compared to homogeneous catalysts is their ability to be recycled and reused in chemical reactions, which has attracted particular attention.

3.3. Reusability of nano catalyst

The ability to the recovery of Cu-MOF catalyst was measured by using after 5 times and no significant change in its catalytic properties was observed. As shown in Fig. 4, the efficiency of the fresh catalyst and the catalyst used have no main different.


Also, FT-IR spectra of the fresh and recycle Cu-MOF catalyst are in good agreement with each other (Fig. S2†).

No leached copper was found in the solution. However, the effect of the homogeneous catalyst was investigated by species leached from the catalyst after hot filtration, and as shown in Fig. S1,† it presented negligible activity in the filtered solution.

Representative spectral data

14-Amino-13-phenyl-2,3,4,13-tetrahydro-1H-benzo[6,7]chromeno[2,3-b]quinoline-7,12-dione (6a). Yield 93 (%); yellow solid; mp = 290–293 dec °C; IR (KBr): 3464, 3376 (NH_2), 1631, 1595 ($\text{C}=\text{O}$) cm^{-1} . ^1H NMR (500 MHz, CDCl_3) δ = 8.44–7.83 (m, 2H), 7.78–7.57 (m, 2H), 7.54–7.15 (m, 5H), 5.24 (s, 1H), 4.19 (s, 2H, NH_2), 2.87–2.80 (m, 2H), 2.45–2.23 (m, 2H), 1.92–1.77 (m, 4H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ = 182.2, 176.7, 154.0, 152.8, 149.7, 149.2, 140.6, 133.0, 132.4, 130.7, 129.8 (2C), 127.9 (2C), 126.6, 125.4, 125.2, 121.5, 113.6, 97.6, 54.2, 34.9, 31.4, 21.9, 21.5, 21.2 ppm; MS (70 eV) m/z 408 [M^+], 331, 105, 77; anal. calcd for $\text{C}_{26}\text{H}_{20}\text{N}_2\text{O}_3$: C, 76.46; H, 4.94; N, 6.86. Found: C, 76.50; H, 4.83; N, 6.91.

14-Amino-13-(4-chlorophenyl)-2,3,4,13-tetrahydro-1H-benzo[6,7]chromeno[2,3-b]quinoline-7,12-dione (6b). Yield 91 (%);



Scheme 2 Synthesis and activation of Cu-MOF.






Fig. 2 pXRD pattern of Cu-MOF.

yellow solid; mp = 290–291 dec °C; IR (KBr): 3463, and 3376 (NH₂), 1631, 1595 (C=O) cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ = 8.15 (dd, *J* = 4.0, 5.0 Hz, 1H), 7.98 (dd, *J* = 4.0, 5.0 Hz, 1H), 7.70 (d, *J* = 5.0 Hz, 1H), 7.69 (d, *J* = 5.0 Hz, 1H), 7.38 (d, *J* = 8.5 Hz, 2H), 7.23 (d, *J* = 8.5 Hz, 2H), 5.27 (s, 1H), 4.31 (s, 2H, NH₂), 2.82–2.81 (m, 2H), 2.42–2.38 (m, 2H), 1.88–1.81 (m, 4H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ = 182.3, 176.8, 154.2, 152.7, 149.8, 149.4, 139.2, 133.2, 132.6, 132.5, 130.6, 129.8, 129.2, 128.1, 125.6, 125.3, 121.2, 113.8, 97.0, 34.3, 31.5, 22.0, 21.5, 21.2 ppm; MS (70 eV) *m/z* 442 [M⁺], 331, 105, 77; anal. calcd for C₂₆H₁₉ClN₂O₃: C, 70.51; H, 4.32; N, 6.33. Found: 70.43; H, 4.45; N, 6.38.

14-Amino-13-(3-methoxyphenyl)-2,3,4,13-tetrahydro-1H-benzo[6,7]chromeno[2,3-*b*]quinoline-7,12-dione (6c). Yield 89 (%); yellow solid; mp = 270–271 dec °C; IR (KBr): 3472, 3375 (NH₂), 1682, 1634, 1594 cm⁻¹. ¹H NMR (500 MHz, CDCl₃) δ = 8.17 (dd, *J* = 5.8, 3.3 Hz, 1H), 8.03 (dt, *J* = 7.3, 3.8 Hz, 1H), 7.70 (dt, *J* = 7.1, 3.6 Hz, 2H), 7.23 (t, *J* = 7.9 Hz, 2H), 7.07 (d, *J* = 7.6 Hz, 1H), 6.99 (t, *J* = 2.2 Hz, 1H), 6.76 (dd, *J* = 8.3, 2.6 Hz, 1H), 5.23 (s, 1H), 4.21 (s, 2H, NH₂), 3.76 (s, 3H, OMe), 2.84 (s, 2H), 2.49–2.20 (m, 2H), 1.86 (s, 4H); ¹³C NMR (125 MHz, CDCl₃) δ = 182.2, 161.2, 159.0, 154.2, 152.6, 149.8, 142.3, 133.0, 132.5, 130.7, 129.9, 128.9, 125.5, 125.3 (2C), 120.3 (2C), 114.0, 113.9, 113.66, 111.7, 54.2, 35.0, 31.4, 21.9, 21.4, 21.2 ppm; MS (70 eV) *m/z* 438 [M⁺], 331, 92, 77; anal. calcd for C₂₇H₂₂N₂O₄: C, 73.96; H, 5.06; N, 6.39. Found: C, 73.89; H, 5.15; N, 6.33.

14-Amino-13-(*p*-tolyl)-2,3,4,13-tetrahydro-1H-benzo[6,7]chromeno[2,3-*b*]quinoline-7,12-dione (6d). Yield 90 (%); yellow solid; mp = 242–245 dec °C; IR (KBr) = 3472, 3375 (NH₂), 1734, 1635, 1594 cm⁻¹. ¹H NMR (500 MHz, CDCl₃) δ = 8.39–7.94 (m, 2H), 7.81–7.54 (m, 2H), 7.41–7.30 (m, 2H), 7.11 (d, *J* = 7.9 Hz, 2H), 5.25 (s, 1H), 4.20 (s, 2H, NH₂), 2.97 (h, *J* = 9.2 Hz, 2H), 2.79–2.50 (m, 2H), 2.28 (s, 3H, Me), 2.21–2.09 (m, 2H), 2.07 (s, 1H), 1.28 (t, *J* = 7.2 Hz, 1H) ppm; ¹³C NMR (125 MHz, CDCl₃) δ = 182.4, 177.0, 162.0, 155.0, 149.2, 147.8, 137.9, 136.5, 133.0, 132.5, 130.7, 129.8, 128.6 (2C), 127.7 (2C), 125.5, 125.3, 121.6, 118.3, 98.2, 33.3, 33.2, 26.0, 21.2, 20.0, 13.1 ppm; MS (70 eV) *m/z* 442 [M⁺], 331, 105, 91; anal. calcd for C₂₇H₂₂N₂O₃: C, 76.76; H, 5.25; N, 6.63. Found: C, 76.70; H, 5.19; N, 6.70.

14-Amino-13-(thiophen-2-yl)-2,3,4,13-tetrahydro-1H-benzo[6,7]chromeno[2,3-*b*]quinoline-7,12-dione (6e). Yield 55 (%); brown solid; mp = 269–271 dec °C; IR (KBr) = 3472, 3378 (NH₂), 1734, 1678, 1635, 1594 cm⁻¹. ¹H NMR (500 MHz, DMSO-*d*₆) δ = 8.11–8.04 (m, 1H), 8.03–7.97 (m, 1H), 7.92–7.66 (m, 2H), 7.27 (d, *J* = 5 Hz, 1H), 7.0 (d, *J* = 3.5 Hz, 1H), 6.86 (dd, *J* = 5.1, 3.6 Hz, 1H), 6.03 (s, 2H, NH₂), 5.84 (s, 1H), 2.61 (d, *J* = 5.5 Hz, 2H), 2.46–2.13 (m, 2H), 1.34–0.97 (m, 2H) ppm; ¹³C NMR (125 MHz, DMSO-*d*₆) δ = 182.0, 177.3, 152.7, 152.4, 151.2, 149.7, 145.5, 134.0, 133.4, 130.6, 130.0, 126.0, 125.5, 125.5, 125.3, 124.5, 121.8, 113.2, 96.7, 31.4, 28.4, 22.5, 21.7, 21.4 ppm; MS (70 eV) *m/z* 414 [M⁺], 331, 105, 77; anal. calcd for C₂₄H₁₈N₂O₃S: C, 69.55; H, 4.38; N, 6.76; S, 7.74. Found: C, 69.50; H, 4.32; N, 6.84; S, 7.79.

13-Amino-12-(*p*-tolyl)-1,2,3,12-tetrahydrobenzo[6,7]chromeno[2,3-*b*]cyclopenta[*e*]pyridine-6,11-dione (6f). Yield 92 (%); yellow solid; mp = 300–302 dec °C; IR (KBr) = 3445, 3355 (NH₂), 1731, 1644, 1582 cm⁻¹. ¹H NMR (500 MHz, CDCl₃) δ = 8.27–7.92 (m, 2H), 7.76–7.59 (m, 2H), 7.34 (d, *J* = 8.0 Hz, 2H), 7.11 (d, *J* = 7.8 Hz, 2H), 5.25 (s, 1H), 4.18 (s, 2H, NH₂), 2.96 (hept, *J* = 8.2,





Fig. 3 SEM image of cubic structure of Cu-MOF.



Table 1 One-pot domino reaction between 2-hydroxy-1,4-naphthoquinone (1 mmol), benzaldehyde (1 mmol), malononitrile (1 mmol) and cyclohexanone (1 mmol) under various conditions

Entry	Solvent (conditions) ^a	Catalyst	Time (h)	Yield ^b (%)
1	EtOH (reflux)	NEt ₃ (5 mol%)/AlCl ₃ (2 step)	12	67
2	H ₂ O (reflux)	NEt ₃ (5 mol%) AlCl ₃ (2 step)	24	Trace
3	EtOH (reflux)	NEt ₃ (5 mol%) AlCl ₃	24	—
4	EtOH (reflux)	Ni-MOF (3 mg)/AlCl ₃	12	30
5	EtOH (reflux)	Al-MOF	12	20
6	EtOH (reflux)	Cu-MOF (3 mg)/AlCl ₃	24	50
7	EtCl ₂ (reflux)	Cu-MOF (3 mg)/AlCl ₃	5	88
8	CH ₂ Cl ₂ (reflux)	Cu-MOF (3 mg)/AlCl ₃	5	40
9	C ₂ Cl ₄ (reflux)	Cu-MOF (3 mg)/AlCl ₃	24	—
10	CH ₂ Cl ₂ (US) ^c	Cu-MOF (3 mg)/AlCl ₃	12	Trace
11	EtCl ₂ (reflux)	Cu-MOF (3 mg)/AlCl ₃	30	30
12	—(120 °C)	Cu-MOF (3 mg)/AlCl ₃	20	Trace
13	EtOH (reflux)	CuAl-MOF	12	40
14	EtCl ₂ (reflux)	Cu-MOF (4 mg)/AlCl ₃	5	93
15	EtCl ₂ (reflux)	Cu-MOF (5 mg)/AlCl ₃	5	93
16	EtCl ₂ (reflux)	Cu-MOF (2 mg)/AlCl ₃	5	73

^a Reaction conditions: benzaldehyde (1 mmol), malononitrile (1 mmol), 2-hydroxy-1,4-naphthoquinone (1 mmol) and cyclohexanone (1.2 mmol).^b Isolated yields. ^c Ultrasound at 80% amplitude and was sonicated for 10 min.

7.3 Hz, 2H), 2.81–2.48 (m, 2H), 2.28 (s, 3H), 2.07 (s, 1H), 1.28 (t, $J = 7.1$ Hz, 1H) ppm; ¹³C NMR (125 MHz, CDCl₃) $\delta = 182.4, 162.0, 155.0, 149.1, 147.8, 137.9, 136.4, 133.0, 132.5, 130.7, 129.8, 128.7, 127.7, 125.5, 125.2, 121.5, 118.2, 98.1, 34.5, 33.2, 26.1, 21.2, 20.0, 13.1$ ppm; MS (70 eV) m/z 408 [M⁺], 317, 91, 77; anal. calcd for C₂₆H₂₀N₂O₃: C, 76.46; H, 4.94; N, 6.86. Found: C, 76.50; H, 4.82; N, 6.92.

13-Amino-14-(*p*-tolyl)-8,9,10,11,12,14-hexahydrobenzo[6,7]chromeno[2,3-*b*]cyclohepta[*e*]pyridine-5,15-dione (6g). Yield 50 (%); brownish solid; mp = 263–266 dec °C; IR (KBr) = 3465, 3378 (NH₂), 1641, 1599 cm⁻¹. ¹H NMR (500 MHz, CDCl₃) $\delta = 8.14$ (s, 1H), 7.99 (s, 1H), 7.67 (s, 2H), 7.33 (d, $J = 7.9$ Hz, 2H), 7.0 (d, $J = 8.5$ Hz, 2H), 5.21 (s, 1H), 4.25 (s, 2H, NH₂), 2.81 (s, 2H), 2.30–2.12 (m, 4H), 1.9–1.4 (m, 5H), 1.26 (d, $J = 7.8$ Hz, 2H) ppm; ¹³C NMR (125 MHz, CDCl₃) $\delta = 182.4, 176.8, 161.6, 158.7, 153.8, 149.8, 137.8, 136.4, 133.0, 132.4, 130.7, 129.8, 128.6$ (2C), 127.7 (2C), 125.5, 125.2, 121.6, 113.6, 97.7, 34.5, 31.4, 21.9, 21.5, 21.2, 20.0, 13.1 ppm; MS (70 eV) m/z 436 [M⁺], 421, 345, 91; anal. calcd for C₂₈H₂₄N₂O₃: C, 77.04; H, 5.54; N, 6.42. Found: C, 77.02; H, 5.59; N, 6.49.

14-Amino-13-(2-chlorophenyl)-2,3,4,13-tetrahydro-1H-benzo[6,7]chromeno[2,3-*b*]quinoline-7,12-dione (6h). Yield 70 (%); yellow solid; mp = 274–277 dec °C; IR (KBr) = 3451, 3383 (NH₂), 1636, 1597 cm⁻¹. ¹H NMR (500 MHz, CDCl₃) $\delta = 8.26$ –8.09 (m, 1H), 8.09–7.92 (m, 1H), 7.79–7.61 (m, 2H), 7.44–7.32 (m, 1H), 7.15 (tt, $J = 7.4, 5.4$ Hz, 2H), 5.63 (s, 1H), 4.54 (s, 2H, NH₂), 2.96–























2.61 (m, 2H), 2.45–2.14 (m, 2H), 1.88–1.65 (m, 4H) ppm; ¹³C NMR (125 MHz, CDCl₃) $\delta = 182.1, 176.8, 153.8, 152.4, 150.1, 149.8, 138.9, 133.2, 132.5, 130.4, 130.5, 130.2, 129.8, 128.5, 127.8, 127.0, 125.6, 125.3, 121.4, 113.3, 97.1, 31.7, 31.3, 22.0, 21.4, 21.2$ ppm; MS (70 eV) m/z 442 [M⁺], 407, 331, 105; anal. calcd for C₂₆H₁₉ClN₂O₃: C, 70.51; H, 4.32; N, 6.33. Found: C, 70.48; H, 4.31; N, 6.40.

14-Amino-13-phenyl-4,13-dihydro-1H,3H-benzo[6,7]chromeno[2,3-*b*]thiopyrano[3,4-*e*]pyridine-7,12-dione (6i). Yield 82 (%); yellow solid; mp = 270–273 dec °C; IR (KBr) = 3451, 3376 (NH₂), 1657, 1595 cm⁻¹. ¹H NMR (500 MHz, DMSO-*d*₆) $\delta = 8.07$ –8.00 (m, 1H), 7.93 (d, $J = 4.8$ Hz, 1H), 7.85–7.77 (m, 2H), 7.57–7.46 (m, 2H), 7.30–6.9 (m, 5H), 5.54 (s, 1H), 3.64–3.53 (m, 1H), 3.40 (d, $J = 16.2$ Hz, 1H), 3.00 (q, $J = 6.2$ Hz, 2H), 2.88 (d, $J = 6.1$ Hz, 2H) ppm; ¹³C NMR (125 MHz, DMSO-*d*₆) $\delta = 181.7, 176.1, 152.9, 150.7, 148.6, 148.3, 148.3, 140.8, 134.1, 133.7, 130.3, 129.9, 128.2$ (2C), 128.1, 127.8 (2C), 126.7, 125.5, 125.4, 122.7, 111.8, 32.6, 30.4, 23.0, 22.6 ppm; MS (70 eV) m/z 426 [M⁺], 349, 315, 303; anal. calcd for C₂₅H₁₈N₂O₃S: C, 70.41; H, 4.25; N, 6.57; S, 7.52. Found: C, 70.40; H, 4.23; N, 6.63; S, 7.61.

14-Amino-13-(4-nitrophenyl)-4,13-dihydro-1H,3H-benzo[6,7]chromeno[2,3-*b*]thiopyrano[3,4-*e*]pyridine-7,12-dione (6j). Yield 78 (%); yellow solid; mp = 284–286 dec °C; IR (KBr) = 3490, 3360 (NH₂), 1637, 1592 cm⁻¹. ¹H NMR (500 MHz, DMSO-*d*₆) $\delta = 8.10$ (d, $J = 8.3$ Hz, 2H), 8.08–8.02 (m, 1H), 7.96–7.90 (m, 1H), 7.86–7.79 (m, 2H), 7.77 (d, $J = 8.2$ Hz, 2H), 6.19 (s, 2H, NH₂), 5.69 (s,



Table 2 One-pot, four-component synthesis of new tacrine derivatives in the presence of Cu-MOF and AlCl_3 in 1,2-dichloroethane under reflux conditions

Entry	Aldehyde	Cycloketone	Yield (%)	Time (h)	Product	Mp ($^{\circ}\text{C}$)
1			95	5	6a	Dec: 290–293
2			91	3.25	6b	Dec: 290–291
3			89	3	6c	Dec: 270–271
4			90	3.35	6d	Dec: 242–245
5			55	6	6e	Dec: 269–271
6			92	2.5	6f	Dec: 300–302
7			50	4.5	6g	Dec: 263–266
8			70	3.5	6h	Dec: 274–277
9			82	4	6i	Dec: 270–273
10			78	5	6j	Dec: 284–286
11			90	4.5	6k	Dec: 250–253






Fig. 4 The recycling of Cu-MOF as catalysts using a model reaction of 2-hydroxynaphthalene-1,4-dione, benzaldehyde, malononitrile following by adding cyclohexanone and AlCl_3 .

1H), 3.57 (d, $J = 16.3$ Hz, 1H), 3.38 (d, $J = 16.0$ Hz, 1H), 2.94 (q, $J = 6.0, 5.4$ Hz, 2H), 2.85 (h, $J = 8.2, 7.3$ Hz, 2H) ppm; ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$) $\delta = 182.0, 177.0, 152.7, 152.0, 150.7, 150.1, 149.2, 145.7, 133.9, 133.4, 130.5, 130.0, 129.5$ (2C), 125.5, 125.2, 122.7 (2C), 121.0, 111.9, 96.8, 33.3, 32.9, 24.0, 22.9 ppm; MS (70 eV) m/z 471 $[\text{M}^+]$, 349, 315, 303; anal. calcd for $\text{C}_{25}\text{H}_{17}\text{N}_3\text{O}_5\text{S}$: C, 63.69; H, 3.63; N, 8.91; S, 6.80. Found: C, 63.60; H, 3.68; N, 8.88; S, 6.76.

14-Amino-13-(4-chlorophenyl)-4,13-dihydro-1H,3H-benzo[6,7]chromeno[2,3-b]thiopyrano[3,4-e]pyridine-7,12-dione (6k). Yield 90 (%); yellow solid; mp = 250–253 dec °C; IR (KBr) = 3450, 3345 (NH_2), 1665, 1595 cm^{-1} . ^1H NMR (500 MHz, $\text{DMSO}-d_6$) $\delta = 8.13$ –8.01 (m, 1H), 8.00–7.91 (m, 1H), 7.85 (t, $J = 4.6$ Hz, 2H), 7.55 (d, $J = 8.1$ Hz, 2H), 7.32 (d, $J = 8.1$ Hz, 2H), 7.09 (s, 2H, NH_2), 5.59 (s, 1H), 3.60 (d, $J = 16.3$ Hz, 1H), 3.40 (d, $J = 16.2$ Hz, 1H), 3.08–2.81 (m, 4H) ppm; ^{13}C NMR (125 MHz, CDCl_3) $\delta = 182.8, 176.5, 152.9, 150.7, 148.8, 139.7, 134.1, 133.6, 131.4, 130.5, 130.2$ (2C), 130.0, 127.7 (2C), 125.5, 125.4, 122.0, 112.0, 97.5, 32.1, 30.5, 23.1, 22.6 ppm; MS (70 eV) m/z 460 $[\text{M}^+]$, 349, 315, 303; anal. calcd for $\text{C}_{25}\text{H}_{17}\text{ClN}_2\text{O}_3\text{S}$: C, 65.15; H, 3.72; 6.08; S, 6.96. Found: C, 65.10; H, 3.83; 6.18; S, 6.90.

4. Conclusion

In conclusion, we have effectively developed and expanded simple and straightforward procedure for one-pot synthesis of tacrine derivatives. One-pot, domino four-component reaction of aldehydes, malononitrile, 2-hydroxy-1,4-naphthoquinone, and cycloketones was carried out in the presence of Cu-MOF and aluminum chloride. Without the requirement to isolate the pyranic intermediate and in a single container, the reaction was conducted under reflux to give the desired products in high yields. Prominent features of the current procedure include a step-economic increasing yield, economic feasibility, capability of recycling the catalyst and covering a broad substrate

scope. The heterogeneous nanocatalyst Cu-MOF presented excellent catalytic performance and can be recycled simply by centrifuging; it can also recover 5 times without considerable loss of efficiency.

We expect this method to be performed easily by pharmacists and chemists and biologically active tacrine derivatives that were synthesized over several stages can be easily synthesized without the need to separate intermediates and purify them.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We would like to thank the Research Council at CCERC Foundation for the financial support of this work. The authors of this article also express their gratitude and appreciation for the efforts made by Tarbiat Modares University in catalyst synthesis.

References

- 1 P. L. Valenzuela, J. S. Morales, C. Fiuza-Luces, P. de la Villa, A. Santos-Lozano and A. Lucia, *J. Am. Med. Dir. Assoc.*, 2019, **20**, 101–102.
- 2 S. Shimohama and J. Kawamata, in *Nicotinic Acetylcholine Receptor Signaling in Neuroprotection*, Springer, Singapore, 2018, pp. 137–158.
- 3 H. Cavdar, M. Senturk, M. Guney, S. Durdagi, G. Kayik, C. T. Supuran and D. Ekinici, *J. Enzyme Inhib. Med. Chem.*, 2019, **34**, 429–437.
- 4 M. Mehta, A. Adem and M. Sabbagh, *Int. J. Alzheimer's Dis.*, 2012, **2012**, 728983.



- 5 S. Ökten, M. Ekiz, Ü. M. Koçyiğit, A. Tutar, İ. Çelik, M. Akkurt, F. Gökalp, P. Taslimi and İ. Gülçin, *J. Mol. Struct.*, 2019, **1175**, 906–915.
- 6 M. Ekiz, A. Tutar and S. Ökten, *Tetrahedron*, 2016, **72**, 5323–5330.
- 7 A. Martínez-Grau and J. Marco, *Bioorg. Med. Chem. Lett.*, 1997, **7**, 3165–3170.
- 8 J. Li, L. Zhang, D. Shi, Q. Li, D. Wang, C. Wang, Q. Zhang, L. Zhang and Y. J. S. Fan, *Synlett*, 2008, 233–236.
- 9 M. T. McKenna, G. R. Proctor, L. C. Young and A. Harvey, *J. Med. Chem.*, 1997, **40**, 3516–3523.
- 10 M. A. Khalilzadeh, A. Hosseini and M. Tajbakhsh, *J. Heterocycl. Chem.*, 2007, **44**, 535–538.
- 11 G. Tai and W. Guo, *Ultrason. Sonochem.*, 2008, **15**, 350–356.
- 12 M. Eghtedari, Y. Sarrafi, H. Nadri, M. Mahdavi, A. Moradi, F. H. Moghadam, S. Emami, L. Firoozpour, A. Asadipour and O. Sabzevari, *Eur. J. Med. Chem.*, 2017, **128**, 237–246.
- 13 L. Pourabdi, M. Khoobi, H. Nadri, A. Moradi, F. H. Moghadam, S. Emami, M. M. Mojtahedi, I. Haririan, H. Forootanfar, A. Ameri, A. Foroumadi and A. Shafiee, *Eur. J. Med. Chem.*, 2016, **123**, 298–308.
- 14 T. Ahmadi, G. Mohammadi Ziarani, P. Gholamzadeh and H. Mollabagher, *Tetrahedron: Asymmetry*, 2017, **28**, 708–724.
- 15 S. Y. Afsar, G. M. Ziarani, H. Mollabagher, P. Gholamzadeh, A. Badiei and A. Soorki, *J. Iran. Chem. Soc.*, 2017, **14**, 577–583.
- 16 G. Mohammadi Ziarani, H. Mollabagher, N. Lashgari and A. Badiei, *Sci. Iran.*, 2018, **25**, 3295–3304.
- 17 G. M. Ziarani, H. Mollabagher, P. Gholamzadeh, A. Badiei and F. Yazdian, *Iran. J. Catal.*, 2018, **8**, 59–67.
- 18 G. Bosica and R. Zammit, *PeerJ*, 2018, **6**, e5065.
- 19 S. Subudhi, D. Rath and K. Parida, *Catal. Sci. Technol.*, 2018, **8**, 679–696.
- 20 J. Lee, O. K. Farha, J. Roberts, K. A. Scheidt, S. T. Nguyen and J. T. Hupp, *Chem. Soc. Rev.*, 2009, **38**, 1450–1459.
- 21 C. de los Ríos, J. L. Marco, M. a. D. C. Carreiras, P. M. Chinchón, A. G. García and M. Villarroya, *Bioorg. Med. Chem.*, 2002, **10**, 2077–2088.
- 22 N. P. Selvam, T. H. Babu and P. T. Perumal, *Tetrahedron*, 2009, **65**, 8524–8530.
- 23 R. Salama, S. Abd El-Hakam, S. Samra, S. El-dafrawy and A. Ahmed, *Int. J. Mod. Chem.*, 2018, **8**, 679–696.
- 24 G. Mohammadi Ziarani, H. Mollabagher, N. Lashgari and A. Badiei, *Sci. Iran.*, 2018, **25**, 3295–3304.
- 25 A. Yazdani-Elah-Abadi, R. Mohebat, M.-T. Maghsoodlou and R. J. P. A. C. Heydari, *Polycyclic Aromat. Compd.*, 2018, **38**, 92–101.
- 26 S. Rostamnia, H. Alamgholiloo and X. Liu, *J. Colloid Interface Sci.*, 2016, **469**, 310–317.
- 27 N. T. S. Phan, T. T. Nguyen, K. D. Nguyen and A. X. T. Vo, *Appl. Catal., A*, 2013, **464–465**, 128–135.
- 28 M. Khoobi, F. Ghanoni, H. Nadri, A. Moradi, M. P. Hamedani, F. H. Moghadam, S. Emami, M. Vosooghi, R. Zadnabad and A. Foroumadi, *Eur. J. Med. Chem.*, 2015, **89**, 296–303.

