


Cite this: *RSC Adv.*, 2020, **10**, 25475

Received 13th March 2020
Accepted 28th June 2020

DOI: 10.1039/d0ra02358b
rsc.li/rsc-advances

Silver-catalyzed regioselective deuteration of (hetero)arenes and α -deuteration of 2-alkyl azaarenes[†]

Baobiao Dong,^{ab} Xuefeng Cong^{id *b} and Na Hao^{*a}


A simple silver-catalyzed regioselective deuteration of (hetero)arenes and α -deuteration of 2-alkyl azaarenes has been described. This strategy provides an efficient and practical avenue to access various deuterated electron-rich arenes, azaarenes and α -deuterated 2-alkyl azaarenes with good to excellent deuterium incorporation utilizing D_2O as the source of deuterium atoms.

Deuterium-labelled organic compounds are of great interest and importance, as they are widely used as internal standards in analytical chemistry,¹ tools for elucidation of reaction mechanisms,² metabolic or pharmacokinetic probes,³ and as biologically active compounds and pharmaceuticals.⁴ Indeed, the first deuterated drug, deutetrabenazine, which is recognized as a different orphan drug for the treatment of chorea *versus* tetraabenazine, was approved by the US Food and Drug Administration in April 2017. As a result, the development of efficient and selective routes for the synthesis of deuterium labeled organic compounds is a subject of increasing interest. Among possible approaches to incorporate deuterium atoms into an organic molecule,⁵ the H/D exchange reaction, which enables the direct deuterium labelling of the desired target molecule without the need to prefunctionalize the starting materials, represents the most straightforward and atom-efficient method. In recent years, extensive studies on H/D exchange reactions of aromatic compounds have been reported, including acid/base promoted pH-dependent H/D exchange reactions of electronically activated aromatic compounds⁶ (Scheme 1a) and transition metal catalyzed site-selective H/D exchange reactions⁷ (Scheme 1b). Among these, transition-metal-catalyzed reactions have gained significant momentum because these catalysts can control the site selectivity of H/D exchange reactions. However, the required ligands, the introduction of directing groups and/or the complexity of catalyst synthesis may not be readily accessible or economically viable. Therefore, there remains room to develop a simple, cost efficient and universal deuteration strategy with broad substrate scope, especially in view of


the growing demand for deuterium-labelled compounds in synthetic chemistry and pharmaceutical industry.

In past decades, silver has demonstrated their high efficiency as special mild Lewis acids, becoming catalysts of first choice for many types of catalytic reactions generally performed under mild reaction conditions and through experimentally simple procedures.⁸ However, despite considerable progress, silver-catalyzed selective H/D exchange reaction has remained much less explored to date.⁹ Quite recently, Huang reported an efficient Ag_2CO_3 -catalyzed H/D exchange of five-membered heteroarenes using D_2O as deuterium source at ambient temperature, but external base such as 1 equiv. of K_2CO_3 and a phosphine ligand were still needed.^{9b} We here developed a practical and selective silver-catalyzed deuteration of (hetero)arenes and α -deuteration of 2-alkyl azaarenes utilizing D_2O as the source of deuterium atom under mild neutral conditions that does not require any other additives (Scheme 1c). This protocol represents a practical and efficient method of silver-catalyzed regioselective H/D exchange reaction under mild


a) Acid/base promoted deuteration of aromatic compounds (pH-dependence)



b) Transition metals catalyzed deuteration of aromatic compounds (C–H activation)



c) Silver-catalyzed regioselective deuteration of (hetero)arenes (This work)



Scheme 1 Selective H/D exchange reactions.

^aDepartment of Pharmaceutical Sciences, School of Pharmacy, Southwest Medical University, Luzhou 646000, China. E-mail: haona@swmu.edu.cn

^bJilin Province Key Laboratory of Organic Functional Molecular Design & Synthesis, Department of Chemistry, Northeast Normal University, Changchun 130024, China. E-mail: congxf010@nenu.edu.cn

[†] Electronic supplementary information (ESI) available: Experimental details, and characterization data. See DOI: 10.1039/d0ra02358b




neutral conditions, efficiently affording a wide range of deuterated electron-rich arenes, azaarenes and α -deuterated 2-alkyl azaarenes through experimentally simple procedures.

Initially, we began our studies by examining the model deuteration of 1,2,3,4-tetrahydroquinoline (**1a**) with D₂O using various commercially available silver salts as the catalyst. To our delight, 94% deuterium incorporation was achieved in the *ortho*- and *para*-positions by using 10 mol% AgOTf in CDCl₃ at 90 °C (Table 1, entry 1). Other silver salts such as Ag₂CO₃, Ag₃PO₄, AgBF₄, AgOAc and Ag₂O were also tested as catalysts, but gave low deuterium incorporation (entries 2–6). AgOTf showed higher activity probably because of the slow release of TfOH in the reaction.^{9a} Control experiment suggests that AgOTf is necessary in this reaction (Table 1, entry 7). We then tested various solvents for this deuteration, and found that THF and 1,4-dioxane can also give good results, but DCE, CH₃CN afforded a slightly low deuterium incorporation (entries 8–11). The reduction of the D₂O amount from 20 to 5 equiv. led to a decrease in deuterium incorporation from 94% to 62% (Table 1, entry 12). When the reaction temperature was decreased to 60 °C, the efficiency of deuterium incorporation dramatically decreased, only giving 58% deuterium incorporation in the *ortho*- and *para*-positions (entry 13). In addition, under standard reaction conditions, other Lewis acids such as Cu(OTf)₂, Zn(OTf)₂, ZnCl₂, FeCl₃, and La(OTf)₃ could also catalyze the H/D exchange, but giving a low deuterium incorporation, especially in the *para*-position.^{10,11}

With the optimized catalytic system in hand, we evaluated the scope and limitations of this selective H/D exchange reaction. We firstly examined the deuteration of various electronic-

Table 1 Optimization of conditions for the silver-catalyzed deuteration of **1a**^a




Entry	[Ag]	Solvent	Deuterium incorporation (D ¹ , D ²)
1	AgOTf	CDCl ₃	94%, 94%
2	Ag ₂ CO ₃	CDCl ₃	11%, 8%
3	Ag ₃ PO ₄	CDCl ₃	21%, 23%
4	AgBF ₄	CDCl ₃	91%, 58%
5	AgOAc	CDCl ₃	21%, 10%
6	Ag ₂ O	CDCl ₃	13%, 10%
7	—	CDCl ₃	<5%, <5%
8	AgOTf	DCE	74%, 81%
9	AgOTf	THF	94%, 93%
10	AgOTf	1,2-Dioxane	90%, 90%
11	AgOTf	CH ₃ CN	77%, 91%
12 ^c	AgOTf	CDCl ₃	62%, 62%
13 ^d	AgOTf	CDCl ₂	58%, 58%

^a Reaction conditions: **1a** (0.5 mmol), [Ag] (0.025 mmol), D₂O (10 mmol), solvent (1.0 mL), 90 °C for 18 h. ^b Deuterium incorporation was determined by ¹H NMR. ^c D₂O (2.5 mmol). ^d At 60 °C.

rich arenes, and the results are listed in Table 2. We were pleased to find out that highly selective H/D exchanges with good to excellent degrees of deuteration were achieved. A variety of substituted 1,2,3,4-tetrahydroquinolines were smoothly converted to the deuterated isotopologues with high levels of deuterium incorporation (**2a–2f**). The bromide and dimethyl-phenylsilyl substituents could be well tolerated in this catalytic system. The deuteration of indoline derivatives as well as *N*-methyl aniline derivatives could be achieved with good to excellent deuterium incorporation in the *ortho*- and *para*-positions (**2g–2l**). The 9,10-dihydroacridine **1m** was also deuterated to afford **2m** in 88% yield with excellent deuterium incorporation. Moreover, other electron-rich aromatic compounds such as 1,3,5-trimethoxybenzene (**1n**) and multi-substituted anisole (**1o**) were also suitable substrates for the selective deuteration affording **2n** and **2o** with good deuterium incorporation. The conversion of *N,N*-disubstituted aniline derivatives such as *N,N*-dimethylaniline under the standard conditions only led to quite poor D-incorporation (<20% D incorporation).

Subsequently, this catalytic system was applied to the deuteration of indole and azole derivatives. The deuteration of these compounds is of great interest and importance, as they

Table 2 Silver-catalyzed selective deuteration of various electron-rich arenes^a



^a Reaction conditions: **1** (0.5 mmol), AgOTf (0.025 mmol), D₂O (10 mmol), CDCl₃ (1.0 mL), 90 °C for 18 h. Isolated yield are given. ^b 1,4-dioxane (1.0 mL) was used.

Table 3 Silver-catalyzed selective deuteration of various azaarenes^a

(Aza)Ar-H 3	AgOTf (5 mol %), D ₂ O (20 equiv) CDCl ₃ , 90 °C, 18 h	(Aza)Ar-D 4
		4a , 96%
		4b , 95%
		4c , 94%
		4d , 95%
		4e , 93%
		4f , 96%
		4g , 95%
		4h , 95%
		4i , 96%
		4j , 95%
		4k , 97%
		4l , 96%
		4m , 95%
		4n , 90%
		4p , 94%
		4q , 98%
		4r , 98%

^a Reaction conditions: 3 (0.5 mmol), AgOTf (0.025 mmol), D₂O (10 mmol), CDCl₃ (1.0 mL), 90 °C for 18 h. Isolated yield are given.

Table 4 Silver-catalyzed α -deuteration of 2-alkyl azaarenes^a

	AgOTf (5 mol %), D ₂ O (20 equiv) THF, 90 °C, 18 h	
		6a , 97%
		6b , 98%
		6c , 96%
		6d , 92%
		6e , 94%
		6f , 96%

^a Reaction conditions: 5 (0.5 mmol), AgOTf (0.025 mmol), D₂O (10 mmol), THF (1.0 mL), 90 °C for 18 h. Isolated yield are given.

and benzothiazole under the standard conditions could not lead to D-incorporation.

Furthermore, by using this silver catalytic system, we also successfully achieved the regioselective deuteration at the α -positions of 2-alkyl azaarenes. As shown in Table 4, the deuteration of 1,2,3,4-tetrahydroacridine gave the deuterated product **6a** efficiently with 95% deuterium incorporation under the current reaction conditions. When the 2-methylquinoxaline was employed, both methyl group and C3 position were readily deuterated with high deuterium incorporation. In addition, the α -positions of alkyl-substituted quinolines **5c**, **5d** and 2-benzylpyridine **5e** could be deuterated, giving the corresponding products in good yields, albeit with 24–40% deuterium incorporation. Also, α -deuteration of oxindole **5f** gave deuterated product **6f** in 96% yield and 91% deuterium incorporation.

Conclusions

In summary, we have developed an efficient silver-catalyzed regioselective deuteration of (hetero)arenes and α -deuteration of 2-alkyl azaarenes using the simple D₂O as the deuterating reagent. The reaction proceeds under neutral mild conditions in high yields with a good to excellent degree of deuterium incorporation. Electron-rich arenes and heteroarenes, such as 1,2,3,4-tetrahydroquinolines, indoline derivatives, *N*-methyl anilines, 1,3,5-trimethoxybenzene and indoles, were all deuterated with high deuterium incorporation. The protocol was also applied to the deuteration of electron-deficient azoles. Moreover, the α -position deuteration of 2-alkyl azaarenes was also achieved under mild neutral conditions.

Conflicts of interest

There are no conflicts to declare.



Acknowledgements

We gratefully acknowledge National Natural Science Foundation of China (NSFC) (21702028), Department of Science and Technology of Jilin Province (20180520206JH), the Collaborative Fund of Luzhou Government and Southwest Medical University (2019LZXNYDJ16) and the Research Fund of Southwest Medical University (2017-ZRQN-037) for financial support.

Notes and references

- 1 (a) J. Atzrodt and V. J. Derdau, Pd- and Pt-catalyzed H/D exchange methods and their application for internal MS standard preparation from a Sanofi-Aventis perspective, *J. Label. Compd. Radiopharm.*, 2010, **53**, 674–685; (b) A. Nakanishi, Y. Fukushima, N. Miyazawa, K. Yoshikawa, T. Maeda and Y. J. Kurobayashi, Quantitation of Rotundone in Grapefruit (*Citrus paradisi*) Peel and Juice by Stable Isotope Dilution Assay, *J. Agric. Food Chem.*, 2017, **65**, 5026–5033; (c) E. Stokvis, H. Rosing and J. H. Beijnen, Stable isotopically labeled internal standards in quantitative bioanalysis using liquid chromatography/mass spectrometry: necessity or not?, *Rapid Commun. Mass Spectrom.*, 2005, **19**, 401–407; (d) C.-Y. Kao and R. W. Giese, Measurement of N7-(2'-Hydroxyethyl)guanine in Human DNA by Gas Chromatography Electron Capture Mass Spectrometry, *Chem. Res. Toxicol.*, 2005, **18**, 70–75.
- 2 (a) E. M. Simmons and J. F. Hartwig, On the Interpretation of Deuterium Kinetic Isotope Effects in C–H Bond Functionalizations by Transition-Metal Complexes, *Angew. Chem., Int. Ed.*, 2012, **51**, 3066–3072; (b) T. Giagou and M. P. Meyer, Kinetic Isotope Effects in Asymmetric Reactions, *Chem.–Eur. J.*, 2010, **16**, 10616–10628.
- 3 (a) E. M. Isin, C. S. Elmore, G. N. Nilsson, R. A. Thompson and L. Weidolf, Use of Radiolabeled Compounds in Drug Metabolism and Pharmacokinetic Studies, *Chem. Res. Toxicol.*, 2012, **25**, 532–542; (b) P. Uhl, G. Fricker, U. Haberkorn and W. Mier, Radionuclides in drug development, *Drug Discovery Today*, 2015, **20**, 198–208; (c) A. E. Mutlib, Application of Stable Isotope-Labeled Compounds in Metabolism and in Metabolism-Mediated Toxicity Studies, *Chem. Res. Toxicol.*, 2008, **21**, 1672–1689; (d) S. Nag, L. Lehmann, G. Kettschau, M. Toth, T. Heinrich, A. Thiele, A. Varrone and C. Halldin, Development of a novel fluorine-18 labeled deuterated fluororasagiline ($[^{18}\text{F}]\text{fluororasagiline-D}_2$) radioligand for PET studies of monoamino oxidase B (MAO-B), *Bioorg. Med. Chem.*, 2013, **21**, 6634–6641; (e) K. Y. Wong, Y. Xu and L. Xu, Review of computer simulations of isotope effects on biochemical reactions: from the Bigeleisen equation to Feynman's path integral, *Biochim. Biophys. Acta, Proteins Proteomics*, 2015, **1854**, 1782–1794.
- 4 (a) A. Katsnelson, Heavy drugs draw heavy interest from pharma backers, *Nat. Med.*, 2013, **19**, 656; (b) Y. Zhang, M. D. Tortorella, Y. Wang, J. Liu, Z. Tu, X. Liu, Y. Bai, D. Wen, X. Lu, Y. Lu and J. Talley, Synthesis of Deuterated Benzopyran Derivatives as Selective COX-2 Inhibitors with Improved Pharmacokinetic Properties, *ACS Med. Chem. Lett.*, 2014, **5**, 1162–1166; (c) Y. Zhu, J. Zhou and B. Jiao, Deuterated Clopidogrel Analogues as a New Generation of Antiplatelet Agents, *ACS Med. Chem. Lett.*, 2013, **4**, 349–352; (d) A. Mullard, Deuterated drugs draw heavier backing, *Nat. Rev. Drug Discovery*, 2016, **15**, 219–221; (e) T. G. Gant, Using Deuterium in Drug Discovery: Leaving the Label in the Drug, *J. Med. Chem.*, 2014, **57**, 3595–3611.
- 5 (a) T. Junk and W. J. Catallo, Hydrogen isotope exchange reactions involving C–H (D, T) bonds, *Chem. Soc. Rev.*, 1997, **26**, 401–406; (b) Y. Sawama, Y. Monguchi and H. Sajiki, Efficient H–D Exchange Reactions Using Heterogeneous Platinum-Group Metal on Carbon–H₂–D₂O System, *Synlett*, 2012, **23**, 959–972; (c) J. Atzrodt, V. Derdau, W. J. Kerr and M. Reid, C–H Functionalisation for Hydrogen Isotope Exchange, *Angew. Chem., Int. Ed.*, 2018, **57**, 3022–3047; (d) J. Atzrodt, V. Derdau, T. Fey and J. Zimmermann, The Renaissance of H/D Exchange, *Angew. Chem., Int. Ed.*, 2007, **46**, 7744–7765.
- 6 (a) A. Martins and M. Lautens, A Simple, Cost-Effective Method for the Regioselective Deuteration of Anilines, *Org. Lett.*, 2008, **10**, 4351–4353; (b) S. Rasku, K. Whl, J. Koskimies and T. Hase, Synthesis of isoflavanoid deuterium labeled polyphenolic phytoestrogens, *Tetrahedron*, 1999, **55**, 3445–3454; (c) J. C. Seibles, D. M. Bollinger and M. Orchin, Synthesis of Perdeuteriobenzo[a]pyrene, *Angew. Chem., Int. Ed. Engl.*, 1977, **16**, 656–657; (d) J. L. Garnett, M. A. Long, R. F. W. Vining and T. Mole, New simple method for rapid, selective aromatic deuteration using organoaluminum dihalide catalysts, *J. Am. Chem. Soc.*, 1972, **94**, 5913–5914; (e) M. Pohjoispää, R. Mera-Adasme, D. Sundholm, S. Heikkinen, T. Hase and K. J. Wähälä, Capricious Selectivity in Electrophilic Deuteration of Methylenedioxy Substituted Aromatic Compounds, *Org. Chem.*, 2014, **79**, 10636–10640.
- 7 (a) W. J. Kerr, D. M. Lindsay, M. Reid, J. Atzrodt, V. Derdau, P. Rojahn and R. Weck, Iridium-catalysed ortho-H/D and -H/T exchange under basic conditions: C–H activation of unprotected tetrazoles, *Chem. Commun.*, 2016, **52**, 6669–6672; (b) W. J. Kerr, M. Reid and T. Tuttle, Iridium-Catalyzed C–H Activation and Deuteration of Primary Sulfonamides: An Experimental and Computational Study, *ACS Catal.*, 2015, **5**, 402–410; (c) B. McAuley, M. J. Hickey, L. P. Kingston, J. R. Jones, W. J. S. Lockley, A. N. Mather, E. Spink, S. P. Thompson and D. J. J. Wilkinson, Convenient and efficient deuteration of functionalized aromatics with deuterium oxide: catalysis by cycloocta-1,5-dienyliridium(I) 1,3-dionates, *J. Label. Compd. Radiopharm.*, 2003, **46**, 1191–1204; (d) D. Hesk, P. R. Das and B. J. Evans, Deuteration of acetanilides and other substituted aromatics using $[\text{Ir}(\text{COD})(\text{Cy}_3\text{P})(\text{Py})]\text{PF}_6$ as catalyst, *J. Label. Compd. Radiopharm.*, 1995, **36**, 497–502; (e) J. R. Heys, A. Y. L. Shu, S. G. Senderoff and N. M. J. Phillips, Deuterium exchange labelling of substituted aromatics using $[\text{IrH}_2(\text{Me}_2\text{CO})_2(\text{PPh}_3)_2]\text{BF}_4$, *J. Label. Compd. Radiopharm.*, 1993, **33**, 431–438; (f) M. J. Hickey, J. R. Jones,



L. P. Kingston, W. J. S. Lockley, A. N. Mather, B. M. McAuley and D. J. Wilkinson, Iridium-catalysed labelling of anilines, benzylamines and nitrogen heterocycles using deuterium gas and cycloocta-1,5-dienyliridium(I) 1,1,1,5,5,5-hexafluoropentane-2,4-dionate, *Tetrahedron Lett.*, 2003, **44**, 3959–3961; (g) D. Giunta, M. Hölscher, C. W. Lehmann, R. Mynott, C. Wirtz and W. Leitner, Room Temperature Activation of Aromatic C–H Bonds by Non-Classical Ruthenium Hydride Complexes Containing Carbene Ligands, *Adv. Synth. Catal.*, 2003, **345**, 1139–1145; (h) B. Groll, M. Schnurch and M. D. Mihovilovic, Selective Ru(0)-Catalyzed Deuteration of Electron-Rich and Electron-Poor Nitrogen-Containing Heterocycles, *J. Org. Chem.*, 2012, **77**, 4432–4437; (i) C. P. Lenges, P. S. White and M. Brookhart, Hydrogen/Deuterium Exchange Reactions and Transfer Hydrogenations Catalyzed by $[C_5Me_5Rh(\text{olefin})_2]$ Complexes: Conversion of Alkoxy silanes to Silyl Enolates, *J. Am. Chem. Soc.*, 1999, **121**, 4385–4396; (j) M. Zhan, H. Jiang, X. Pang, T. Zhang, R. Xu, L. Zhao, Y. Liu, Y. Gong and Y. Chen, A convenient method for the Ru(0)-catalyzed regioselective deuteration of N-alkyl-substituted anilines, *Tetrahedron Lett.*, 2014, **55**, 5070–5073; (k) S. Ma, G. Villa, P. S. Thuy-Boun, A. Homs and J. Q. Yu, Palladium-Catalyzed *ortho*-Selective C–H Deuteration of Arenes: Evidence for Superior Reactivity of Weakly Coordinated Palladacycles, *Angew. Chem., Int. Ed.*, 2014, **53**, 734–737; (l) A. M. Krause-Heuer, N. R. Yepuri, T. A. Darwish and P. J. Holden, Mild Conditions for Deuteration of Primary and Secondary Arylamines for the Synthesis of Deuterated Optoelectronic Organic Molecules, *Molecules*, 2014, **19**, 18604–18617; (m) H. Sajiki, N. Ito, H. Esaki, T. Maesawa, T. Maegawa and K. Hirota, Aromatic ring favorable and efficient H–D exchange reaction catalyzed by Pt/C, *Tetrahedron Lett.*, 2005, **46**, 6995–6998; (n) V. Derdau, J. Atzrodt, J. Zimmermann, C. Kroll and F. Brückner, Hydrogen-Deuterium Exchange Reactions of Aromatic Compounds and Heterocycles by NaBD₄-Activated Rhodium, Platinum and Palladium Catalysts, *Chem.-Eur. J.*, 2009, **15**, 10397–10404; (o) D. Zhao, H. Luo, B. Chen, W. Chen, G. Zhang and Y. Yu, Palladium-Catalyzed H/D Exchange Reaction with 8-Aminoquinoline as the Directing Group: Access to *ortho*-Selective Deuterated Aromatic Acids and β -Selective Deuterated Aliphatic Acids, *J. Org. Chem.*, 2018, **83**, 7860–7866; (p) Y. Sawama, K. Park, T. Yamada and H. Sajiki, New Gateways to the Platinum Group Metal-Catalyzed Direct Deuterium-Labeling Method Utilizing Hydrogen as a Catalyst Activator, *Chem. Pharm. Bull.*, 2018, **66**, 21–28; (q) J. Krüger, B. Manmontri and G. Fels, Iridium-Catalyzed H/D Exchange, *Eur. J. Org. Chem.*, 2005, 1402–1408; (r) T. Yamada, K. Park, N. Yasukawa, K. Morita, Y. Monguchi, Y. Sawama and H. Sajiki, Mild and Direct Multiple Deuterium-Labeling of Saturated Fatty Acids, *Adv. Synth. Catal.*, 2016, **358**, 3277–3282.

8 (a) M. Naodovic and H. Yamamoto, Asymmetric Silver-Catalyzed Reactions, *Chem. Rev.*, 2008, **108**, 3132–3148; (b) Y. Yamamoto, Silver-Catalyzed C_{sp}–H and C_{sp}–Si Bond Transformations and Related Processes, *Chem. Rev.*, 2008, **108**, 3199–3222; (c) Q.-Z. Zheng and N. Jiao, Ag-catalyzed C–H/C–C bond functionalization, *Chem. Soc. Rev.*, 2016, **45**, 4590–4627; (d) G. Fang and X. Bi, Silver-catalysed reactions of alkynes: recent advances, *Chem. Soc. Rev.*, 2015, **44**, 8124–8173; (e) K. Sekine and T. Yamada, Silver-catalyzed carboxylation, *Chem. Soc. Rev.*, 2016, **45**, 4524–4532; (f) M. P. Muñoz, Silver and platinum-catalysed addition of O–H and N–H bonds to alkenes, *Chem. Soc. Rev.*, 2014, **43**, 3164–3183.

9 (a) An example on the catalytic H/D exchange of arenes in acidic solvents, in which silver salt was examined, see: D. Munz, M. Webster-Gardiner, R. Fu, T. Strassner, W. A. Goddard III and T. B. Gunnoe, Proton or Metal? The H/D Exchange of Arenes in Acidic Solvents, *ACS Catal.*, 2015, **5**, 769–775; (b) An example on the catalytic H/D exchange of hydrogen isotope exchange of five-membered heteroarenes by Ag₂CO₃/JohnPhos, see: E.-C. Li, G.-Q. Hu, Y.-X. Zhu, H.-H. Zhang, K. Shen, X.-C. Hang, C. Zhang and W. Huang, Ag₂CO₃-Catalyzed H/D Exchange of Five-Membered Heteroarenes at Ambient Temperature, *Org. Lett.*, 2019, **21**, 6745–6749.

10 Other Lewis acids were also tested under the standard condition, however, gave a lower deuterium incorporation, especially in the *para*-position, Cu(OTf)₂ (75% D *ortho*, 75% D *para*), Zn(OTf)₂ (60% D *ortho*, 20% D *para*), ZnCl₂ (24% D *ortho*, 14% D *para*), FeCl₃ (75% D *ortho*, 40% D *para*), and La(OTf)₃ (<10% D *ortho*, <10% D *para*).

11 Recently, Werner and coworkers reported B(C₆F₅)₃-catalyzed regioselective deuteration of electron-rich aromatic and heteroaromatic compounds, see: W. Li, M.-M. Wang, Y. Hu and T. Werner, B(C₆F₅)₃-Catalyzed Regioselective Deuteration of Electron-Rich Aromatic and Heteroaromatic Compounds, *Org. Lett.*, 2017, **19**, 5768–5771.

12 (a) T. Eicher and S. Hauptmann, *The Chemistry of Heterocycles*, Wiley-VCH, Weinheim, 2003; (b) M. Ishikura, T. Abe, T. Choshi and S. Hibino, Simple indole alkaloids and those with a non-rearranged monoterpenoid unit, *Nat. Prod. Rep.*, 2013, **30**, 694–752; (c) V. Sharma, P. Kumar and D. Pathak, Biological Importance of the Indole Nucleus in Recent Years: A Comprehensive Review, *J. Heterocycl. Chem.*, 2010, **47**, 491–502; (d) A. J. Kochanowska-Karamyan and M. T. Hamann, Marine Indole Alkaloids: Potential New Drug Leads for the Control of Depression and Anxiety, *Chem. Rev.*, 2010, **110**, 4489–4497; (e) J. S. Carey, D. Laffan, C. Thomson and M. T. Williams, Analysis of the reactions used for the preparation of drug candidate molecules, *Org. Biomol. Chem.*, 2006, **4**, 2337–2347.

