


 Cite this: *RSC Adv.*, 2020, 10, 18953

# Studies on asymmetric total synthesis of (–)-β-hydrastine via a chiral epoxide ring-opening cascade cyclization strategy†

 Jihui Li,<sup>a</sup> Tianxiao Wu,<sup>a</sup> Xinjing Song,<sup>a</sup> Yang Zheng,<sup>a</sup> Jiaxin Meng,<sup>a</sup> Qiaohua Qin,<sup>a</sup> Yongxiang Liu,<sup>id</sup> abc Dongmei Zhao<sup>id</sup> \*a and Maosheng Cheng<sup>a</sup>

Herein, facile and enantioselective approaches to synthesize the core phthalide tetrahydroisoquinoline scaffold of (–)-β-hydrastine via both a CF<sub>3</sub>COOH-catalyzed (86% ee) and KHMDS-catalyzed (78% ee) epoxide ring-opening/transesterification cascade cyclization from chiral epoxide under very mild conditions are described. The key elements include a highly enantioselective epoxidation using the Shi ketone catalyst and an intramolecular CF<sub>3</sub>COOH-catalyzed cascade cyclization in one pot, and a late-stage C-3' epimerization under MeOK/MeOH conditions as the key steps to achieve the first total synthesis of (–)-β-hydrastine (up to 81% ee).

Received 4th April 2020

Accepted 11th May 2020

DOI: 10.1039/d0ra03038d


[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## Introduction

Natural products that feature multiple stereocenters and complex molecular topologies hold a special fascination to chemists. Phthalide tetrahydroisoquinoline alkaloids, such as hydrastine<sup>1</sup> (**1**, Fig. 1), are known to possess interesting and diverse biological properties.<sup>2</sup> In light of their fascinating architectural attributes and unique structures, as well as promising biological activities, phthalide tetrahydroisoquinoline alkaloids have been attractive and challenging targets of synthetic and medicinal chemists for decades. However, due to the small quantities of the natural products including hydrastine (**1**), a comprehensive biological evaluation of this class of alkaloids could not be accomplished. To accelerate further investigation of pharmacological activities and SAR studies, we embarked on the asymmetric total syntheses of this class of alkaloids.

Hydrastine (**1**), originally isolated from *Corydalis longipes*<sup>3</sup> has been used as an antiseptic,<sup>4</sup> as a uterine hemostatic<sup>4</sup> and as a potent competitive antagonist at mammalian GABA<sub>A</sub> receptors (CD<sub>50</sub> 0.16 mg kg<sup>-1</sup>, i.v.).<sup>5</sup> In 2016, (–)-β-hydrastine [(–)-β-**1**] was found to exhibit potent antitumor activity against human lung adenocarcinoma cells.<sup>6</sup> Structurally, hydrastine (**1**)

comprises a 1,2,3,4-tetrahydroisoquinoline and a phthalide ring and these two units are linked together by two chiral centers at their C-1 and C-3' carbons to form either *erythro*- or *threo*-isomer (Fig. 1).<sup>7</sup> Because of unique structural features and interesting biological properties of hydrastine, the design of new protocols for the construction of this privileged scaffold has attracted considerable interest. Since the work for the synthesis of hydrastine (**1**) by Robinson and co-workers using the direct condensation strategy in 1931 (Scheme 1a),<sup>8</sup> several synthetic studies<sup>9,10</sup> had been reported for the synthesis of hydrastine (**1**). Among them, (±)-β-hydrastine and (±)-α-hydrastine were obtained from the substituted β-phenylethylmethylamide of a substituted phthalidecarboxylic acid by dehydrated ring-closure and following reduction in 1950 (Scheme 1b),<sup>9</sup> and the semi-synthesis of (±)-β-hydrastine and (±)-α-hydrastine from natural berberine was reported (Scheme 1c).<sup>10</sup> In addition, several other approaches<sup>11,12</sup> to (±)-β-hydrastine and (±)-α-hydrastine were reported, such as the electrochemical<sup>11a</sup> and zinc-promoted reductive coupling<sup>11b</sup> (Scheme 1d). Alternatively, the phthalide tetrahydroisoquinoline skeleton of hydrastine (**1**) could also be synthesized through the Bischler–Napieralski


 Fig. 1 The *erythro*- and *threo*-form of hydrastine (**1**).


<sup>a</sup>Key Laboratory of Structure-Based Drug Design & Discovery of Ministry of Education, Shenyang Pharmaceutical University, Shenyang, 110016, People's Republic of China. E-mail: medchemzhao@163.com

<sup>b</sup>Wuyia College of Innovation, Shenyang Pharmaceutical University, Shenyang 110016, People's Republic of China

<sup>c</sup>Institute of Drug Research in Medicine Capital of China, Benxi, 117000, People's Republic of China

† Electronic supplementary information (ESI) available: Experimental details, characterization data, NMR spectra and HPLC chromatograms for products. See DOI: 10.1039/d0ra03038d






Scheme 1 The known strategic approaches to hydrastine. (a) The direct condensation strategy. (b) The dehydrated ring-closure strategy. (c) The semi-synthesis strategy. (d) The reductive coupling strategy. (e) The Bischler–Napieralski cyclization strategy.

reaction<sup>12</sup> (Scheme 1e) in the similar fashion of the synthesis of noscapine.<sup>13</sup> Although these methods successfully achieved the synthesis of hydrastine (**1**), an inherent drawback associated with all of the above-mentioned strategies was the lack of enantioselectivity. Thus, the efficient, practical and highly enantioselective construction of phthalide tetrahydroisoquinoline structures still constitutes a synthetic challenge. To our best knowledge, no attempt to synthesize (–)-β-hydrastine [(–)-β-**1**] with high enantioselectivity has been reported.

Our research group has been exploring a unified acid-catalyzed epoxide ring-opening/intramolecular transesterification cascade cyclization strategy for synthesis of phthalide tetrahydroisoquinoline alkaloids, in which the C and D rings are constructed simultaneously in one step.<sup>14</sup> The method was successfully applied to the diastereoselective total



Scheme 2 (a) Our previous work to the total synthesis of (±)-β-hydrastine. (b) Our proposed synthetic strategy to the total synthesis of (–)-β-hydrastine in this study.

synthesis of (±)-β-hydrastine *via* an  $S_N1$ -like mechanism (Scheme 2a).<sup>14</sup> In order to investigate further applications of this cascade cyclization strategy for enantioselective total syntheses of phthalide tetrahydroisoquinoline alkaloids, we postulated that the chiral epoxide was utilized as a key stereocontrol element in addition to a reactivity control element to enantioselectively construct the phthalide tetrahydroisoquinoline scaffold based on an  $S_N1$ -type acid-catalyzed or an  $S_N2$ -type base-catalyzed cyclization strategy (Scheme 2b). As the Shi ketone catalyst and other variants are synthetically accessible,<sup>15,16</sup> we have the ability to generate epoxide substrate bearing the necessary stereochemistry based on the chosen cascade direction. In this paper, we document the success of the cascade cyclization strategy in the form of the first enantioselective total synthesis of (–)-β-hydrastine.


## Results and discussion

In designing a synthetic route toward (–)-β-hydrastine, we employed the chiral epoxide in order to control both the reactivity and stereoselectivity and envisioned a crucial cascade cyclization transformation, namely, an acid-catalyzed cascade cyclization. Our retrosynthetic analysis of (–)-β-hydrastine was outlined in Scheme 3. The target natural compound (–)-β-hydrastine could be accessed from (–)-α-hydrastine *via* epimerization at C-3', and (–)-α-hydrastine could be obtained from (–)-α-**2** after methylation. Then, we speculated that the phthalide tetrahydroisoquinoline core of (–)-α-**2** could be constructed by an acid-catalyzed epoxide ring-opening/transesterification cascade cyclization of the (R,R)-**3**, which could be prepared from (E)-stilbene **4** by asymmetric epoxidation with Shi ketone catalyst.<sup>15</sup> Moreover, the stilbene **4** might be constructed from iodide **5** and substituted styrene **6** by a Pd-catalyzed Heck coupling reaction.<sup>14,17</sup> The iodide **5** could be synthesized by iodization of *N*-protected phenethylamine **7**,<sup>14</sup> which could be obtained from commercially available vanillin (**9**). In addition,






**Table 1** Optimization of asymmetric epoxidation of (*E*)-stilbene **4** for the acid-catalyzed cascade cyclization



Entry <sup>a</sup>	Ketone <b>13</b> (equiv.)	Conditions <sup>b</sup>	Yield <sup>c</sup> (%)	ee <sup>d</sup> (%)
1	3.0	A	ND <sup>e</sup>	—
2	3.0	B	ND <sup>e</sup>	—
3	5.0	B	ND <sup>e</sup>	—
4	0.3	C	ND <sup>e</sup>	—
5	0.3	D	ND <sup>e</sup>	—
6	1.6	D <sup>f</sup>	ND <sup>e</sup>	—
7	2.0	D <sup>f</sup>	ND <sup>e</sup>	—
8	3.0	D <sup>f</sup>	ND <sup>e</sup>	—
9	3.0	D <sup>f,g</sup>	Trace <sup>h</sup>	—
10	1.6	D <sup>f,g</sup>	Trace <sup>h</sup>	—
11	1.6	D <sup>g</sup>	Trace <sup>h</sup>	—
12 <sup>i</sup>	0.3	E	ND <sup>e</sup>	—
13	3.0	E	Trace <sup>j</sup>	—
14 <sup>k</sup>	3.0	F	42	86
15 <sup>k</sup>	1.0	F <sup>l</sup>	24	79
16 <sup>m</sup>	3.0	G	25	58
17 <sup>m</sup>	3.0	H	40	71

<sup>a</sup> Substrate **4** (0.1 mmol). <sup>b</sup> Condition A: Oxone (5.6 equiv.), NaHCO<sub>3</sub> (18.9 equiv.), CH<sub>3</sub>CN-CH<sub>2</sub>Cl<sub>2</sub>-H<sub>2</sub>O (v/v/v = 1 : 4 : 5), 0 °C, 48 h. Condition B: Oxone (5.0 equiv.), NaHCO<sub>3</sub> (15.5 equiv.), Bu<sub>4</sub>NHSO<sub>4</sub> (5 mol%), CH<sub>3</sub>CN-aq. Na<sub>2</sub>EDTA (4 × 10<sup>-4</sup> M) (v/v = 1.5 : 1), 0 °C, 2.0 h. Condition C: Oxone (2.02 equiv.), K<sub>2</sub>CO<sub>3</sub> (4.04 equiv.), Bu<sub>4</sub>NHSO<sub>4</sub> (5 mol%), CH<sub>3</sub>CN-DMM-0.05 M aq. Na<sub>2</sub>HPO<sub>4</sub>-0.05 M aq. KH<sub>2</sub>PO<sub>4</sub> (pH = 7.0) (v/v/v = 1 : 2 : 1), 0 °C, 24 h. Condition D: Oxone (1.38 equiv.), K<sub>2</sub>CO<sub>3</sub> (5.8 equiv.), Bu<sub>4</sub>NHSO<sub>4</sub> (5 mol%), CH<sub>3</sub>CN-DMM-0.05 M Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub>·10H<sub>2</sub>O of aq. Na<sub>2</sub>EDTA (4 × 10<sup>-4</sup> M) solution (v/v/v = 1 : 2 : 2), 0 °C, 1.5 h. Condition E: Oxone (1.38 equiv.), K<sub>2</sub>CO<sub>3</sub> (5.8 equiv.), Bu<sub>4</sub>NHSO<sub>4</sub> (5 mol%), CH<sub>3</sub>CN-0.05 M Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub>·10H<sub>2</sub>O of aq. Na<sub>2</sub>EDTA (4 × 10<sup>-4</sup> M) solution (v/v = 3 : 2), 0 °C to rt, 24 h. Condition F: Oxone (4.6 equiv.), K<sub>2</sub>CO<sub>3</sub> (18.6 equiv.), Bu<sub>4</sub>NHSO<sub>4</sub> (5 mol%), CH<sub>3</sub>CN-0.05 M Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub>·10H<sub>2</sub>O of aq. Na<sub>2</sub>EDTA (4 × 10<sup>-4</sup> M) solution (v/v = 3 : 2, 25 mL), 0 °C, 4.5 h. Condition G: 30% H<sub>2</sub>O<sub>2</sub> (30 equiv.), CH<sub>3</sub>CN-1.0 M K<sub>2</sub>CO<sub>3</sub> in 4 × 10<sup>-4</sup> M of EDTA (v/v = 2 : 1), 0 °C, 36 h. Condition H: 30% H<sub>2</sub>O<sub>2</sub> (30 equiv.), CH<sub>3</sub>CN-EtOH-CH<sub>2</sub>Cl<sub>2</sub> (v/v/v = 1 : 1 : 2), 2.0 M K<sub>2</sub>CO<sub>3</sub> in 4 × 10<sup>-4</sup> M of EDTA, 0 °C, 36 h. <sup>c</sup> Isolated yield of (-)-α-2. <sup>d</sup> The enantiomeric excess was determined by chiral HPLC (Chiralpak IH). <sup>e</sup> The epoxide (**R,R**)-**3** was not detected by TLC and the starting material **4** was recovered. <sup>f</sup> Oxone (1.38 equiv.) and K<sub>2</sub>CO<sub>3</sub> (5.8 equiv.) were used. <sup>g</sup> The reactions were stopped after 2 h for -10 °C. <sup>h</sup> Most of the starting material **4** was recovered. <sup>i</sup> Substrate **4** (1.0 mmol). <sup>j</sup> Most of the epoxide (**R,R**)-**3** was decomposed. <sup>k</sup> Substrate **4** (0.3 mmol). <sup>l</sup> The reactions were stopped after 8 h for 0 °C. <sup>m</sup> Substrate **4** (0.4 mmol). DMM = dimethoxymethane.

With highly enantioselective (-)-α-2 in hand, we were pleased to synthesize (-)-β-hydrastine [(-)-β-1] using already established conditions for two transformations.<sup>14</sup> The *N*-methylation of (-)-α-2 occurred under classic Eschweiler-Clarke reaction conditions,<sup>14</sup> thus yielding (-)-α-hydrastine [(-)-α-1]. Remarkably, without purification of (-)-α-2, (-)-α-hydrastine was prepared from **4** in the three-step yield of 30% (see Scheme 5). Finally, a late-stage epimerization of (-)-α-1 at C-3' under MeOK/MeOH conditions



**Scheme 7** Syntheses of (*E*)-stilbenes **16a-c**.


furnished (-)-β-hydrastine [(-)-β-1] with 81% ee, which was in agreement with authentic reference standards for this natural product<sup>1</sup> (see the ESI†).

Therefore, (-)-β-hydrastine had been synthesized with a high enantiomeric excess (81% ee) by using CF<sub>3</sub>COOH-catalyzed cascade cyclization reaction and C-3' epimerization under MeOK/MeOH conditions as the key steps (Scheme 5). Inspired by the ring-opening reaction of epoxides in basic conditions *via* an S<sub>N</sub>2-type mechanism,<sup>20</sup> we also assumed that the core phthalide tetrahydroisoquinoline scaffold with the *R* and *S* configurations at C-1 and C-3' could be enantioselectively structured based on a base-catalyzed epoxide ring-opening/transesterification cascade cyclization strategy in order to avoid the late-stage epimerization reaction at C-3'. Correspondingly, the chiral epoxide with (*S,S*) configurations was chosen as the substrate to direct the cascade reaction under basic conditions. To test the working hypothesis, initially, we prepared a series of racemic model substrates [(±)-**17**] through DMDO-mediated epoxidation<sup>14</sup> from the corresponding (*E*)-stilbenes (**16**), which were subjected to base-catalyzed cascade cyclization model reactions. Similarly, the precursors **16** for epoxidation were prepared in three steps, namely, *N*-protection, I<sub>2</sub>/CF<sub>3</sub>COOAg-mediated iodination and following Pd-catalyzed Heck coupling reaction (Scheme 7).

With the (*E*)-stilbenes **16a-c** in hand, we focused our attention on the base-catalyzed cascade cyclization model reactions. Initially, we chosen trifluoroacetyl<sup>21</sup> as the *N*-protection group to examine the reactivity in the presence 60% NaH<sup>21</sup> in 1,4-dioxane, unfortunately, leading to no desired cyclization product (±)-β-**18a** (Table 2, entry 1). After that, we further optimized the reaction conditions of the model reaction, and the results were summarized in Table 2. Using (±)-**17a** as the model substrate, firstly, extensive screening of different bases (KHMDS, LDA, and *n*-BuLi), solvents (THF and 1,4-dioxane), and reaction temperatures resulted in no desired (±)-β-**18a** or the decomposition of (±)-**17a** (Table 2, entries 2–7). In our opinion, it was probably due to that the electron-withdrawing ability of trifluoroacetyl group was not strong enough to fulfill the deprotonation process of amino in the presence of strong bases, and then failing to facilitate desired cascade cyclization reaction. In light of the above reason, the 4-(trifluoromethyl)benzenesulfonyl, a much stronger electron-withdrawing group,



Table 2 Optimization of racemic model substrates [(±)-17] for the base-catalyzed cascade cyclization




Entry <sup>a</sup>	17	Base (equiv.)	Solvent	Temp. (°C)	T (h)	Yield <sup>b</sup> (%)
1	17a	60% NaH (2.5)	1,4-Dioxane	0–65	12	— <sup>c</sup>
2	17a	KHMDS (1.2)	1,4-Dioxane	0–65	12	ND <sup>d</sup>
3	17a	KHMDS (1.2)	1,4-Dioxane	0–84	12	— <sup>c</sup>
4	17a	KHMDS (1.2)	Dry THF	0–65	12	ND <sup>d</sup>
5	17a	KHMDS (1.2)	Dry THF	0 to rt	12	ND <sup>d</sup>
6	17a	LDA (1.2)	Dry THF	0–65	12	— <sup>c</sup>
7	17a	<i>n</i> -BuLi (1.2)	Dry THF	–78 to rt	12	— <sup>c</sup>
8	17b	KHMDS (1.2)	Dry THF	0–65	12	11
9	17b	KHMDS (1.5)	1,4-Dioxane	0–65	12	10
10	17b	<b>KHMDS (1.5)</b>	<b>Dry THF</b>	<b>0–65</b>	<b>5.0</b>	<b>30</b>
11	17c	KHMDS (1.5)	Dry THF	0–65	12	ND <sup>d</sup>
12	17c	NaHMDS (1.5)	Dry THF	0–65	12	ND <sup>d</sup>
13	17c	KHMDS (1.5)	1,4-Dioxane	0–84	12	— <sup>c</sup>
14	17c	60% NaH (2.5)	1,4-Dioxane	0–65	12	15
15	17c	60% NaH (2.5)	1,4-Dioxane	0–65	24	10
16	17c	60% NaH (2.5)	1,4-Dioxane	0–60	12	16
17	17c	60% NaH (1.5)	1,4-Dioxane	0–60	5.0	19

<sup>a</sup> The substrate **16** (0.1 mmol) was dissolved in solvent and then the base was added in dropwise. <sup>b</sup> Isolated yield of (±)-β-18 in two steps. <sup>c</sup> Most of the epoxide (±)-17 was decomposed. <sup>d</sup> The (±)-β-18 was not detected by TLC and the epoxide (±)-17 was recovered. NaHMDS = sodium bis(trimethylsilyl)amide. KHMDS = potassium bis(trimethylsilyl)amide. LDA = lithium diisopropylamide.


was used and the corresponding (±)-17b was subjected to the model reactions (Table 2, entries 8–10). To our delight, the desired cyclization product (±)-β-18b was generated in 11% yield when the reaction was carried out by action of 1.0 M KHMDS in THF at 0–65 °C for 12 h (Table 2, entry 8). Encouraged by this result, we further optimized the reaction conditions by changing reaction solvent and temperature. As displayed in Table 2, the reaction solvent had little effect on the yield of (±)-β-18b (Table 2, entry 9). A decrease in the reaction time from 12 h to 5 h resulted in a three-time increase in yield for the same reaction solvent and temperature (Table 2, entry 10). In addition, considering that the steric hindrance of 4-(trifluoromethyl)

benzenesulfonyl might lead to lower yields, a mesyl-protected model substrate (±)-17c was prepared for further investigation of model reactions (Table 2, entries 11–17). The results revealed that no (±)-β-18c was delivered from (±)-17c in both KHMDS and NaHMDS conditions (Table 2, entries 11–13). However, the model substrate (±)-17c was conducted to 60% NaH in 1,4-dioxane, resulting in the desired (±)-β-18c in 15% yield (Table 2, entry 14). Moreover, attempts were made to improve the yields by extending the reaction time, lowering the reaction temperature and decreasing the equiv. of NaH, but the improvement was not significant (Table 2, entries 16–17). Finally, the optimal conditions were determined to be the model substrate (±)-17b in THF at 0–65 °C for 5 h in the presence of KHMDS (1.5 equiv.) with stirring (Table 2, entry 10).

Based on the abovementioned results, the (*E*)-stilbene **16b** was smoothly transformed to phthalide-tetrahydroisoquinoline-containing scaffold (–)-β-18b in 8% (30% brsm) yield over two steps with the enantiomeric excess (ee) of 78% (Scheme 8), where asymmetric epoxidation of **16b** with Shi catalyst *ent*-13 (ref. 15 and



Scheme 8 Synthesis of the phthalide tetrahydroisoquinoline core of (–)-β-hydrastine in method B.



Scheme 9 Synthesis of (*S,S*)-catalyst *ent*-13.



16) (Scheme 9) delivered (**S,S**)-**17b**, and the resulting epoxide underwent KHMDS-catalyzed cascade cyclization to give corresponding (–)-**β-18b**. After that, the total synthesis of (–)-**β-hydrastine** would be completed through the *N*-deprotection and the known methylation<sup>14</sup> reaction. Regarding to the late-stage *N*-deprotection of (–)-**β-18b**, several conditions, such as Mg/MeOH,<sup>22</sup> SmI<sub>2</sub>/THF,<sup>23</sup> Na-naphthalide/THF,<sup>24</sup> had been screened, but failing to afford (–)-**β-2** (see the ESI†) and further studies on the late-stage *N*-deprotection are still in progress.

## Conclusions

In conclusion, we have completed an efficient and stereo-controlled approach to total synthesis of the phthalide tetrahydroisoquinoline alkaloid (–)-**β-hydrastine** [(–)-**β-1**]. The key elements include a high enantioselective epoxidation using the Shi ketone catalyst and an intramolecular acid-catalyzed cascade cyclization in one pot, and a late-stage epimerization. Furthermore, it should be noted that the facile and enantioselective construction of the core phthalide tetrahydroisoquinoline scaffold was made possible by the CF<sub>3</sub>COOH-catalyzed or KHMDS-catalyzed epoxide ring-opening cascade cyclization from chiral epoxide under very mild conditions. In this reaction, the phthalide tetrahydroisoquinoline scaffold was enantioselectively constructed with one C–N bond and one C–C bond formed simultaneously. Efforts toward the asymmetric total syntheses of other phthalide tetrahydroisoquinoline alkaloids, such as (–)-**α-noscapine**, and the structure-related alkaloids, are currently ongoing in our laboratory. To conclude, the chemistry described herein would serve as an efficient alternative strategy to synthesize biologically active phthalide tetrahydroisoquinoline alkaloids and their derivatives.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We gratefully acknowledge the Program for Innovative Research Team of the Ministry of Education and the Program for Liaoning Innovative Research Team in University.

## References

- G. Blasko, D. J. Gula and M. Shamma, *J. Nat. Prod.*, 1982, **45**, 105–122.
- (a) S. Y. Yin, Y. M. Kim, J. J. Lee, C. M. Jin, Y. J. Yang, S.-K. Park, S. K. Yoo and M. K. Lee, *Biol. Pharm. Bull.*, 2007, **30**, 1547–1550; (b) S. Y. Yin, Y. M. Kim, J. J. Lee, C. M. Jin, Y. J. Yang, J. J. Ma, M. H. Kang, M. Kai and M. K. Lee, *Neuropharmacology*, 2004, **47**, 1045–1052; (c) M. K. Lee, Y. H. Zhang, J. S. Shin and S. S. Lee, *Med. Sci. Res.*, 1997, **25**, 619–620.
- (a) R. N. Jha, M. B. Pandey, A. K. Singh, S. Singh and V. P. Singh, *Nat. Prod. Res.*, 2009, **23**, 250–255; (b) M. Goel, U. P. Singh, R. N. Jha, V. B. Pandey and M. B. Pandey, *Folia Microbiol.*, 2003, **48**, 363–368.
- K. Goswami, V. Yadava, V. Padmanabhan and S. Banerjee, *Acta Crystallogr.*, 1993, **49**, 504–506.
- J. Huang and G. A. R. Johnston, *Br. J. Pharmacol.*, 1990, **99**, 727–730.
- B. Guo, X. Li, S. Song, M. Chen, M. Cheng, D. Zhao and F. Li, *Oncol. Rep.*, 2016, **35**, 2246–2256.
- (a) M. Shamma and V. S. Georgiev, *Tetrahedron Lett.*, 1974, **15**, 2339–2342; (b) K. Iwasa, M. Kamiguchi, M. Sugiura and N. Takao, *J. Nat. Prod.*, 1987, **50**, 1083–1088.
- E. Hope, F. L. Pyman, F. G. P. Remfry and R. Robinson, *J. Chem. Soc.*, 1931, 236–247.
- R. D. Haworth, A. R. Pinder and R. Robinson, *Nature*, 1950, **165**, 529.
- (a) J. L. Moniot and M. Shamma, *J. Org. Chem.*, 1979, **44**, 4337–4342; (b) J. L. Moniot and M. Shamma, *J. Am. Chem. Soc.*, 1976, **98**, 6714–6715.
- (a) T. Shono, Y. Usui and H. Hamaguchi, *Tetrahedron Lett.*, 1980, **21**, 1351–1354; (b) T. Shono, H. Hamaguchi, M. Sasaki, S. Fujita and K. Nagami, *J. Org. Chem.*, 1983, **48**, 1621–1628; (c) K. Orito, M. Miyazawa, R. Kanbayashi, M. Tokuda and H. Sugimoto, *J. Org. Chem.*, 1999, **64**, 6583–6596.
- L. Na, D. Zhao, S. Song, L. Sun and M. Cheng, *Acta Chim. Sin.*, 2011, **69**, 356–361.
- Z. Varga, G. Blasko, G. Dornyei and C. Szantay, *Acta Chim. Hung.*, 1991, **128**, 831–837.
- J. Li, Y. Liu, X. Song, T. Wu, J. Meng, Y. Zheng, Q. Qin, D. Zhao and M. Cheng, *Org. Lett.*, 2019, **21**, 7149–7153.
- (a) Z.-X. Wang, Y. Tu, M. Frohn, J.-R. Zhang and Y. Shi, *J. Am. Chem. Soc.*, 1997, **119**, 11224–11235; (b) Y. Tu, Z.-X. Wang and Y. Shi, *J. Am. Chem. Soc.*, 1996, **118**, 9806–9807.
- (a) J. Kang, G. J. Lim, S. K. Yoon and M. Y. Kim, *J. Org. Chem.*, 1995, **60**, 564–577; (b) S. Mio, Y. Kumagawa and S. Sugai, *Tetrahedron*, 1991, **47**, 2133–2144.
- D. Shabashov and O. Daugulis, *Org. Lett.*, 2006, **8**, 4947–4949.
- B. Wang, X.-Y. Wu, O. A. Wong, B. Nettles, M.-X. Zhao, D. Chen and Y. Shi, *J. Org. Chem.*, 2009, **74**, 3986–3989.
- (a) L. Shu and Y. Shi, *Tetrahedron*, 2001, **57**, 5213–5218; (b) L. Shu and Y. Shi, *Tetrahedron Lett.*, 1999, **40**, 8721–8724.
- (a) T. Angelini, F. Fringuelli, D. Lanari, F. Pizzo and L. Vaccaro, *Tetrahedron Lett.*, 2010, **51**, 1566–1569; (b) N. Oguni, Y. Miyagi and K. Itoh, *Tetrahedron Lett.*, 1998, **39**, 9023–9026; (c) D. L. Fox, A. A. Robinson, J. B. Frank and R. N. Salvatore, *Tetrahedron Lett.*, 2003, **44**, 7579–7582.
- D. Knight, S. L. E. Overman and G. Pairedeau, *J. Am. Chem. Soc.*, 1995, **117**, 5776–5788.
- (a) C. Qi, F. Hasenmaile, V. Gandon and D. Leboeuf, *ACS Catal.*, 2018, **8**, 1734–1739; (b) M. Nagamoto, T. Yanagi, T. Nishimura and H. Yorimitsu, *Org. Lett.*, 2016, **18**, 4474–4477.
- V. Pace, J. P. Rae and D. J. Procter, *Org. Lett.*, 2014, **16**, 476–479.
- N. Chakor, S. Dallavalle, L. Scaglioni and L. Merlini, *Eur. J. Org. Chem.*, 2010, 6217–6223.

