RSC Advances

PAPER

Cite this: RSC Adv., 2020, 10, 26006

Design, synthesis, biological evaluation and molecular docking study of novel pyridoxine– triazoles as anti-Alzheimer's agents†

Tiy[a](http://orcid.org/0000-0002-5700-8236)s Pal,^a Saipriyanka Bhimaneni,^b Abha Sharma **D**^{*a} and S. J. S. Flora D^{*b}

A series of multi-target natural product-pyridoxine based derivatives were designed, synthesized, characterized and evaluated as anti-Alzheimer agents. In vitro testing revealed the multi-functional properties of compounds such as inhibition of acetylcholinesterase (AChE), antioxidant and metal chelation. Among the series, 5i derivative was found most potent AChE inhibitor, possess antioxidant potential and chelating metal ions. Further binding interaction of 5i with AChE was studied using molecular docking, showed interaction with both PAS and CAS site of AChE. In silico predictions were also performed to predict toxicity and ADME properties of the molecule 5i and found within drug likeness range. Therefore, 5i could be a promising multi-functional compound that can be used for further development of novel drug for Alzheimer disease. **PAPER**
 EXERCT ACTIVE CONSIST AND THE VALUE CONSIST CONSIST AND CREAT CONSIST CONSIST AND CREAT CONSIST CONSIST THREE CONSISTS AND CONSIST CONSISTS THE CONSISTS AND CREAT CONSISTS THE CONSISTS ARE CONSISTENT THREE CONSI

Received 4th June 2020 Accepted 3rd July 2020 DOI: 10.1039/d0ra04942e

rsc.li/rsc-advances

1 Introduction

Alzheimer's disease (AD) is the most challenging neurodegenerative disease affecting mostly the population above 65 years of age. The perplexing nature of the disease makes it all more difficult for its ministration. More than 50 million are suffering globally and the number is likely to get hiked up to 150 million by the year 2050.¹ The multifaceted nature of the disease with several interconnected etiologies led to the idea of developing multi-target directed ligands (MTDLs). MTDLs in the field of neurodegenerative diseases is now a very signicant area of research developed by academia and industry.^{2,3} The design of MTDL comprises two or more pharmacophore moieties which are responsible for targeting multiple targets which are accountable for the pathogenesis of the disease. Therefore, MTDLs are single molecules with multiple functions which are strategically designed to serve the purpose of developing potential new chemical entities that can be used against AD.

The currently available Food and Drug Administration (FDA) approved drugs mainly focus on the cholinergic hypothesis, as one of the most excavated areas to slow down the progression of AD. Cholinergic hypothesis, amyloid-β cascade hypothesis, metal dyshomeostasis, tau hypothesis and oxidative stress hypothesis are some of the identified etiologies responsible for this disease. Damage to cholinergic neurons lead to cognitive decline which

showed the way to develop acetylcholinesterase (AChE) inhibitors providing symptomatic relief to the AD patients. However, this age related disease poses a problem of uncontrollable bio-burden of metal ions in the brain. This metal dyshomeostasis in turn induces oxidative stress and formation of amyloid- β (A β) plaques. Due to this interconnection of etiologies, classical metal chelator cannot do any benefit to this existing problem of the disease. Therefore MTDLs is needed in this juncture. Increase in biometals amplifies reactive oxygen species (ROS) which includes augmentation of lipid peroxidation, protein oxidation, nitration and glycol oxidation. Oxidative stress aggravates the pathophysiological conditions of neuronal damage.⁴ A molecule having additional anti-oxidant potency along with AChE inhibitory potential and metal chelating ability can be developed as a potential anti-AD agent. Studies have stated that peripheral anionic site (PAS) binding of AChE may promote A β aggregation. For PAS binding, aromatic rings with pi–pi stacking are recommended as derived from the docking studies with donepezil.⁵

In light of the concepts mentioned above, and to focus on our efforts to develop a new class of anti-Alzheimer's agents as MTDLs, we focused on the synthesis of pyridoxine based 1,2,3 triazoles and its biological evaluation. We assessed the synthesized compounds for AChE inhibition, antioxidant potency, metal chelation ability using in vitro assays as screening methods. We predicted various pharmacokinetic properties and toxicity of the best compound of the series using in silico tools.

2 Results and discussion

2.1 Design

Nitrogen heterocycles have long been of medical interest⁶⁻⁸ while, triazoles falling under the category of nitrogen

a Department of Medicinal Chemistry, National Institute of Pharmaceutical Education and Research, Raebareli, India. E-mail: abha.sharma@niperraebareli.edu.in

b Department of Regulatory Toxicology, National Institute of Pharmaceutical Education and Research, Raebareli, India. E-mail: sjsflora@hotmail.com

[†] Electronic supplementary information (ESI) available: Spectra data. See DOI: 10.1039/d0ra04942e

Paper RSC Advances and the contract of the con

heterocycles have been the moiety of interest. Numerous molecules with triazole moiety have been reported. In this regard, Rastegari et al. reported 1,2,3-triazole chromenone carboxamide derivatives as AChE inhibitors.⁹ Jalili-Baleh et al. developed 3-phenylcoumarin lipoic acid based triazole derivatives as anti-Alzheimer's agents.¹⁰ Fig. 1 provides a summary of few triazole and pyridoxine containing molecules reported recently as potential anti-Alzheimer's agents.⁹⁻¹⁵

Pyridoxine is known to reduce homocysteine levels in Alzheimer's patients. Homocysteine is derived from amino acid methionine and is a non-protein homologue of cysteine. The directly attached hydroxyl group to the pyridine nucleus might be playing a role in chelation of metal and also for the antioxidant properties. Pyridoxine has successfully completed the phase 3 clinical trials as a vitamin which can reduce homocysteine levels in AD affected patients. The circulating high level of homocysteine in AD is responsible for deposition of amyloid β plaques.^{16,17} Pyridoxine–resveratrol have been reported as only pyridoxine based compounds against AD. However, pyridoxine based triazoles have not been reported and therefore, herein we report this new class of compounds.

Taking into the consideration the multifaceted nature of the disease, we designed pyridoxine based 1,2,3-triazoles as an MTDL which might be an effective agents against AD. We have designed our target molecules containing both the pyridoxine and triazole moiety as a new class of MTDL using linking approach. Fig. 2 provides the possible hypothesis behind the designing of our target molecules.

2.2 Chemistry

We synthesized pyridoxine based triazoles derivatives (Scheme 1). The target molecules were synthesized via four step reaction scheme using pyridoxine hydrochloride (1) as starting material. In the first step, commercially available pyridoxine hydrochloride (1) was protected at the 3- and 4-hydroxy groups using ptoluenesulfonic acid monohydrate and dry acetone in the

Fig. 2 Design of target molecules chemistry.

Fig. 1 Triazole and pyridoxine containing molecules as anti-Alzheimer's agents.

presence of 2,2-dimethoxypropane under argon atmosphere. Then compound 2 was treated with propargyl bromide using NaH as base and dry THF as solvent under argon atmosphere to give compound 3. In the next step, various substituted benzyl azides were generated in situ from benzyl bromides followed by synthesis of triazole moiety after addition of compound 3 using the classical CuAAC, generating compounds 4(a–q). Finally, target compounds 5(a–q) were obtained through deprotection of the isopropylidene group.

2.3 In vitro assay of AChE inhibition

The *in vitro* activity of the new synthesized molecules was done using AChE from electric eel (EeAChE) following Ellman's method.²⁵ All the molecules were tested for their ability to inhibit AChE activity, keeping donepezil as the reference compound. The IC_{50} values of the molecules are summarized in Table 1. AChE inhibition remains a very critical end point in slowing down the progression of the disease, and thus we selected to determine the ability of our compounds to inhibit the aforesaid enzyme.

2.3.1 Structure activity relationship. Our target molecule contains triazole and pyridoxine moieties incorporated into it. Before starting our investigation on our library of compounds $5(a-q)$, we sought to find out the inhibitory activity of pyridoxine on AChE and observed it to be inactive against the enzyme; these results correlated well with the studies of Drtinova et al.¹⁸ We also sought to find out the effect of triazole moiety and substitution on the aromatic ring on enzymatic activity.

The unsubstituted 5a did not show much inhibition even at the highest concentration (10 mM), thus can be considered

inactive towards the enzyme. With the substitution of methyl group at the para position (5b) it however, showed inhibitory activity as compared to 5a. By changing the position of methyl group from para to meta (5c) and ortho (5d) position, a marked change was noted. A *meta* substitution of the electron donating group (EDG) offered a better inhibitory activity almost twice potent as compared to 5b, while the ortho alkyl substitution proved even better. Substitution of methoxy group at the $3rd$ position (5e) and at $3rd$ and $5th$ position of the aromatic ring (5f) had no inhibitory activity, however, methyl substitution proved better than these. A double methoxy substitution was found more efficacious compared with its single substitution.

We further explored by trying different halides as substituents and their effect on the inhibition on the enzyme. We also synthesized F, Cl and Br derivatives at different position. Halogens are supposed to have +R effect on the aromatic ring, despite of their position. When the effect of halogen is observed at the *para* position, fluoro substituent $(5g)$ showed lesser efficacy compared to $-Cl(5j)$ and $-Br(5m)$ derivatives. However, we did not find a uniform trend in activity as we moved down the periodic table. Chloro substitution at the *para* position was more potent in comparison to bromo substitution. However, meta substitution of halogens showed much better results as compared to the para substitution. Change in position from para to meta did not prove successful in improving potency in chloro substitution (5k), as there was no such change in the IC_{50} value. Bromine at the 3^{rd} position of the aromatic ring $(5n)$ was more active in comparison to the fluoro substitution (5h). We even tried to find out the effect of substitution when two positions on the aromatic ring are occupied with the halogen atoms. We thus included fluorine in $3rd$ and $5th$ position (5i) and in **EXACO Access Article is the constructed on 1990** and the comparison are the published on 09 July 2020. The comparison of the comparison of the common 2 weak creative Common 2020. The article is likely are the common and

Scheme 1 Reagents and conditions-(i) 2,2-dimethoxypropane, p-toluenesulfonic acid monohydrate, anhydrous acetone, under argon atmosphere 20 h, rt (ii) NaH, propargyl bromide, dry THF, reflux under argon atmosphere, 4 h (iii) benzyl bromide, NaN₃, triethylamine, CuSO₄·5H₂O (10 mol%), ascorbic acid (25 mol%), t-BuOH : H₂O (1 : 1), reflux for 6-12 h (iv) THF, 10% conc. sulphuric acid, 70 °C, 6-12 h.

Table 1 Results of AChE inhibition and anti-oxidant values

 $a_{IC_{50}}$ values is the concentration of the enzyme required to decrease the enzyme activity by 50% and are the mean of three independent experiments, represented in mean \pm SEM (SEM = standard error mean). ^b The mean \pm SEM of three independent experiments. Data are represented as ORAC-FL values of Trolox equivalents (μ M of tested compound/ μ M of Trolox). ^c n.a = not active. ^d N.T = not tested.

another compound, chlorine at $3rd$ and $4th$ position (5l). Inclusion of two fluorine atoms at both the *meta* positions, did not show change in potency but there was a spurt in % of enzyme inhibition. To summarize the effect of di-substitution on the aromatic ring, 5i was considered the best as compared to 5l and 5f. Owing to the fact that fluorine and methoxy groups are both ring activators, the difference in potency and percentage of

inhibition between the two can be due to the high electronegativity and small size of fluorine.

We tried with trifluoro-methyl substitution at para, meta and $ortho$ positions (50, 5p, 5q). Trifluoro-methyl substitution at the ortho position was the best amongst the three both from the perspective of potency and inhibitory activity. However, when compared with methyl substitution, 5b was double potent in

Table 2 Percentage of AChE inhibition of 5c, 5d, 5h, 5i, 5n, 5q and donepezil at different concentrations of the compounds^a

	$156 \mu M$		312 µM		$625 \mu M$		1.25 mM		$2.5 \text{ }\mathrm{mM}$		5 mM		10 mM	
	$\%$	\pm SEM	$\%$	\pm SEM	$\%$	\pm SEM	$\%$	\pm SEM	$\%$	\pm SEM	$\%$	\pm SEM	$\%$	\pm SEM
5с	22.49	3.62	26.92	0.55	39.11	2.48	53.8	3.13	64.11	1.79	73.43	1.13	81.92	0.85
5d	33.53	6.44	36.41	6.8	51.15	5.55	58.56	$\overline{4}$	69.57	2.3	81.3	2.36	86.03	0.63
5h	19.04	0.29	25.49	2.22	35.1	1.59	56.37	1.85	60.48	1.24	66.56	0.26	69.02	0.36
5i	14.97	4.12	26.58	3.92	35.38	0.47	43	1.52	61.41	2.82	66.78	$0.1\,$	77.89	0.15
5n	14.69	1.15	25.79	2.73	41.99	1.42	57.39	4.99	70.16	4.11	75.88	2.44	86.28	1.38
5q	22.49	3.62	26.92	0.55	39.11	2.48	53.8	3.13	64.11	1.79	73.43	1.13	81.92	0.85
Donepezil	89.07	1.19	94.5	0.81	97.44	0.73	100.88	0.93	99.7	1.04	99.74	0.69	99.95	0.48
								2.4 Antioxidant study Aging is associated with oxidative stress resulting in generation of reactive oxygen species. We determined the oxygen radical absorption capacity-fluoroscein (ORAC-FL) values for the entire						
								series. The ORAC-FL values are given in Table 1 and are a measure of the total antioxidant capacity. We generated peroxyl						
								free radicals using the free radical generator 2,2-azobis(2-						
								amidino-propane)dihydrochloride (AAPH) and fluorescein as						
								the fluorescent probe in this assay at 37 °C. The total antioxidant						
								capacity of the molecules were determined in equivalents to						
								Trolox (vitamin E analogue), keeping both at same concentration						
respect to 50. We got the same observation for meta substitution as well, 5c was twice in terms of potency as compared to 5p. Methyl substitution at the ortho position was marginally better in terms of trifluoro-methyl substitution at the same position. Methyl group is considered as ring activator, while trifluoro- methyl group as ring deactivator. Therefore, the observations confirm that EDG is best suited for inhibitory activity against AChE, when pyridoxine based triazoles are concerned. Table 2 gives a detailed information of percentage of inhibition of AChE at different concentrations of by 5c, 5d, 5h, 5i, 5n, 5q and the reference compound, donepezil. Fig. 3 shows the comparative results of percentage of inhibition of AChE by 5c, 5d, 5h, 5i, 5n,								(8 μ M). One of the reasons for choosing pyridoxine as a phar-						
5q derivatives at 10 mM concentration, taking donepezil as the								macophore was for it's antioxidant nature. We observed all our						
reference standard.								compounds to have anti-oxidant property as shown in Table 1.						
		To summarize all our observations with respect to our stan-						Vitamin B_6 being a dietary supplement was tested using the						
dard compound donepezil, our molecules required a higher														
concentration to show their inhibitory activity. Donepezil showed								ORAC-FL method. All the compounds showed almost similar						
almost 89% inhibition at the highest dilution in which we tested								anti-oxidant nature, however, pyridoxine showed the most effi-						
our compounds (156 μ M) and 65% inhibition at 5 μ M. However,								cacy in terms of an anti-oxidant. Pyridoxine was taken as a refer- ence compound to evaluate the entire series in this experiment.						

To summarize all our observations with respect to our standard compound donepezil, our molecules required a higher concentration to show their inhibitory activity. Donepezil showed almost 89% inhibition at the highest dilution in which we tested our compounds (156 μ M) and 65% inhibition at 5 μ M. However, at the highest concentration, that is, at 10 mM concentration, compounds showed a maximum of 86% inhibitory activity towards AChE while donepezil exhibited 100% activity.

Fig. 3 Percentage of inhibition of AChE by 5c, 5d, 5h, 5i, 5n, 5q derivatives at 10 mM concentration. Data presented here is mean \pm SEM of three independent experiments.

2.4 Antioxidant study

Therefore, from the observations of Table 1, we selected 5c, 5d, 5h, 5i, 5n and 5q on the basis of AChE inhibitory activity and antioxidant property of the compounds for further evaluations.

2.5 Metal chelation study

Chelation of bio-metals will make the compounds prove as MTDLs and be an added advantage. We tested 5c, 5d, 5h, 5i, 5n and 5q against bio-metals Fe³⁺, Cu²⁺, Zn²⁺, Al³⁺ which are responsible for AD, using UV spectrophotometer within the wavelength range of 200 to 600 nm. All these bio-metals are responsible for the pathophysiological conditions in AD. We found a marked ability within the compounds to chelate $Fe³⁺$. Compounds 5c, 5i and 5q chelated both Fe^{3+} and Al^{3+} , 5i chelated the metals most with a 222% increase in absorbance at 292 nm which also includes a bathochromic shift of about 3 nm in case of Fe^{3+} and an increase of 104% in absorbance at 292 nm which too included a bathochromic shift of 4 nm in case of Al^{3+} . The observations for all the compounds studied for metal chelation a study has been tabulated in Table 3.

Although, aluminium being the most abundant metal on the earth's crust, it do not pose an immediate health concern when being exposed to the human body. Chronic exposure to aluminium in drinking water has been studied for it's association with AD. To add more to the pathophysiology, aluminium is known to enhance ROS generation and stimulated iron based generation of ROS species. Both these metals are also known to be associated with the senile plaques.¹⁹ Therefore, 5i has the ability to chelate both the trivalent cations which can add well to the property of the compound as a MTDL as shown in Fig. 4a.

The main challenge for chelation therapy of metals is selectivity towards metal cations. So we sought to check the selectivity towards essential metals responsible for proper physiological functions of our body. We included Ca^{2+} , Mg²⁺, $Na⁺$ and $K⁺$ in our study and interestingly found out that 5i did not chelate these metal ions at all (Fig. 4a). The spectrum of 5i– $Fe³⁺$ complex was quite different from the parent spectrum and those by other metals. Owing to the fact, that our biological system comprise a mixture of metal ions, we sought to check the affinity of our compound 5i towards $Fe³⁺$ in the presence of other metals (Fe $^{3+}$, Zn $^{2+}$, Cu $^{2+}$, Ca $^{2+}$, Mg $^{2+}$, Na $^+$ and K $^+$) present in equal amount. However, we excluded Al^{3+} from the study as the purpose was to check affinity towards Fe^{3+} and Al^{3+} does not form an essential part of our system. We observed the characteristic curve of $5i-Fe^{3+}$ as $5i$ showed affinity towards Fe^{3+} in the metal pool (Fig. 4b).

We studied the stoichiometric association of the $Fe³⁺$ -5i complex by keeping the amount of ligand fixed and titrating the amount of metal in increasing concentration ranging from 0– 100μ M using UV spectrophotometer. The spectral changes were recorded at nm and a graph was plotted (Fig. 4c). There was an increase in intensity at 293 nm and a decrease at 327 nm, on successive increase in concentration of $Fe³⁺$. We analysed the data and plotted the Job's plot, at 239 nm and 327 nm where on extrapolation we got a mole fraction of 0.5 in both the wavelength confirming formation of $1:1$ stoichiometric complex between 5i and $Fe³⁺$ (Fig. 4d and e).

Good et al. performed a laser microprobe study or the LAMMA study where it was found selective deposition of Al and Fe in the neurofibrillary tangles.²⁰ Therefore, 5i being a selective aluminium and iron chelator can be a potential inhibitor against these potentially toxic metals and can surely halt the mechanism of pathological important structures.

^a Data represented here is percentage change in absorbance; (+) indicating a decrease in absorbance and (-) as increase; calculated by the formulation in the formulation h [(abs of tested compound – abs of complex)/abs of tested compound] \times 100. b Data represented here is shift in nm; (+) indicates a hypsochromic shift and (-) indicates a bathochromic shift as calculated from the formula [λ_{max} of tested compound $-\lambda_{\text{max}}$ of complex].

Fig. 4 (a) Metal chelation spectra of 5i. (b) Selectivity spectrum of 5i from metal pool comprising of Fe³⁺, Zn²⁺, Cu²⁺, Ca²⁺, Mg²⁺, Na⁺ and K⁺; 5i formed complex with Fe³⁺ as identified from it's characteristic curve shown in (a). (c) Titration spectra of Fe³⁺ with 5i with 20 µL aliquots of metal solution to in increase the concentration of Fe³⁺ from 0-100 μ M [a-k], with successive increment of 10 μ M. (d) Job's plot of 5i at 293 nm. (e) Job's plot of 5i at 327 nm.

2.6 In silico study

2.6.1 Molecular docking study. We investigated on the binding mode of pyridoxine based triazole derivatives with AChE. We studied the binding interaction of 5i with AChE using molecular docking in Schrodinger glide. The Fig. 5 represents the molecular interactions of 5i with the active site amino acid residues of the enzyme AChE. The molecular docking was done with recombinant human AChE (PDB ID 4EY5) and it's 2D representation has been shown in Fig. 5.²¹ The interactions mainly observed were with the peripheral anionic site of the enzyme (PAS site) with hydrophobic amino acids Tyr 124 and Phe 338 having π – π interactions with triazole nucleus of 5i. On

the other hand, pyridine nucleus of the pyridoxine moiety was seen to have $\pi-\pi$ interaction with hydrophobic Trp 286. The principal amino acid of PAS, that is, Trp 86, was found in the vicinity of the ligand.²² The hydroxyl group of the pyridoxine was found to form hydrogen bond with Arg 296. The catalytic anionic site (CAS) amino acids Ser 203 and His 447 were found in close proximity of the ligand. The dock score of 5i calculated from the molecular docking study was -8.50 kcal J⁻¹.

The ligand showed interactions with the PAS site amino acids of AChE and binding to the PAS site of the enzyme is known to inhibit A β aggregation due to the homologous domains of the β -neurexin with PAS site, itself serving as

a ligand. Thus in silico results of molecular docking with ligand validated the results obtained from the in vitro AChE experiments.

2.6.2 Predictive ADME properties of 5i by computational methods. Taking the help of SwissADME (http:// www.swissadme.ch), we predicted the drug-like properties, pharmacokinetic and physicochemical properties of 5i. This open access web tool comprised proficient methods of predictive tools like BOILED EGG, 23 Bioavailability Radar. The software presents an early estimation of these ADME parameters (for Absorption, Distribution, Metabolism and Excretion) and provides sufficient light to the physicochemical parameters which is needed to be improved during formulation. The in silico result predicted by the aforesaid software is presented in Fig. 6a and b. The molecule 5i was found to possess good drug likeness properties and was well within the Bioavailability Radar. The Bioavailability Radar gives a first glance of the druglikeness of the molecule. The spider web like structure given inside Fig. 6a is the Bioavailability Radar. The pink region is the optimal range for every property. Daina et al. explained well about the web tool for prediction of drug likeness properties. All the predicted properties of the compound 5i by the web tool have been put up in Fig. 6a.²⁴

Fig. 6b is the BOILED EGG representation of the permeation properties of the molecule 5i. The figure displays two parts, the white part representing those set of values which allows gastric permeation and the yolk or the yellow region representing those set of values which allows blood–brain-barrier permeation. The graph is plotted against W log $P[a \log P(n\text{-octanol/water parti-})]$ tion coefficient) method developed by Wildman and Crippen] versus TPSA or Topological Polar Surface Area in the BOILED EGG representation. The blue dot in the figure written as Molecule 1 is of 5i.

2.6.3 Toxicity prediction of 5i by computational methods. The in silico toxicity prediction can reduce the time, cost and the number of animal experiments. We used the ProTox software for this in silico study, which identifies the LD_{50} value of the median lethal dose of a drug after analysing the similarity of the compounds and also identifies the toxic fragments.²⁵ It is an open access web tool (http://www.tox.charite.de/tox), which displays strong results at (sensitivity, specificity and precision of 76, 95 and 75%, respectively) which are validated by diverse external set. Fig. 7 displays the oral toxicity prediction results which show the molecule 5i to have a predicted LD_{50} of 3350 mg kg^{-1} , falling under the Class 5 of toxicity of chemical classification. The molecule is predicted to be inactive in case of hepatotoxicity, carcinogenicity, mutagenicity, cytotoxicity, all

Fig. 6 (a) Predicted ADME properties of 5i as obtained from the report generated by SwissADME software. (b) BOILED EGG representation of 5i (denoted as Molecule 1 and represented by blue dot in the figure). From the BOILED EGG representation, the yolk denotes the points which allows passive blood–brain-barrier permeation, while the white region denotes points for passive gastrointestinal tract absorption. The graph is plotted against W log P[a log P(n-octanol/water partition coefficient) method developed by Wildman and Crippen] versus TPSA or Topological Polar Surface Area.

Fig. 7 Report of toxicity prediction as generated by ProTox-II online tool.

nuclear signalling pathways and stress response pathways. The predicted LD_{50} value of 5i is well in correlation with the fact that pyridoxine itself rarely causes toxicity when given in very high doses. The in silico results generated gives us intuitive idea that 5i can be considered as a much safer molecule.

3 Conclusion

Here, we report the design, synthesis and biological evaluations of a novel class of pyridoxine based triazole as multi-target directed ligands. Pyridoxine itself is very crucial for the homocysteine balance in AD patients and as well as considered an important vitamin for our daily diet. However, as proved from our experiment, it is not an AChE inhibitor. Out of the seventeen synthesized pyridoxine based triazoles, six showed good AChE inhibition and antioxidant potency. Meta and ortho substitution on the aromatic ring with EDG or ring activator proved beneficiary for AChE activity. With further metal chelation studies, we found 5i as the best compound of this new series. 5i has good AChE inhibitory activity (IC₅₀ = 1.56 ± 0.02) mM) with 77.89% of inhibition at highest concentration and possessed antioxidant property having ORAC-FL value of 1.21 \pm 0.28 equivalent to Trolox, whose ORAC-FL value is taken as 1. The *in vitro* results for 5i has been validated with the help of molecular docking with AChE by investigating it's binding poses. The compound 5i chelated $Fe³⁺$ as the spectra showed about 220% increase in absorbance and a 104% increase was seen in case of Al^{3+} , exhibiting bathochromic shift in both. Selectivity towards targeted metal ion has always been

a challenge in metal chelation therapeutics, henceforth, further throwing light on the ability of $5i$ to selectively chelate $Fe³⁺$ from a pool of metal ions, gives it an added advantage. None of the literature of triazole based anti-Alzheimer's agents have reported selectivity towards metal ions with a drastic change in absorbance. We concluded our work with in silico studies of drug likeness, pharmacokinetic properties of 5i, along with toxicity predictions which predicted it to be under Class 5 according to the toxicity labelling of chemicals. Our initial ndings can open up new therapeutic areas in multi-target directed ligands against AD.

4 Experimental section

4.1 Chemistry

4.1.1 General remarks. All chemical reagents and solvents were purchased from Sigma Aldrich, Spectrochem Pvt. Ltd, Alfa Aesar and used without further purification. Reactions were monitored by Thin Layer Chromatography (TLC) using Merck silica gel 60 F254 pre-coated plates (0.25 mm) and the compounds were examined under UV chamber (Ultra violet fluorescence analysis cabinet-PSI). Column chromatography was performed with silica gel (100–200 mesh). All products were characterized by $\rm ^1H$ NMR, $\rm ^{13}C$ NMR and IR spectroscopy. $\rm ^1H,$ $\rm ^{19}F$ and 13C NMR were recorded on JEOL RESONANCE ECZ500R at 500 MHz, 470 MHz and 125 MHz respectively with chemical shift (δ) expressed in parts per million in CDCl₃ at 27 °C downfield from TMS as an internal standard and coupling in Hertz (Hz). Splitting patterns are described as singlet (s),

doublet (d), triplet (t), multiplet (m), doublet of doublet (dd) and doublet of triplet (dt). For 1 H NMR, 13 C NMR reference for CDCl₃ appeared at 7.26 and 77.367 ppm respectively. For 1 H NMR, ¹³C NMR reference for DMSO- d_6 appeared at 2.50 ppm and 39.52 ppm respectively. IR spectra were recorded on a BRUKER ECO-ATR Spectrometer. HRMS was measured on LC-MS by Agilent Technologies LC-MS and purity check was done using HPLC on a HPLC by Agilent Technologies. The retention time in HPLC is represented as t_R and calculated in minutes (min).

4.1.2 General procedure for the acetonide protection of (5 hydroxy-6-methylpyridine-3,4-diyl)dimethanol. To a suspension of pyridoxine hydrochloride (5.00 g, 24.3 mmol) in dry acetone (75 mL) under argon atmosphere, added 2,2-dimethoxypropane $(50.1$ mL, 408 mmol) and stirred for 10 min.²⁶ Following this, ptoluenesulfonic acid monohydrate (18.5 g, 97.2 mmol) was added and the reaction was stirred in room temperature for 20 hours. Completion of the reaction was monitored by TLC. The reaction mixture was neutralized with aqueous sodium bicarbonate, concentrated under reduced pressure, and the dark brown solution was extracted with dichloromethane (40 mL \times 3). The organic phase was washed with brine and dried over anhydrous $Na₂SO₄$. The DCM layer was concentrated in vacuum to provide a crude product, which was washed with hexane to give compound 2 as white crystals in 86% yield. Paper

Vece Metallion (i.e., the May 121 Mat therefore for a persistent of the presence of the Social Access Article is licensed under the Society Articles. By the Society Articles Commons Articles Articles In the Society

4.1.2.1 (2,2,8-Trimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-yl) methanol (2). White solid crystals; mp-110-113 °C; $R_f = 0.5$ (ethyl acetate/hexane 3 : 2); ¹H NMR (500 MHz, chloroform-*d*) δ 7.77 (s, 1H, N=CH) δ 4.91 (s, 2H, O–CH₂) δ 4.52 (s, 2H, CH₂–OH) δ 2.34 (s, 3H, CH₃) δ 1.53 (s, 6H, 2 \times CH₃); ¹³C NMR (125 MHz, chloroform-d) d 145.44, 145.09, 138.42, 130.59, 125.24, 99.41, 58.24, 57.88, 24.53, 18.24; IR $\text{(cm}^{-1}\text{)}$ v: 3851.17, 3619.70, 1695.98, 1213.27.

4.1.3 General procedure for synthesis of 2,2,8-trimethyl-5- $((prop-2-yn-1-yloxy)$ methyl $)-4H-[1,3]$ dioxino $[4,5-c]$ pyridine. To a suspension of sodium hydride (1.5 eq.) in THF (40 mL), added (2,2,8-trimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-yl)methanol (2) $(1.00 \text{ g}, 4.7 \text{ mmol})$ under argon atmosphere and refluxed for 30 min. Following this, propargyl bromide (2 eq.) was added dropwise for 1 h and the reaction mixture was refluxed for 4 h. Completion of the reaction was monitored by TLC. The reaction mixture was concentrated under reduced pressure, quenched with saturated ammonium chloride solution and extracted with dichloromethane (40 mL \times 3). The organic phase was washed with brine and dried over anhydrous $Na₂SO₄$. The DCM layer was concentrated in vacuum to provide a crude product, which was purified by column on silica using 20 : 80 ratio of hexane-DCM solvent system to provide compound 3 in 75% yield.

4.1.3.1 2,2,8-Trimethyl-5-((prop-2-yn-1-yloxy)methyl)-4H-[1,3] dioxino[4,5-c]pyridine (3). Brown oil; $R_f = 0.6$ (DCM/methanol 12 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.99 (s, 1H, N= CH) δ 4.9 (s, 2H, O–CH₂) δ 4.71 (s, 2H, CH₂–OH) δ 4.46 (s, 2H, CH₂-C) δ 2.4 (s, 3H, C<u>H₃</u>) δ 1.5 (s, 6H, 2 \times C<u>H₃</u>); IR (cm⁻¹) ν : 3292.03, 2992.75, 2931.93, 2858.50, 2114.38, 1067.29, 1025.50.

4.1.4 General procedure for synthesis of substituted 5-(((1 benzyl-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2,2,8-trimethyl- $4H-[1,3]$ dioxino $[4,5-c]$ pyridine $4(a-q)$. Different derivatives were

prepared via the reaction of benzyl bromides (1.1 mmol) with sodium azide (1 mmol) in the presence of triethylamine in t-BuOH/water $(1:1)$. After around 30 min, 2,2,8-trimethyl-5- $((prop-2-yn-1-yloxy)methyl)-4H-[1,3]dioxino[4,5-c]pyridine (3) (1$ mmol) in absolute *t*-BuOH (1 mL) was added to the solution in the presence of $CuSO₄·5H₂O$ (10 mol%) and freshly prepared ascorbic acid solution (25 mol%), at 50 $^{\circ}$ C and stirred at 70 $^{\circ}$ C for 12 h. After completion of the reaction (checked by using TLC), the reaction mixture was passed through a slurry of Celite in ethyl acetate. The filtrate was concentrated under reduced pressure. The organic residue was extracted with EtOAc (20 mL \times 3). The organic phase was washed with brine and dried over anhydrous $Na₂SO₄$. The solvent was evaporated to dryness under reduced pressure to afford compounds 4(a–q). The compounds were finally washed with hexane.

4.1.5 General procedure for deprotection of compounds $4(a-q)$. The corresponding compounds $4(a-q)$ (0.60–0.65 mmol) was dissolved in THF (5 mL), and 10% HCl solution (4 mL) was added. The reaction mixture was refluxed for 6-11 h under an argon atmosphere. After complete reaction, THF was evaporated under reduced pressure. The residue was neutralized with aqueous saturated sodium bicarbonate solution, and extracted with EtOAc (20 mL \times 3). The organic phase was washed with brine and dried over anhydrous $Na₂SO₄$. The solvent was evaporated to dryness under reduced pressure to 5(a–q) in 85– 96% yield. The compounds were finally washed with hexane.

4.1.5.1 5-(((1-Benzyl-1H-1,2,3-triazol-4-yl)methoxy)methyl)-4- (hydroxymethyl)-2-methylpyridin-3-ol (5a). Brown oil; yield 95%; $R_{\rm f}$ = 0.4 (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroformd) δ 7.75 (s, 1H, N=CH) δ 7.39 (s, 1H, C=CH–N) δ 7.36–7.31 (m, 3H, Ar–H) δ 7.21 (dd, J = 7.2, 2.2, 2H, Ar–H) δ 5.45 (s, 2H, N–CH₂) δ 4.95 (s, 2H, CH₂-O) δ 4.43 (s, 2H, CH₂-OH) δ 4.43 (s, 2H, CH₂-O) δ 2.40 (s, 3H, N–C–CH₃); ¹³C NMR (125 MHz, chloroform-*d*) d 152.05, 148.52, 144.80, 139.90, 134.26, 131.57, 129.28, 128.21, 127.54, 122.73, 68.25, 62.28, 59.77, 54.38, 18.67; IR $\text{(cm}^{-1}\text{)}$ v: 3141.08, 2926.77, 2863.95, 2363.77, 2246.90, 1615.93, 1225.02, 1213.27; HRMS (m/z) : calculated for C₁₈H₂₀N₄O₃ [M + H]⁺ 341.1691; found, 341.1611; HPLC purity-99%; $t_R = 9.86$ min.

4.1.5.2 4-(Hydroxymethyl)-2-methyl-5-(((1-(4-methylbenzyl)- 1H-1,2,3-triazol-4-yl)methoxy)methyl)pyridin-3-ol (5b). Yellow sticky solid; yield 88%; $R_{\rm f} = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.83 (s, 1H, N=CH) δ 7.35 (s, 1H, C= CLN δ = 7.14 (q, J = 8.1, 4H, Ar– H) δ 5.42 (s, 2H, N– CH_2) δ 4.97 (s, 2H, CH₂-OH) δ 4.52 (s, 2H, CH₂-O) δ 4.42 (s, 2H, CH₂-O) δ 2.45 (s, 3HN–C–C \underline{H}_3), δ 2.33 (s, 3H, Ar–C \underline{H}_3); ¹³C NMR (125 MHz, chloroform-d) δ 144.76, 140.56, 139.15, 134.72, 131.53, 131.29, 130.05, 129.94, 128.28, 122.42, 68.35, 61.47, 59.67, 54.22, 21.23, 18.85. IR (cm^{-1}) v: 3145.19, 2930.77, 2869.75, 2367.37, 2248.92, 1620.39, 1221.03, 1210.28; HRMS (m/z): calculated for C₁₉H₂₃N₄O₃ [M + H]⁺ 355.1770; found, 355.1772; HPLC purity-97%; $t_R = 11.20$ min.

4.1.5.3 4-(Hydroxymethyl)-2-methyl-5-(((1-(3-methylbenzyl)- 1H-1,2,3-triazol-4-yl)methoxy)methyl)pyridin-3-ol (5c). Yellow oil; yield 87%; $R_{\rm f} = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.83 (s, 1H, N=CH) δ 7.37 (s, 1H, C=CH–N) δ 7.24–7.21 (m, 1H, Ar–H) δ = 7.16 (d, J = 7.6 Hz, 1H, Ar–H) δ 7.05–7.02 (m, 2H, Ar–H) δ 5.42 (s, 2H, N–CH₂) δ 4.97 (s, 2H,

CH₂-OH) δ 4.52 (s, 2H, CH₂-O) δ 4.43 (s, 2H, CH₂-O) δ 2.45 (s, 3H, N–C–C H_3), δ 2.32 (s, 3H, Ar–C H_3); ¹³C NMR (125 MHz, chloroform-d) d 140.38, 139.18, 134.11, 131.64, 129.78, 129.16, 128.97, 127.45, 125.67, 125.31, 122.58, 68.34, 61.68, 59.60, 54.43, 21.39, 18.75. IR (cm^{-1}) v: 3146.20, 2931.47, 2868.75, 2369.58, 2249.22, 1622.19, 1223.10, 1211.32; HRMS (m/z): calculated for C₁₉H₂₃N₄O₃ [M + H]⁺ 355.1770; found, 355.1768; HPLC purity-96%; $t_R = 10.30$ min.

4.1.5.4 4-(Hydroxymethyl)-2-methyl-5-(((1-(2-methylbenzyl)-

1H-1,2,3-triazol-4-yl)methoxy)methyl)pyridin-3-ol (5d). Darkgreen semisolid; yield 93%; $R_f = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.83 (s, 1H, N=C<u>H</u>) δ 7.29 (s, 1H, C=CH–N) δ = 7.30–7.25 (m, 1H, Ar–H), δ 7.20 (dd, J = 7.2, 3.6 Hz, 2H, Ar- \underline{H}), δ 7.13 (d, $J = 8.0$, 1H, Ar- \underline{H}) δ 5.48 (s, 2H, N–CH₂) δ 4.97 (s, 2H, CH₂–OH) δ 4.52 (s, 2H, CH₂–O) δ 4.42 (s, 2H, CH₂–O) δ 2.45 (s, 3H, N–C–CH₃), δ 2.24 (s, 3H, Ar–CH₃); ¹³C NMR (125 MHz, chloroform-d) d 148.80, 143.86, 137.03, 132.11, 131.21, 129.61, 129.44, 126.83, 68.39, 61.62, 59.57, 52.56, 19.04; $\text{IR (cm}^{-1})$ v: 3147.20, 2930.70, 2870.59, 2372.85, 2247.31, 1620.20, 1221.70, 1215.36; HRMS (m/z): calculated for $C_{19}H_{23}N_4O_3$ [M + H]⁺ 355.1770; found, 355.1774; HPLC purity-95%; $t_{\rm R} = 10.92$ min.

4.1.5.5 4-(Hydroxymethyl)-5-(((1-(3-methoxybenzyl)-1H-1,2,3 triazol-4-yl)methoxy)methyl)-2-methylpyridin-3-ol (5e). Yellow sticky solid; yield 90%; $R_{\rm f}$ = 0.4 (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.78 (s, 1H, N=CH) δ 7.40 (s, 1H, C= CH–N) δ 7.27 (d, J = 7.6 Hz, 1H, Ar–H) δ 6.86 (dd, J = 8.3, 2.5 Hz, 1H, Ar–H) δ 6.80 (d, J = 7.5 Hz, 1H, Ar–H) δ 6.74 (d, J = 1.6 Hz, 1H, Ar- \underline{H}) δ 5.42 (s, 2H, N-C \underline{H}_2) δ 4.96 (s, 2H, C \underline{H}_2 -OH) δ 4.46 (s, 2H, CH₂-O) δ 4.43 (s, 2H, CH₂-O) δ 3.75 (s, 3H, Ar-O-CH₃), δ 2.42 (s, 3H, N–C–CH₃); ¹³C NMR (125 MHz, chloroform-*d* and DMSO-d₆) δ 156.30, 149.66, 144.88, 140.99, 134.97, 127.76, 124.99, 118.81, 118.55, 72.91, 68.02, 64.64, 60.09, 58.73, 23.86; $\text{IR (cm}^{-1})$ v: 3248.36, 2932.56, 2873.62, 2375.56, 2248.32, 1620.20, 1231.80, 1218.36; HRMS (m/z): calculated for $C_{19}H_{23}N_4O_4$ [M + H]⁺ 371.1719; found, 371.1719; HPLC purity-96%; $t_{\rm R} = 10.24$ min.

4.1.5.6 5-(((1-(3,5-Dimethoxybenzyl)-1H-1,2,3-triazol-4-yl) methoxy)methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5f). Yellow oil; yield 85%; $R_{\rm f} = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.83 (s, 1H, N=CH) δ 7.41 (s, 1H, C=

CH–N) δ 7.25 (s, 1H, Ar–H) δ 6.40 (s, 1H, Ar–H), δ 6.36 (s, 1H, Ar– H), δ 5.39 (s, 2H, N–CH₂) δ 4.96 (s, 2H, CH₂–OH) δ 4.50 (s, 2H, CH₂–O) δ 4.44 (s, 2H, CH₂–O) δ 3.74 (s, 6H, Ar–O–CH₃), δ 2.44 (s, 3H, N–C–C H_3); ¹³C ¹³C NMR (125 MHz, chloroform-d and DMSO-d6) d 161.43, 136.37, 127.44, 122.72, 106.32, 100.48, 68.30, 61.88, 59.74, 55.54, 54.41, 18.79.; IR $\rm (cm^{-1})$ $\rm \nu$: 3245.63, 2893.60, 2892.36, 2376.78, 2369.32, 1623.20, 1241.81, 1278.23; HRMS (m/z) : calculated for C₂₀H₂₅N₄O₅ [M + H]⁺ 401.1825; found, 401.1834; HPLC purity-97%; $t_R = 10.72$ min.

4.1.5.7 5-(((1-(4-Fluorobenzyl)-1H-1,2,3-triazol-4-yl)methoxy) methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5g). Brown oil; yield 93%; $R_{\rm f} = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.82 (s, 1H, N=CH) δ 7.38 (s, 1H, C=CH-N) δ 7.24–7.20 (m, 2H, Ar–H), δ 7.03 (t, J = 8.6 Hz, 2H, Ar–H) δ 5.44 (s, 2H, N-C \underline{H}_2) δ 4.98 (s, 2H, C \underline{H}_2 -OH) δ 4.51 (s, 2H, C \underline{H}_2 -O) δ 4.44 (s, 2H, CH₂–O) δ 2.44 (s, 3H, N–C–CH₃); ¹³C NMR (125

MHz, chloroform-d) δ 148.67, 145.34, 140.30, 131.58, 130.16, 130.09, 122.49, 116.41, 116.24, 68.38, 61.78, 59.64, 53.64, 18.75; ¹⁹F NMR (470 MHz, chloroform-d) δ -112.10 to -112.17 (m); IR $\text{(cm}^{-1}\text{)}$ v: 3247.63, 2896.70, 2912.86, 2378.87, 2367.85, 1613.10, 1245.81, 1218.23; HRMS (m/z) : calculated for C₁₈H₂₀N₄O₃F [M + H ⁺ 359.1519; found, 359.1514; HPLC purity-99.8%; t_R = 10.32 min.

4.1.5.8 5-(((1-(3-Fluorobenzyl)-1H-1,2,3-triazol-4-yl)methoxy) methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5h). Yellow oil; yield 86%; $R_{\rm f} = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.81 (s, 1H, N=CH) δ 7.42 (s, 1H, C=CH–N) δ 7.33 (td, $J = 8.0$, 5.9 Hz, 1H, Ar- \underline{H}), δ 7.06-6.99 (m, 2H, Ar- \underline{H}) δ 6.92 (d, J = 9.1 Hz, 1H, Ar–H) δ 5.47 (s, 2H, N–CH₂) δ 4.98 (s, 2H, CH₂-OH) δ 4.51 (s, 2H, CH₂-O) δ 4.45 (s, 2H, CH₂-O) δ 2.44 (s, 3H, N-C-C \underline{H}_3); ¹³C NMR (125 MHz, chloroform-d) δ 151.98, 148.68, 145.06, 140.25, 131.55, 131.00, 130.94, 127.45, 123.65, 122.69, 116.15, 115.99, 115.24, 115.07, 68.37, 61.86, 59.69, 53.72, 18.73; ¹⁹F NMR (470 MHz, chloroform-d) δ -111.20 (ddd, $J = 9.0, 6.3, 2.2$ Hz); IR (cm^{-1}) v: 3245.63, 2893.70, 2911.86, 2375.77, 2377.65, 1619.20, 1286.81, 1220.23; HRMS (m/z): calculated for $C_{18}H_{20}N_4O_3F$ [M + H]⁺ 359.1519; found, 359.1519; HPLC purity-98%; $t_R = 10.30$ min. Open Access Article. Published on 09 July 2020. Downloaded on 5/4/2024 6:52:27 PM. This article is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported Licence.](http://creativecommons.org/licenses/by-nc/3.0/) **[View Article Online](https://doi.org/10.1039/d0ra04942e)**

4.1.5.9 5-(((1-(3,5-Difluorobenzyl)-1H-1,2,3-triazol-4-yl) methoxy)methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5i). Yellow sticky solid; yield 96%; $R_f = 0.3$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d + DMSO-d₆) δ 7.87 (s, 1H, N= CH) δ 7.48 (s, 1H, C=CH–N) δ 6.85–6.79 (m, 3H, Ar–H), δ 5.56 (s, 2H, N–CH₂) δ 4.91 (s, 2H, CH₂–OH) δ 4.60 (s, 2H, CH₂–O) δ 4.52 (s, 2H, CH₂-O) δ 2.42 (s, 3H, N-C-CH₃); ¹³C NMR (125 MHz, chloroform-d and DMSO- d_6) δ 156.20, 149.88, 144.87, 143.80, 135.68, 132.50, 128.31, 115.92, 115.71, 108.78, 72.95, 68.04, 64.48, 57.54, 23.92; ¹⁹F NMR (470 MHz, chloroform-d) δ -103.42 to -103.51 (m) ; IR $\text{(cm}^{-1}\text{)}$ ν : 3240.35, 2885.91, 2900.62, 2378.85, 2378.63, 1622.31, 1278.78, 1223.73; HRMS (m/z): calculated for $C_{18}H_{19}N_4O_3F_2$ [M + H]⁺ 377.1425; found, 377.1422; HPLC purity-100%; $t_{\rm R} = 10.88$ min.

4.1.5.10 5-(((1-(4-Chlorobenzyl)-1H-1,2,3-triazol-4-yl) methoxy)methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5j). Yellow sticky solid; yield 85%; $R_f = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.80 (s, 1H, N=C<u>H</u>) δ 7.40 $(s, 1H, C=CH-N)$ δ 7.32 (d, J = 8.4 Hz, 2H, Ar–H), δ 7.16 (d, J = 8.3 Hz, 2H, Ar-H δ 5.43 (s, 2H, N-CH₂) δ 4.97 (s, 2H, CH₂-OH) δ 4.48 (s, 2H, CH₂-O) δ 4.44 (s, 2H, CH₂-O) δ 2.43 (s, 3H, N-C-CH₃); ¹³C NMR (125 MHz, chloroform-d) δ 148.53, 145.03, 139.96, 135.10, 132.76, 131.63, 129.53, 129.50, 127.57, 122.62, 68.32, 62.04, 59.72, 53.62, 29.78, 18.65; IR $\rm (cm^{-1})$ v: 3224.69, 2895.81, 2910.25, 2376.68, 2379.36, 1627.91, 1278.70, 1210.35; HRMS (m/z) : calculated for C₁₈H₂₀N₄O₃Cl [M + H]⁺ 375.1224; found, 375.1222; HPLC purity-97%; $t_R = 11.56$ min.

4.1.5.11 5-(((1-(3-Chlorobenzyl)-1H-1,2,3-triazol-4-yl) $methoxy)$ methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5k). Yellow oil; yield 88%; $R_{\rm f} = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.84 (s, 1H, N=CH) δ 7.42 (s, 1H, C= CH-N) δ 7.34-7.27 (m, 2H, Ar-H), δ 7.23 (s, 1H, Ar-H) δ 7.10 (d, J $=$ 7.2 Hz, 1H) δ 5.45 (s, 2H, N–C $_{\text{H}_2}$) δ 4.99 (s, 2H, C $_{\text{H}_2}$ –OH) δ 4.54 (s, 2H, CH₂–O) δ 4.45 (s, 2H, CH₂–O) δ 2.45 (s, 3H, N–C–CH₃); ¹³C NMR (125 MHz, chloroform-d) δ 151.92, 149.51, 148.78,

145.10, 130.60, 129.26, 128.25, 126.24, 122.65, 68.43, 61.68, 59.66, 53.67, 18.81; IR $\text{(cm}^{-1}\text{)}$ v: 3220.69, 2898.81, 2914.38, 2369.82, 2381.41, 1628.85, 1280.01, 1210.56; HRMS (m/z): calculated for $C_{18}H_{20}N_4O_3Cl$ $[M + H]^+$ 375.1229; found, 375.1225; HPLC purity-97%; $t_R = 10.32$ min.

4.1.5.12 5-(((1-(3,4-Dichlorobenzyl)-1H-1,2,3-triazol-4-yl)

methoxy)methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5l). Yellow oil; yield 92%; $R_{\rm f} = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform- d + DMSO- d_6) δ 7.75 (s, 1H, N=CH) δ 7.64 (s, 1H, C=CH–N) δ 7.24–7.20 (m, 2H, Ar–H), δ 7.06 (s, 1H, Ar-H) δ 5.40 (s, 2H, N-CH₂) δ 4.78 (s, 2H, CH₂-OH) δ 4.48 (s, 2H, CH₂–O) δ 4.38 (s, 2H, CH₂–O) δ 2.30 (s, 3H, N–C–CH₃); ¹³C NMR (125 MHz, chloroform-d and DMSO- d_6) δ 135.45, 131.07, 130.10, 127.62, 123.37, 68.19, 64.43, 59.76, 52.57, 19.27; IR $\text{(cm}^{-1}\text{)}$ v: 3225.72, 2890.23, 2926.83, 2370.82, 2380.62, 1629.85, 1282.01, 1217.61; HRMS (m/z) : calculated for C₁₈H₁₉N₄O₃Cl₂ [M + H]⁺ 410.0912; found, 410.0865; HPLC purity-98%; $t_R = 10.32$ min.

4.1.5.13 5-(((1-(4-Bromobenzyl)-1H-1,2,3-triazol-4-yl)methoxy) methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5m). Yellow sticky solid; yield 91%; $R_{\rm f}$ = 0.6 (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.84 (s, 1H, N=CH) δ 7.49 (d, J = 8.3 Hz, 2H, Ar-H δ 7.48 (s, 1H, C=CH-N) δ 7.11 (d, J = 8.3, 2H, Ar–H) δ 5.43 (s, 2H, N–CH₂) δ 4.98 (s, 2H, CH₂–OH) δ 4.53 (s, 2H, CH₂–O) δ 4.44 (s, 2H, CH₂–O) δ 2.45 (s, 3H, N–C–CH₃)¹³C NMR (125 MHz, chloroform-d) d 148.75, 133.25, 132.48, 129.80, 123.24, 122.56, 68.42, 61.70, 59.64, 53.69, 29.78; IR $\rm (cm^{-1})$ ν : 3226.24, 2892.83, 2924.81, 2374.22, 2386.92, 1685.23, 1280.81, 1216.56; HRMS (m/z) : calculated for C₁₈H₂₀N₄O₃Br [M + H]⁺ 419.0719; found, 419.0715; HPLC purity-99%; $t_R = 11.91$ min.

4.1.5.14 5-(((1-(3-Bromobenzyl)-1H-1,2,3-triazol-4-yl)methoxy) methyl)-4-(hydroxymethyl)-2-methylpyridin-3-ol (5n). Yellow oil; yield 91%; $R_{\rm f} = 0.6$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-d) δ 7.85 (s, 1H, N=CH) δ 7.48 (s, 1H, C=CH–N) δ 7.41 (d, J = 8.3 Hz, 2H, Ar–H) δ 7.23 (s, 1H, Ar–H) δ 7.15 (d, J = 7.7, 1H, Ar-H) δ 5.44 (s, 2H, N-CH₂) δ 4.98 (s, 2H, CH₂-OH) δ 4.53 (s, 2H, CH₂-O) δ 4.45 (s, 2H, CH₂-O) δ 2.45 (s, 3H, N-C-CH₃); ¹³C NMR (125 MHz, chloroform-d) δ 132.20, 131.15, 130.85, 126.72, 123.30, 117.51, 68.44, 61.74, 59.67, 53.60, 36.88 IR (cm^{-1}) v: 3225.20, 2890.63, 2923.72, 2373.15, 2388.85, 1685.57, 1282.74, 1210.78; HRMS (m/z): calculated for $C_{18}H_{20}N_4O_3Br[M+H]^+$ 419.0719; found, 419.0715; HPLC purity-95%; $t_{\rm R} = 11.79$ min.

4.1.5.15 4-(Hydroxymethyl)-2-methyl-5-(((1-(4-(tri uoromethyl)benzyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl) pyridin-3-ol (50). Yellow sticky solid; yield 95%; $R_f = 0.4$ (DCM/ methanol 19 : 1); ¹H NMR (500 MHz, chloroform-*d*) δ 7.80 (s,

1H, N=CH) δ 7.62 (d, J = 8.2 Hz, 2H, Ar-H) δ 7.43 (s, 1H, C= CH–N) 7.33 (d, $J = 8.1$, 2H, Ar–H) δ 5.53 (s, 2H, N–CH₂) δ 4.98 (s, 2H, CH₂-OH) δ 4.49 (s, 2H, CH₂-O) δ 4.45 (s, 2H, CH₂-O) δ 2.43 (s, 3H, N–C–C H_3); ¹³C NMR (125 MHz, chloroform-d) δ 151.97, 148.67, 145.22, 140.19, 138.21, 131.49, 128.35, 127.46, 126.26, 122.75, 68.40, 61.94, 59.72, 53.67, 18.74; 19F NMR (470 MHz, chloroform-d) δ -62.71 (s); IR $\rm (cm^{-1})$ v: 3115.52, 2895.96, 2925.63, 2375.38, 2390.58, 1182.06, 1683.57, 1281.54, 1213.61; HRMS (m/z) : calculated for C₁₉H₂₀N₄O₃F₃ [M + H]⁺ 409.1488; found, 409.1489; HPLC purity-97%; $t_R = 12.57$ min.

4.1.5.16 4-(Hydroxymethyl)-2-methyl-5-(((1-(3-(tri-

uoromethyl)benzyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl) *pyridin-3-ol (5p).* Yellow oil; yield 87%; $R_f = 0.4$ (DCM/methanol) 19 : 1); ¹H NMR (500 MHz, chloroform-*d*) δ 7.82 (s, 1H, N=C<u>H</u>) δ 7.62 (s, 1H, C=CH–N) δ 7.51 (d, J = 4.4 Hz, 1H, Ar–H) δ 7.49 (d, $J = 7.8$ Hz, 1H, Ar–H) δ 7.44 (s, 1H, Ar–H) δ 7.41 (d, $J = 7.7$ Hz, 1H, Ar-H) δ 5.54 (s, 2H, N-CH₂) δ 4.98 (s, 2H, CH₂-OH) δ 4.52 (s, 2H, CH₂–O) δ 4.46 (s, 2H, CH₂–O) δ 2.44 (s, 3H, N–C–CH₃)¹³C NMR (125 MHz, chloroform-d) δ 140.23, 140.11, 135.32, 131.44, 129.95, 125.88, 124.84, 122.68, 68.40, 59.69, 53.74, 29.78; ¹⁹F NMR (470 MHz, chloroform-*d*) δ -62.64 (s); IR (cm⁻¹) v: 3116.27, 2896.56, 2923.18, 2374.38, 2391.60, 1180.28, 1682.66, 1280.45, 1216.68; HRMS (m/z) : calculated for C₁₉H₂₀N₄O₃F₃ [M + H^{\dagger} 409.1488; found, 409.1492; HPLC purity-96%; $t_R = 8.70$ min.

4.1.5.17 4-(Hydroxymethyl)-2-methyl-5-(((1-(2-(tri uoromethyl)benzyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl) pyridin-3-ol (5q). Brown oil; yield 89%; $R_f = 0.4$ (DCM/methanol 19 : 1); ¹H NMR (500 MHz, chloroform-*d*) δ 7.82 (s, 1H, N=C<u>H</u>) δ 7.62 (s, 1H, C=CH–N) δ 7.51 (d, J = 4.4 Hz, 1H, Ar–H) δ 7.49 (d, $J = 7.8$ Hz, 1H, Ar-H $) \delta$ 7.44 (s, 1H, Ar-H $) \delta$ 7.40 (d, $J = 7.7$ Hz, 1H, Ar–H) δ 5.54 (s, 2H, N–CH₂) δ 4.98 (s, 2H, CH₂–OH) δ 4.52 (s, 2H, CH₂–O) δ 4.46 (s, 2H, CH₂–O) δ 2.44 (s, 3H, N–C–CH₃); ¹³C NMR (125 MHz, chloroform-d) d 152.85, 132.87, 130.47, 129.06, 126.41, 123.04, 68.43, 61.65, 59.65, 50.39, 19.04; 19F NMR (470 MHz, chloroform-d) δ –58.88 (s); IR (cm⁻¹) v: 3117.83, 2897.75, 2927.20, 2378.42, 2390.56, 1185.75, 1683.20, 1283.30, 1215.78; HRMS (m/z) : calculated for C₁₉H₂₀N₄O₃F₃ [M + H]⁺ 409.1488; found, 409.1490; HPLC purity-98%; $t_R = 12.09$ min. Paper Watch, 136.66, 139.41, 136.34, 132.4, 132.4, 6:43, 6:43, 14.5.16 4/3pdrograde)

2020. 332.4, 133.4, 142.4, 132.4, 132.4, 132.4, 2020. 2020. 2020. Article. Published United Uncertification-Noncommercial 3.0 Uncertifi

4.2 Biological activity

4.2.1 Inhibition experiment of AChE. For evaluation of AChE activity of the derivatives, we followed a modified Ellman's method using AChE from Electrophorus electricus²⁷. After adding 0.1 M phosphate buffer solution (pH-8, $120 \mu L$) to every well, AChE enzyme (0.5 U mL^{-1} , 10 μ L) along with different concentrations of test compounds (20 µL) were added and incubated at room temperature for 15 minutes. After incubation, reaction mixture comprising of 75 mM acetylthiocholine iodide and 10 mM 5,5'-dithiobis-2-nitrobenzoic acid (DTNB) were added to 96 well plate to generate yellow colouration due to 5-thio-2-nitro-benzoic acid. Absorbance was measured immediately after addition of the reaction mixture in a kinetic loop for 10 minutes at 412 nm at an interval of 1 minute using BioTek Synergy HTX Multimode Microplate reader. All the samples were assayed in triplicate and IC_{50} values were determined and expressed as mean \pm SEM as generated from Graph Pad Prism 8.0 software.

4.2.2 Antioxidant assay. The antioxidant capacity of all the compounds in our series were measured using oxygen radical absorbance capacity fluorescein (ORAC-FL) method previously described by Fang Lei et al.²⁸ with some modifications. The total reaction mixture was kept to 200 μ L and 75 mM phosphate buffer solution was used for every assay. In dark 96 well plate, 25 μ L of sample (8 μ M)/Trolox (8 μ M)/buffer was added along with 150 nM fluorescein solution (150 μ L) and incubated in dark at $37 °C$ for 30 minutes. Following this, free radical initiator,

RSC Advances **Paper** Paper **Paper Paper Paper**

 12 mM AAPH (25 µL) was added and immediately after addition, fluorescence readings were taken at excitation 485 nm and emission at 535 nm in a kinetic loop for 15 minutes at 1 minute interval. All the samples were assayed in triplicate in BioTek Synergy HTX Multimode Microplate reader. The area under the fluorescence decay curve (AUC) was calculated for every sample and total antioxidant capacity of the compound was determined by substractingit's AUC from the AUC of the blank. ORAC-FL values are expressed as Trolox equivalents and ORAC-FL value of Trolox is taken as 1. The ORAC-FL values of three individual experiments were expressed as expressed as mean \pm SEM as generated from Graph Pad Prism 8.0 software. **PSC** Advances Compare the control of the state of the computer control of the state on 1 control of the state on 1 control of the state of the

4.3 Metal chelation study

Metal chelation studies of the tested compounds were done using UV spectrophotometer (Agilent Technologies 500) within the wavelength range of 200 to 600 nm at 37 $^{\circ}$ C. Solutions of FeCl₃, CuCl₂, ZnSO₄·H₂O, AlCl₃·6H₂O, KCl, NaCl, CaSO₄, $Mg(OH)_2 \cdot 6H_2O$ and compounds to be tested were prepared, such that the resulting solution comprised 100 μ M each of ligand and metal and incubated at 37° C for 30 min. For taking the spectra of the compounds alone, metal solution was replaced with Milli Q water.

For affinity study of 5i towards Fe, all metals Fe^{3+} , Zn^{2+} , Cu^{2+} , Ca^{2+} , Mg^{2+} , Na⁺ and K⁺ were mixed, such that the reaction mixture contained equal concentration each including the ligand $5i$ as 100 μ M.

For stoichiometric studies, 100 μ M solution of 5i was taken in the cuvette and to it 20 μ L aliquots of 1 mM FeCl₃ was added till the final concentration reached of metal reached 100 μ M. The concentration of FeCl₃ ranged from 10-100 μ M and the titration was done in 37 °C.

4.4 In silico study

4.4.1 Molecular docking. The computational analyses of the compounds were done on Red Hat 5.0 Linux platform with Intel core 4 quad processor and 16 GB of RAM. The crystal structure of AChE (PDB ID 4EY5) was imported into Maestro software (Schrodinger Release 2019-3). The water molecules were removed and hydrogen bonds were added to the protein structure to make it adapt to an environment of pH 7, using protein preparation wizard, a function of maestro. Protein prep wizard utility was used to optimize the position of heavy atoms. The prepared protein was docked against the pyridoxine based triazoles by using the extra precession mode.

4.4.2 ADME prediction by SwissADME. The SwissADME software was accessed from http://www.swissadme.ch. The cdx. format of the structure was imported and the list of SMILES was entered and by clicking on the option RUN, the predictive properties were generated.

4.4.3 Toxicity prediction using ProTox-II. The online tool ProTox-II is a free open access web tool and accessed from http://www.tox.charite.de/tox. The input was provided with an integrated PubMed search of pyridoxine and then structural changes were done as per required for generating 5i. The results were automatically generated as shown in Section 2.6.3.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Tiyas Pal and Saipriyanka Bhimaneni are thankful to Department of Pharmaceutical, Ministry of Chemicals and Fertilizers for providing financial assistance. Spectral study was carried out at Central Instrumental Facility NIPER-R. NIPER-R/ Communication/123.

References

- 1 P. Christina, World Alzheimer's Report 2018, Alzheimer's Dis. Int. world Alzheimer Rep., 2018, pp. 1–48.
- 2 A. Cavalli, M. L. Bolognesi, A. Minarini, M. Rosini, V. Tumiatti, M. Recanatini and C. Melchiorre, Multi-Target-Directed Ligands to Combat Neurodegenerative Diseases, J. Med. Chem., 2008, 51(7), 2326.
- 3 R. R. Ramsay, M. R. Popovic-Nikolic, K. Nikolic, E. Uliassi and M. L. Bolognesi, A Perspective on Multi-Target Drug Discovery and Design for Complex Diseases, Clin. Transl. Med., 2018, 7(3), 1–14.
- 4 P. Mecocci and M. C. Polidori, Antioxidant Clinical Trials in Mild Cognitive Impairment and Alzheimer's Disease, Biochim. Biophys. Acta, Mol. Basis Dis., 2012, 1822(5), 631– 638.
- 5 M. A. Silva, A. S. Kiametis and W. Treptow, Donepezil Inhibits Acetylcholinesterase via Multiple Binding Modes at Room Temperature, J. Chem. Inf. Model., 2020, DOI: 10.1021/acs.jcim.9b01073.
- 6 H. Tachallait, A. Bouyahya, A. Talha, Y. Bakri, N. Dakka, L. Demange, R. Benhida and K. Bougrin, Concise Synthesis and Antibacterial Evaluation of Novel 3-(1,4-Disubstituted-1,2,3-Triazolyl)Uridine Nucleosides, Arch. Pharm., 2018, 351(11), 1–11.
- 7 A. Aziz Ali, D. Gogoi, A. K. Chaliha, A. K. Buragohain, P. Trivedi, P. J. Saikia, P. S. Gehlot, A. Kumar, V. Chaturvedi and D. Sarma, Synthesis and Biological Evaluation of Novel 1,2,3-Triazole Derivatives as Anti-Tubercular Agents, Bioorg. Med. Chem. Lett., 2017, 27(16), 3698–3703.
- 8 M. Irfan, B. Aneja, U. Yadava, S. I. Khan, N. Manzoor, C. G. Daniliuc and M. Abid, Synthesis, QSAR and Anticandidal Evaluation of 1,2,3-Triazoles Derived from Naturally Bioactive Scaffolds, Eur. J. Med. Chem., 2015, 93, 246–254.
- 9 A. Rastegari, H. Nadri, M. Mahdavi, A. Moradi, S. Sara, N. Edraki, F. Homayouni and B. Larijani, Bioorganic Chemistry Design, Synthesis and Anti-Alzheimer's Activity of Novel 1,2,3-Triazole-Chromenone Carboxamide Derivatives, Bioorg. Chem., 2019, 83, 391–401.
- 10 L. Jalili-Baleh, H. Nadri, H. Forootanfar, A. Samzadeh-Kermani, T. T. Küçükkılınç, B. Ayazgok, M. Rahimifard, M. Baeeri, M. Doostmohammadi, L. Firoozpour, S. N. A. Bukhari, M. Abdollahi, M. R. Ganjali, S. Emami,

M. Khoobi and A. Foroumadi, Novel 3-Phenylcoumarin– Lipoic Acid Conjugates as Multi-Functional Agents for Potential Treatment of Alzheimer's Disease, Bioorg. Chem., 2018, 79, 223–234.

- 11 X. Yang, X. Qiang, Y. Li, L. Luo, R. Xu, Y. Zheng, Z. Cao, Z. Tan and Y. Deng, Pyridoxine-Resveratrol Hybrids Mannich Base Derivatives as Novel Dual Inhibitors of AChE and MAO-B with Antioxidant and Metal-Chelating Properties for the Treatment of Alzheimer's Disease, Bioorg. Chem., 2017, 71, 305–314. Paper Water and A. Furnomell, Wavel A. Phenyleaumarin-

Metabologica Sarticle Commercial Agent Commercial Access 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020, 2020,
	- 12 M. Yazdani, N. Edraki, R. Badri, M. Khoshneviszadeh, A. Iraji and O. Firuzi, Multi-Target Inhibitors against Alzheimer Disease Derived from 3-Hydrazinyl 1,2,4-Triazine Scaffold Containing Pendant Phenoxy Methyl-1,2,3-Triazole: Design, Synthesis and Biological Evaluation, Bioorg. Chem., 2019, 84, 363–371.
	- 13 Z. Najafi, M. Mahdavi, M. Saeedi, E. Karimpour-Razkenari, R. Asatouri, F. Vafadarnejad, F. H. Moghadam, M. Khanavi, M. Sharifzadeh and T. Akbarzadeh, Novel Tacrine-1,2,3-Triazole Hybrids: In vitro, In vivo Biological Evaluation and Docking Study of Cholinesterase Inhibitors, Eur. J. Med. Chem., 2017, 125, 1200–1212.
	- 14 A. Kaur, S. S. Narang, A. Kaur, S. Mann, N. Priyadarshi, B. Goyal, N. K. Singhal and D. Goyal, Multifunctional Mono-Triazole Derivatives Inhibit AB_{42} Aggregation and Cu^{2+} -Mediated A β_{42} Aggregation and Protect against A β_{42} -Induced Cytotoxicity, Chem. Res. Toxicol., 2019, 32(9), 1824–1839.
	- 15 M. R. Jones, E. L. Service, J. R. Thompson, M. C. P. Wang, I. J. Kimsey, A. S. Detoma, A. Ramamoorthy, M. H. Lim and T. Storr, Dual-Function Triazole-Pyridine Derivatives as Inhibitors of Metal-Induced Amyloid-β Aggregation, Metallomics, 2012, 4(9), 910–920.
	- 16 J. M. Zhuo and D. Praticò, Acceleration of Brain Amyloidosis in an Alzheimer's Disease Mouse Model by a Folate, Vitamin B6 and B12-Deficient Diet, Exp. Gerontol., 2010, 45(3), 195-201.
	- 17 A. Hashim, L. Wanga, K. Junej, Y. Yeb, Y. Zhao and L. J. Ming, Vitamin B6s Inhibit Oxidative Stress Caused by Alzheimer's Disease-Related CuII-b-Amyloid Complexes-Cooperative Action of Phospho-Moiety, Bioorg. Med. Chem. Lett., 2011, 21(21), 6430–6432.
	- 18 L. Drtinova, P. Dobes and M. Pohanka, Low Molecular Weight Precursor Applicable for Alzheimer Disease Drugs Synthesis (AChE and BChE Inhibition, BACE Inhibition,

Antioxidant Properties and in Silico Modulation), J. Appl. Biomed., 2014, 12(4), 285–290.

- 19 A. Campbell, The Potential Role of Aluminium in Alzheimer's Disease, Nephrol., Dial., Transplant., 2002, 17, 17–20.
- 20 P. F. Good, D. P. Perl, L. M. Bierer and J. Schmeidler, Selective Accumulation of Aluminum and Iron in the Neurofibrillary Tangles of Alzheimer's Disease: A Laser Microprobe (LAMMA) Study, Ann. Neurol., 1992, 31(3), 286– 292.
- 21 H. Akrami, B. F. Mirjalili, M. Khoobi, H. Nadri, A. Moradi, A. Sakhteman, S. Emami, A. Foroumadi and A. Shafiee, Indolinone-Based Acetylcholinesterase Inhibitors: Synthesis, Biological Activity and Molecular Modeling, Eur. J. Med. Chem., 2014, 84, 375–381.
- 22 G. Johnson and S. Moore, The Peripheral Anionic Site of Acetylcholinesterase: Structure, Functions and Potential Role in Rational Drug Design, Curr. Pharm. Des., 2005, 12(2), 217–225.
- 23 A. Daina and V. Zoete, A BOILED-Egg to Predict Gastrointestinal Absorption and Brain Penetration of Small Molecules, ChemMedChem, 2016, 1117–1121.
- 24 A. Daina, O. Michielin and V. Zoete, SwissADME: A Free Web Tool to Evaluate Pharmacokinetics, Drug-Likeness and Medicinal Chemistry Friendliness of Small Molecules, Sci. Rep., 2017, 7, 1–13.
- 25 M. N. Drwal, P. Banerjee, M. Dunkel, M. R. Wettig and R. Preissner, ProTox: A Web Server for the in Silico Prediction of Rodent Oral Toxicity, Nucleic Acids Res., 2014, 42, 3–8.
- 26 N. L. Morozowich, A. L. Weikel, J. L. Nichol, C. Chen, L. S. Nair, C. T. Laurencin and H. R. Allcock, Polyphosphazenes Containing Vitamin Substituents: Synthesis, Characterization, and Hydrolytic Sensitivity, Macromolecules, 2011, 44(6), 1355–1364.
- 27 M. Pohanka, M. Hrabinova, K. Kuca and J. P. Simonato, Assessment of Acetylcholinesterase Activity Using Indoxylacetate and Comparison with the Standard Ellman's Method, Int. J. Mol. Sci., 2011, 12(4), 2631–2640.
- 28 K. Pérez-Cruz, M. Moncada-Basualto, J. Morales-Valenzuela, G. Barriga-González, P. Navarrete-Encina, L. Núñez-Vergara, J. A. Squella and C. Olea-Azar, Synthesis and Antioxidant Study of New Polyphenolic Hybrid-Coumarins, Arabian J. Chem., 2018, 11(4), 525–537.