RSC Advances



View Article Online

View Journal | View Issue

PAPER



Cite this: RSC Adv., 2020, 10, 29829

A simple method for the synthesis of *N*difluoromethylated pyridines and 4-pyridones/ quinolones by using BrCF₂COOEt as the difluoromethylation reagent⁺

Albert Gandioso,‡ Mohamed El Fakiri,‡ Anna Rovira‡ and Vicente Marchán 吵 *

We describe a novel transition metal-free method for the synthesis of *N*-difluoromethylated pyridines and 4-pyridones/quinolones by using readily available ethyl bromodifluoroacetate as a fluorine source. The formation of *N*-difluoromethylated pyridines involves a two-step process in which *N*-alkylation by ethyl bromodifluoroacetate is followed by *in situ* hydrolysis of the ester and decarboxylation. Besides optimizing the *N*-difluoromethylation conditions and assessing the influence of steric and electronic effects on the outcome of the reaction, we have synthesized the *N*-difluoromethylated analogues of two fluorophores and demonstrated that their spectroscopic properties can be improved through replacement of *N*-CH₃ group by *N*-CF₂H.

Received 20th July 2020 Accepted 7th August 2020

DOI: 10.1039/d0ra06322c

rsc.li/rsc-advances

Introduction

The introduction of fluorinated scaffolds in organic compounds is a well-established approach in medicinal, agricultural and biomaterial sciences for the modification of biologicallyrelevant properties such as binding affinity, metabolic stability, lipophilicity, and bioavailability.¹ Non-invasive diagnostic tools such as positron emission tomography (PET) and magnetic resonance imaging (MRI) are also based on compounds incorporating radioactive (¹⁸F) and non-radioactive (¹⁹F) fluorinated moieties, respectively.² Remarkably, about half of the most successful drugs are based on fluorine-containing compounds and more than three out of ten drugs approved by the FDA in the last two years contain this atom.³ In this context, there is a clear demand for novel methods to expand the current fluorination toolbox in organic synthesis.

Among fluorinated scaffolds, the difluoromethyl group (CF_2H) has attracted great interest owing to its unique physicochemical properties. It is considered a lipophilic bioisostere of hydroxyl and thiol functional groups capable of hydrogen bonding,⁴ which has been exploited to improve membrane permeability and binding affinity of biologically-active compounds.⁵ Despite these promising properties, the chemistry of difluoromethylation has been less extensively studied compared with that of trifluoromethylation. The late-stage

introduction of CF₂H group into organic substrates has been mainly investigated by metal-mediated photocatalytic and thermal processes,⁶ and by transition-metal-free strategies.^{6a,b,7}

Pyridine and its derivatives are found in natural products (*e.g.*, vitamins, coenzymes and alkaloids) and in a wide number of drugs and agrochemicals.⁸ More than 100 pyridinecontaining drugs have been approved by the FDA,⁹ making this nitrogen-containing heterocycle a privileged scaffold in medicinal chemistry. Remarkably, *N*-methylpyridinium moieties are found in several marketed drugs and in biologically active compounds such as antibacterials and anticancer agents and, consequently, replacement of the methyl group by CF_2H offers an unprecedented opportunity to discover novel drug candidates with improved (or simply different) biological activities. Similarly, *N*-difluoromethylation of pyridine might have a strong impact in the bioimaging field since many fluorescent probes incorporate *N*-alkylpyridinium moieties.¹⁰

The quinolin-4(1*H*)-one is also a common structural motif in many bioactive compounds, including well-known antibacterial agents such as fluoroquinolones.¹¹ Since many quinolone antibiotics are *N*-alkylated and incorporate at least one fluorine atom in their chemical structure, the *N*-difluoromethylation of the quinolin-4(1*H*)-one scaffold offers an exceptional opportunity to develop new antibiotics with an expanded spectrum and high efficacy.

Despite the enormous potential that the *N*-difluoromethylation reaction of pyridine and derivatives offers, only two methods have been reported to date (Scheme 1). In 1996, Röschenthaler and co-workers described the two-step *N*difluoromethylation reaction of DMAP by using CF_2Br_2 in the presence of activated copper, followed by hydrogenation with

Departament de Química Inorgànica i Orgànica, Secció de Química Orgànica, IBUB, Universitat de Barcelona, Martí i Franquès 1-11, E-08028, Barcelona, Spain. E-mail: vmarchan@ub.edu

[†] Electronic supplementary information (ESI) available: Experimental procedures and characterization data of all compounds. See DOI: 10.1039/d0ra06322c

[‡] These authors contributed equally to this work.

tributyltin hydride.12 However, the ozone depletion potential of CF₂Br₂ and manipulation risks advises against the use of this reagent. More recently, Zafrani, Gershonov and collaborators have described the difluorocarbene-mediated N-difluoromethylation of trialkyl amines using diethyl bromodifluoromethylphosphonate and a fluoride source.13 Although the reaction was optimized for the synthesis of N-difluoromethyltrialkylammonium salts, the method proved to be adequate for the N-difluoromethylation of DMAP. On the other hand, the two examples reported in the literature for the synthesis of Ndifluoromethylated 4-quinolones are based on the use of CHBrF₂ as fluorinating reagent of a quinolin-4-ol derivative,¹⁴ and on the photogeneration of difluorocarbene by deiodination and deacetylation of ethyl difluoroiodoacetate under basic conditions.15 Examples on the synthesis of N-difluoromethyl-2pyridone derivatives are based on the use of ClCF₂COONa/18crown-6 and BrCF₂COOEt/K₂CO₃.¹⁶ Herein, we describe a new straightforward transition-metal-free method for the synthesis of N-difluoromethylated pyridines based on the use of ethyl bromodifluoroacetate (1), which is a cheap, safe and commercially available reagent (Scheme 1). Based on this reaction, a simple procedure for the synthesis of N-difluoromethylated 4pyridones and 4-quinolones has been also discovered.

Results and discussion

Initially, DMAP was chosen as a model substrate to investigate the use of ethyl bromodifluoroacetate (1) as a potential *N*difluoromethylating reagent of pyridine. First, DMAP was reacted with 1 (5 mol equiv.) in ACN as a solvent (HPLC quality) at 60 °C for 24 h. Reversed-phase UV-HPLC-MS analysis (Fig. S1, S2 and Table S1†) analysis revealed the presence of compound **2a** (24%) resulting from the *N*-alkylation of the pyridine heterocycle together with a small amount of unreacted DMAP (4%) (Scheme 2). To our delight, both the carboxylic acid derivative **2b** (32%) and the *N*-difluoromethylated pyridinium salt **3** (40%) were also identified in the crude, which indicates that hydrolysis of the ester and decarboxylation take place *in situ* in the reaction media affording the expected *N*-difluoromethylated derivative **3**. Although halodifluoromethyl



Scheme 1 (A and B) Previously reported methods for the *N*-difluoromethylation of DMAP, and (C) synthesis of *N*-difluoromethylated pyridines and 4-pyridones described in this work.



Scheme 2 *N*-Difluoromethylation of DMAP using ethyl bromodifluoroacetate (1).

compounds have been extensively used as difluorocarbene synthons in difluoromethylation reactions,¹⁷ the presence of **2a** and **2b** supports a direct nucleophilic attack of the pyridine nitrogen on the electrophilic carbon of **1** rather than the generation of a difluorocarbene intermediate. In order to confirm this mechanism, compound **2b** was isolated by reversed-phase HPLC and fully characterized by ¹H, ¹³C and ¹⁹F-NMR, IR and HR ESI-MS (see ESI†). It is worth noting that the ester intermediate (**2a**) could not be isolated since it was completely transformed into **2b** during the purification by silica column chromatography.

Based on these results, we focused on the optimization of the reaction conditions for the *N*-difluoromethylation of DMAP, and the progress of the reaction was investigated by capillary electrophoresis analysis (Fig. S3 and S4†). As shown in Table 1 (entry 2), the amount of 3 was increased (83%) when THF (which was previously eluted through basic alumina) was used instead of ACN (56%). By contrast, the outcome of the reaction was negatively affected in distilled THF (Table 1, entry 3): the amount of *N*-alkylation did not significantly change but the transformation of the *N*-alkylated intermediates into 3 was considerably reduced. Very interestingly, the *N*-difluoromethylation of DMAP was quantitative (entry 5) when a 1 : 1 (v/ v) mixture of neutralized THF and ACN was used as a solvent. Again, the transformation of the intermediates into 3 was

Table 1 Optimization of the reaction conditions for the N-difluoromethylation of DMAP^a

Entry	Compound 1 equiv.	Solvent	Conversion ^d [%]
1	5	ACN	56 (95)
2	5	THF^{b}	83 (86)
3	5	THF^{c}	30 (93)
4	5	Dioxane	25 (56)
5	5	$ACN/THF^{b}(1:1)$	100 (100)
6	5	$ACN/THF^{c}(1:1)$	43 (100)
7	3	$ACN/THF^{b}(1:1)$	100 (100)
8	1.5	$ACN/THF^{b}(1:1)$	100 (100)
9	1.1	$ACN/THF^{b}(1:1)$	84 (84)
10^e	5	$ACN/THF^{b}(1:1)$	100 (100)

^{*a*} Reactions were performed on a 0.08 mmol scale in 2 mL of the solvent at 60 °C for 24 h. ^{*b*} THF was eluted through basic alumina. ^{*c*} THF was distilled over sodium and benzophenone. ^{*d*} Conversion yield was determined by capillary electrophoresis analysis. The percentage value corresponds to the amount of **3** in the reaction mixture and the value in parenthesis to the sum of the amount of **3** and **2a** + **2b**. ^{*c*} Reaction was carried out under microwave irradiation during 2 h at 60 °C.



Expansions of the ¹H (left) and ¹³C NMR (right) spectra of **3**



Fig. 2 N-Difluoromethylation of pyridine derivatives (4-7) and pyridine-containing fluorophores (8-9). ^a Conversion yield was determined by ¹H NMR (4a–6a) or HPLC-MS (7a, 8b–9b) analysis. ^b Isolated yield.

reduced when distilled THF was combined with ACN (entry 6), suggesting that the presence of basic traces in aluminaneutralized THF facilitates the in situ hydrolysis of the ester and decarboxylation (see below). With the optimized conditions in hand, the reaction was repeated at large scale and the pure Ndifluoromethylated compound was isolated (yield: 92%) and fully characterized by ¹H, ¹³C and ¹⁹F-NMR and HR ESI-MS. Fig. 1 shows the aromatic region of the ¹H-NMR of 3 where a characteristic triplet at 8.54 ppm resulting from the coupling of the fluorine atoms to the proton of the CF₂H group is observed. Similarly, the carbon of the CF₂H group appears as a triplet at 111 ppm in the ¹³C NMR spectrum (Fig. 1), and the fluorine atoms as a doublet in the ¹⁹F NMR spectrum (see ESI†).

Further evidence of the two-step process for the N-difluoromethylation of pyridine and of the source of proton in the final N-difluoromethylated product (3) was obtained by investigating the reaction of DMAP with 1 by NMR in a 98:2 (v/v) mixture of CD₃CN and D₂O. Very interestingly, a deuterated analogue of 3 (compound 3-d) was obtained as a major product (97%). Indeed, the CF₂D group of 3-d appears in the ¹⁹F NMR spectrum (Fig. S5[†]) as a triplet, which results from the coupling of the fluorine atoms to deuterium. The presence of a doublet (3% relative to the triplet) in the ¹⁹F NMR spectrum and of a small triplet in the ¹H NMR spectrum (Fig. S6[†]) confirmed the formation of compound 3, which can be attributable the residual non-deuterated water present both in D₂O and in CD₃CN. As expected, HR ESI-MS analysis (Fig. S7[†]) showed that the mass of 3-d was one unit higher than that of 3 (Table S2[†]) These results confirmed that hydrolysis of the ester intermediate 2a to the carboxylic acid 2b can be mediated by traces of water present in the solvent, which would also be source of the proton of the CF₂H group in the final product. The fact that around 20% of compound 2b was spontaneously transformed into the N-difluoromethylated pyridinium salt (3) after standing in CD₃CN solution at room temperature (see ¹H NMR in Fig. S8[†]), also confirms the ease with which decarboxylation takes place from the carboxylic acid derivative. This is in contrast with the slow decarboxylation rate described for 2-aryl-2,2-difluoroacetic acid derivatives which require the addition of additives and high temperature heating.17,18

Once established the best combination of solvents, we investigated the effect of reducing the amount of 1 on the reaction efficiency. As shown in Table 1, the reaction between



Fig. 3 Comparison of the normalized absorption (left) and fluorescence (right) spectra of N-CH₃ fluorophores (8a-9a) and their N-CF₂H analogues (8b-9b) in ACN



Scheme 3 Synthesis of *N*-difluoromethylated 4-pyridone and 4quinolone (A) and proposed pathway for the formation of compound **10b** and **10c** in the reaction of 4-methoxypyridine with **1** (B).

DMAP and **1** still gave good conversion yields at almost equimolar ratios between both reactants (entry 9), being quantitative with 1.5 mol equiv. of **1** after 24 h at 60 °C (entry 8). In all cases, a single product (**3**) was always detected in the reaction crude by capillary electrophoresis (Fig. S4†). Importantly, the reaction time can be reduced (up to 2 h) by using microwave irradiation (entry 10).

To gain insight into the scope and limitations of this novel synthetic method, the N-difluoromethylation of different pyridine-containing substrates was examined. First, the influence of the steric hindrance around the heterocyclic nitrogen of pyridine was investigated by evaluating the N-difluoromethylation of pyridine (4), 2-picoline (5) and 2,6-lutidine (6) (Fig. 2). All the reactions were carried out under the DMAPoptimized conditions (Table 1, entry 5), and the conversion yield was determined by ¹H NMR (Fig. S9-S11[†]) by comparing the integration of diagnostic peaks in the starting compounds (4-6) and in the N-difluoromethylated analogues (4a-6a). The conversion yield of pyridine and 2-methylpyridine into their Ndifluoromethylated derivatives was found 63% and 81%, respectively. On the one hand, the reduced reactivity of pyridine towards 1 compared with DMAP seems to be a consequence of the absence of an electron-donating group that activates the electron-deficient pyridine heterocycle. On the other hand, the electron-donating effect of the methyl group in 2-picoline seems to compensate the increased steric hindrance around the nitrogen atom, leading to a higher conversion into the Ndifluoromethylated compound compared with pyridine. In the case of 2,6-lutidine, NMR spectra of the reaction crude showed the lowest conversion yield among the three pyridine derivatives investigated (<5%), which confirms that in this case steric effects prevail over the electron-donating effect of the two methyl substituents. In good agreement with these results, the conversion yield for the N-difluoromethylation of ethyl isonicotinate was found very low (<5%), both under the DMAPoptimized conditions and under microwave irradiation, which

indicates that the ethyl bromodifluoroacetate-based method does not tolerate the presence of electron-withdrawing groups in the pyridine.

N-Alkylation of heterocycles has been widely exploited to modulate the photophysical and physicochemical properties of organic fluorophores,¹⁰ and many fluorescent mitochondrial probes incorporate lipophilic positively charged moieties such as N-methyl pyridinium.¹⁹ In this context, we envisaged that replacement of the N-methyl group in pyridine-containing fluorophores by N-CF₂H could have a strong impact on the spectroscopic properties of the compounds. Our group has recently described a new class of push-pull fluorophores, nicknamed COUPYs, based on the N-alkylation of a pyridine moiety in a novel coumarin scaffold (e.g., compound 8 in Fig. 2) which exhibit several attractive photophysical properties, including emission in the far-red/NIR region.20 Taking into account that absorption and emission maxima can be red-shifted through the incorporation of strong electron-withdrawing groups via Nalkylation of the pyridine heterocycle,^{20a,b} here we focused on investigating the N-difluoromethylation of the parent COUPY scaffold (8) to further explore the scope of the methodology. To our delight, 8 was efficiently N-difluoromethylated by using 1 and the expected COUPY derivative (8b) was isolated by silica column chromatography as a dark blue solid (yield: 83%) and fully characterized by NMR and HRMS. Similarly, N-difluoromethylation of acridine orange (9) with 1 provided a novel Ndifluoromethylated analogue (9b) of this commercially available dye. Having at hand compounds 8b and 9b, we investigated the effect of replacing the N-CH3 group in the parent compounds (8a and 9a, respectively; see Fig. 2) by N-CF₂H on their spectroscopic properties (Fig. 3). Remarkably, the absorption maximum of **8b** ($\lambda_{abs} = 580$ nm) was considerably red-shifted (ca. 32 nm) with respect to the parent compound 8a ($\lambda_{abs} =$ 548 nm), and the same tendency was found with the acridine derivatives ($\lambda_{abs} = 497$ nm for **9a** and $\lambda_{abs} = 532$ nm for **9b**). The emission maxima of the compounds were also red-shifted with respect to the parent compounds ($\lambda_{em} = 638 \text{ nm for } \mathbf{8b} \nu s. \lambda_{em} =$ 603 nm for 8a, and $\lambda_{em} = 565$ nm for 9b vs. $\lambda_{em} = 524$ nm for 9a), which reproduces the effect of replacing the N-CH₃ group with *N*-CF₂H on the compounds' absorption maxima.

Finally, it is worth noting that during the evaluation of the substrate scope of the N-difluoromethylation method with pyridine derivatives containing electron-donating groups, an unexpected result was revealed. HPLC-MS analysis for the reaction of 4-methoxypyridine (10) with 1 showed the disappearance of 10 but neither the major product (83%) nor the minor product (17%) were the expected N-difluoromethylated salt (10a) (Scheme 3). MS data of the major compound was consistent with the formation of the N-difluoromethylated pyridin-4-one (10b), and the minor product was characterized as the N-methylated derivative of 4-methoxypyridine (10c). As a possible mechanism, we postulate the nucleophilic attack of the bromide anion on the methoxy substituent of 10a, which would generate 10b and CH₃Br. The formation of 10c can be explained by N-methylation of unreacted 4-methoxypyridine with methyl bromide (Scheme 3). A similar mechanism has been proposed by Ma and collaborators for the conversion of 4-

Paper

methoxypyridines into *N*-methyl-4-pyridones.²¹ Compound **10b** was isolated by silica column chromatography (55% yield) and fully characterized by NMR spectroscopy and HR ESI-MS. ¹H and ¹⁹F-NMR spectra showed the expected multiplicities for the proton and fluorine atoms of the CF₂H group (triplet and doublet, respectively), and ¹³C-NMR spectrum provided further evidence of the structure of the pyridin-4-one moiety: in addition to the expected triplet for the CF₂H group, a characteristic peak due to the carbonyl function was observed around 180 ppm. Similarly, reaction of 4-methoxyquinoline (**11**) with ethyl bromodifluoroacetate provided the corresponding *N*-difluoromethylated 4-quinolin-4-one (**11a**) with a 60% yield after purification (Scheme 3).

Conclusions

In summary, a novel transition metal-free synthetic method for the N-difluoromethylation of pyridine-containing substrates has been developed. The procedure involves the use of a cheap, safe and readily available reagent, ethyl bromodifluoroacetate (1), which provides the corresponding N-difluoromethylated pyridinium salts in moderate to good yields after reaction with the parent pyridine compounds. The formation of N-difluoromethylated pyridines involves a two-step process in which Nalkylation by ethyl bromodifluoroacetate is followed by in situ ester hydrolysis and decarboxylation By investigating the substrate scope and limitations of the method with a series of pyridine-containing molecules, we have observed that steric and electronic effects of the substituents in the heterocycle play an important role both on the effectiveness of the reaction and on the structure of the final compound. Remarkably, the presence of a methoxy group at the para position relative to the nitrogen atom of the pyridine ring results in the formation of a N-difluoromethylated pyridin-4-one structure, while the presence of electron-withdrawing groups difficult the reaction. This new facile N-difluoromethylating method should allow the synthesis of novel biologically active compounds in which the N-methyl group in N-methyl pyridinium and N-methyl-4pyridone/quinolone moieties could be easily replaced by N-CF₂H. In addition, we anticipate that N-difluoromethylation of conventional pyridine-containing organic fluorophores will allow to improve their spectroscopic properties, opening the door to novel fluorescent probes for bioimaging applications. Work is in progress in our laboratory to apply this new method to the synthesis of other N-difluoromethylated heterocycles, including imidazole, indole and pyrrol.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by funds from the Spanish Government (MCIU/AEI/FEDER, UE; grant CTQ2017-84779-R) and the Generalitat de Catalunya (2017 DI 072). The authors acknowledge helpful assistance of Josep Martí (capillary electrophoresis), Dr Francisco Cárdenas (NMR) and Dr Irene Fernández and Laura Ortiz (MS) from CCiTUB. A. R. and A. G. were recipient fellows of the University of Barcelona.

Notes and references

- 1 (*a*) T. Liang, C. N. Neumann and T. Ritter, Introduction of Fluorine and Fluorine-Containing Functional Groups, *Angew. Chem., Int. Ed.*, 2013, **32**, 8214; (*b*) N. A. Meanwell, Fluorine and Fluorinated Motifs in the Design and Application of Bioisosteres for Drug Design, *J. Med. Chem.*, 2018, **61**, 5822.
- 2 (a) A. F. Brooks, J. J. Topczewski, N. Ichiishi, M. S. Sanford and P. H. J. Scott, Late-stage [¹⁸F] fluorination: new solutions to old problems, *Chem. Sci.*, 2014, 5, 4545; (b) I. Tirotta, V. Dichiarante, C. Pigliacelli, G. Cavallo, G. Terraneo, F. B. Bombelli, P. Metrangolo and G. Resnati, ¹⁹F Magnetic Resonance Imaging (MRI): From Design of Materials to Clinical Applications, *Chem. Rev.*, 2015, 115, 1106.
- 3 (a) Y. Zhou, J. Wang, Z. Gu, S. Wang, W. Zhu, J. L. Aceña, V. A. Soloshonok, K. Izawa and H. Liu, Next Generation of Fluorine-Containing Pharmaceuticals, Compounds Currently in Phase II-III Clinical Trials of Major Pharmaceutical Companies: New Structural Trends and Therapeutic Areas, Chem. Rev., 2016, 116, 422; (b) H. Mei, J. Han, S. Fustero, M. Medio-Simon, D. M. Sedgwick, C. Santi, R. Ruzziconi and V. A. Soloshonok, Fluorine-Containing Drugs Approved by the FDA in 2018, Chem.-Eur. J., 2019, 25, 11797; (c) B. G. de la Torre and F. Albericio, The Pharmaceutical Industry in 2019. An Analysis of FDA Drug Approvals from the Perspective of Molecules, Molecules, 2020, 25, 745.
- 4 C. D. Sessler, M. Rahm, S. Becker, J. M. Goldberg, F. Wang and S. J. Lippard, CF₂H, a Hydrogen Bond Donor, *J. Am. Chem. Soc.*, 2017, **139**, 9325.
- 5 (*a*) N. A. Meanwell, Synopsis of Some Recent Tactical Application of Bioisosteres in Drug Design, *J. Med. Chem.*, 2011, 54, 2529; (*b*) Y. Zafrani, D. Yeffet, G. Sod-Moriah, A. Berliner, D. Amir, D. Marciano, E. Gershonov and S. Saphier, Difluoromethyl Bioisostere: Examining the "Lipophilic Hydrogen Bond Donor" Concept, *J. Med. Chem.*, 2017, 60, 797.
- 6 (a) Y. Lu, C. Liu and Q.-T. Chen, Recent Advances in Difluoromethylation Reaction, Curr. Org. Chem., 2015, 19, 1638; (b) D. E. Yerien, S. Barata-Vallejo and A. Postigo, Difluoromethylation Reactions of Organic Compounds, Chem.-Eur. J., 2017, 23, 14676; (c) Z. Feng, Q.-Q. Min, X.-P. Fu, L. An and X. Zhang, . Chlorodifluoromethane-triggered formation of difluoromethylated arenes catalysed by palladium, Nat. Chem., 2017, 9, 918; (d) Z. Feng, Y.-L. Xiao and X. Zhang, Transition-Metal (Cu, Pd, Ni)-Catalyzed Difluoroalkylation via Cross-Coupling with Difluoroalkyl Halides, Acc. Chem. Res., 2018, 51, 2264; (e) X.-P. Fu, X.-S. Xue, X.-Y. Zhang, Y.-L. Xiao, S. Zhang, Y.-L. Guo, X. Leng, K. N. Houk and X. Zhang, Controllable

catalytic difluorocarbene transfer enables access to diversified fluoroalkylated arenes, *Nat. Chem.*, 2019, **11**, 948.

- 7 (a) P. Dai, X. Yu, P. Teng, W.-H. Zhang and C. Deng, Visible-Light- and Oxygen-Promoted Direct Csp²-H Radical Difluoromethylation of Coumarins and Antifungal Activities, *Org. Lett.*, 2018, 20, 6901; (b) A. Lemos, C. Lemaire and A. Luxen, Progress in Difluoroalkylation of Organic Substrates by Visible Light Photoredox Catalysis, *Adv. Synth. Catal.*, 2019, 361, 1500; (c) Y. Huang, Z. Lin, Y. Chen, S. Fang, H. Jiang and W. Wu, Transition-metal-free N-difluoromethylation of hydrazones with TMSCF₂Br as the difluoromethylation reagent, *Org. Chem. Front.*, 2019, 6, 2462; (d) X. Xu and F. Liu, Transition-metal-free radical tri-/difluoromethylation of *N*,*N*-dialkylhydrazones with sodium sulfinates, *Org. Chem. Front.*, 2017, 4, 2306.
- 8 Y. Higashio and T. Shoji, Heterocyclic compounds such as pyrrole, pyridines, pyrrolidine, piperidine, indole, imidazol and pyrazines, *Appl. Catal., A*, 2004, **260**, 251.
- 9 E. Vitaku, D. T. Smith and J. T. Njardarson, Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceutical, *J. Med. Chem.*, 2014, 57, 10257.
- 10 (a) Q. Zheng and L. D. Lavis, Development of photostable fluorophores for molecular imaging, *Curr. Opin. Chem. Biol.*, 2017, **39**, 32; (b) L. G. Freidus, P. Pradeep, P. Kumar, Y. E. Choonara and V. Pillay, Alternative fluorophores designed for advanced molecular imaging, *Drug Discovery Today*, 2018, **23**, 115.
- 11 (a) T. D. M. Pham, Z. M. Ziora and M. A. T. Blaskovich, Quinolone antibiotics, *MedChemComm*, 2019, 10, 1719; (b)
 P. Ghosh and S. Das, Synthesis and Functionalization of 4-Quinolones – A Progressing Story, *Eur. J. Org. Chem.*, 2019, 4466.
- 12 A. Kolomeitsev, R.-M. Schoth, E. Lork and G.-V. Röschenthaler, Facile new method for synthezising N-polyfluoroalkylated heterocycles – molecular structure of N-(bromodifluoromethyl)-4-dimethylaminopyridinium bromide, *Chem. Commun.*, 1996, 335.
- 13 Y. Zafrani, D. Amir, L. Yehezkel, M. Madmon, S. Saphier, N. Karton-Lifshin and E. Gershonov, Chemoselective N-Difluoromethylation of Functionalized Tertiary Amines, *J. Org. Chem.*, 2016, **81**, 9180.
- 14 J. Tani, Y. Mushika and T. Yamaguchi, Studies on biologically active halogenated compounds. IV. Synthesis and antibacterial activity of fluorinated quinoline derivatives, *Chem. Pharm. Bull.*, 1982, **30**, 3530.
- 15 Y. Li, M. Rao, Z. Fan, B. Nian, Y. Yuan and J. A. Cheng, A visible-light-irradiated electron donor-acceptor complex-promoted radical reaction system for the CH perfluoroalkylation of quinolin-4-ols, *Tetrahedron Lett.*, 2019, **60**, 151046.

- 16 (a) J. Hu, W. Zhang and F. Wang, Selective difluoromethylation and monofluoromethylation reactions, *Chem. Commun.*, 2009, 7465; (b) M. Ando, T. Wada and N. Sato, Facile One-Pot Synthesis of N-Difluoromethyl-2-pyridone Derivatives, *Org. Lett.*, 2006, 8(17), 3805.
- 17 (a) X. Ma, S. Mai, Y. Zhou, G.-J. Cheng and Q. Song, Dual role of ethyl bromodifluoroacetate in the formation of fluorine-containing heteroaromatic compounds, *Chem. Commun.*, 2018, 54, 8960; (b) Z. Feng, Q.-Q. Min and X. Zhang, Access to Difluoromethylated Arenes by Pd-Catalyzed Reaction of Arylboronic Acids with Bromodifluoroacetate, *Org. Lett.*, 2019, 18, 44.
- 18 K. Fujikawa, Y. Fujioka, A. Kobayashi and H. Amii, A New Method for Aromatic Difluoromethylation: Copper-Catalyzed Cross-Coupling and Decarboxylation Sequence from Aryl Iodides, *Org. Lett.*, 2011, 13, 5560.
- 19 W. Xu, Z. Zeng, J.-H. Jiang, Y.-T. Chang and L. Yuan, Discerning the Chemistry in Individual Organelles with Small-Molecule Fluorescent Probes, *Angew. Chem., Int. Ed.*, 2016, **55**, 13658.
- 20 (a) A. Gandioso, R. Bresolí-Obach, A. Nin-Hill, M. Bosch, M. Palau, A. Galindo, S. Contreras, A. Rovira, C. Rovira, S. Nonell and V. Marchán, Redesigning the Coumarin Scaffold into Small Bright Fluorophores with Far-Red to Near-Infrared Emission and Large Stokes Shifts Useful for Cell Imaging, J. Org. Chem., 2018, 83, 1185; (b) A. Gandioso, M. Palau, R. Bresolí-Obach, A. Galindo, A. Rovira, M. Bosch, S. Nonell and V. Marchán, High Photostability in Nonconventional Coumarins with Far-Red/NIR Emission through Azetidinyl Substitution, J. Org. Chem., 2018, 83, 11519; (c) A. Rovira, A. Gandioso, M. Goñalons, A. Galindo, A. Massaguer, M. Bosch and Marchán, Solid-Phase Approaches for Labeling V. Targeting Peptides with Far-Red Emitting Coumarin Fluorophores, J. Org. Chem., 2019, 84, 1808; (d) V. Novohradsky, A. Rovira, C. Hally, A. Galindo, G. Vigueras, A. Gandioso, M. Svitelova, R. Bresolí-Obach, H. Kostrhunova, L. Markova, J. Kasparkova, S. Nonell, J. Ruiz, V. Brabec and V. Marchán, Towards Novel Photodynamic Anticancer Agents Generating Superoxide Anion Radicals: A Cyclometalated Ir^{III} Complex Conjugated to a Far-Red Emitting Coumarin, Angew. Chem., Int. Ed., 2019, 58, 6311; (e) A. Rovira, M. Pujals, A. Gandioso, M. López-Corrales, M. Bosch and V. Marchán, Modulating Photostability and Mitochondria Selectivity in Far-Red/NIR Emitting Coumarin Fluorophores through Replacement of Pyridinium by Pyrimidinium, J. Org. Chem., 2020, 85, 6086.
- 21 X. Yi, J. Chen, X. Xu and Y. Ma, Solvent and substituent effects on the conversion of 4-methoxypyridines to N-methyl-4-pyridones, *Synth. Commun.*, 2017, **47**, 872.