


 Cite this: *RSC Adv.*, 2020, 10, 40582

 Received 8th October 2020
 Accepted 23rd October 2020

DOI: 10.1039/d0ra08580d

rsc.li/rsc-advances

A scalable and green one-minute synthesis of substituted phenols†

 Vijayaragavan Elumalai  and Jørn H. Hansen *

A mild, green and highly efficient protocol was developed for the synthesis of substituted phenols *via ipso*-hydroxylation of arylboronic acids in ethanol. The method utilizes the combination of aqueous hydrogen peroxide as the oxidant and H₂O₂/HBr as the reagent under unprecedentedly simple and convenient conditions. A wide range of arylboronic acids were smoothly transformed into substituted phenols in very good to excellent yields without chromatographic purification. The reaction is scalable up to at least 5 grams at room temperature with one-minute reaction time and can be combined in a one-pot sequence with bromination and Pd-catalyzed cross-coupling to generate more diverse, highly substituted phenols.

Introduction

Phenols are indispensable in synthesis and constitute a privileged structural scaffold present in pharmaceuticals, natural products and synthetic polymers, among many other applications.¹ Consequently, considerable attention has been focused on the synthesis of phenol and its derivatives.² Traditional methods include nucleophilic aromatic substitution of aryl halides,³ transition-metal catalyzed processes,⁴ hydroxylation of benzene,⁵ oxidation of cumene⁶ and hydrolysis of diazo compounds.⁷ These methods have some major limitations, including harsh reaction conditions, reagent toxicities and the need for additives and complex ligands. The *ipso*-hydroxylation of boronic acids is an alternative route for the synthesis of phenols which avoids many of the problems with previous approaches. The boronic acids have low toxicity, higher stability in air, are easy to handle and display high functional group diversity. Aryl and heteroarylboronic acids are readily available and have been employed in several studies on *ipso*-hydroxylation chemistry. These mainly involved transition metal catalysis⁸ (Scheme 1A-i), photocatalysis⁹ (Scheme 1A-ii) or stoichiometric oxidants such as H₂O₂,¹⁰ oxone,¹¹ *N*-oxides,¹² organic hypervalent iodine(III),¹³ benzoquinone,¹⁴ mCPBA,¹⁵ NaBO₃ (ref. 16) and molecular oxygen¹⁷ (Scheme 1A-iii). Moreover, numerous variants have been reported recently for the synthesis of substituted phenols from boronic acids.¹⁸ Despite the high efficiency of many of these protocols, they are often characterized by high temperatures and long reaction times, the use of transition metals, necessity of strong base addition and

the use of harmful solvents. Furthermore, in many of these methods, scaled-up synthesis of phenols has not been demonstrated. When summarizing published papers on the use of hydrogen peroxide,^{18g} as one of the simplest and most available green oxidants, surprisingly, we find that virtually all of them report using extra additives of some kind and relatively long reaction times.

Recognizing the power of the transformation in the synthesis of diverse phenols, we set out to find mild, practical and simple reaction conditions with a minimum of additives, high chemical yields and with a clear green profile. Herein we report a very rapid, catalyst/additive-free and practical protocol for the synthesis of phenols *via* the *ipso*-hydroxylation of arylboronic acids using hydrogen peroxide in ethanol. The protocol represents a significant advance in the field as it works with unprecedented efficiency under ambient, green conditions and affords substantially improved chemical yields of phenols with only one minute reaction time and without flash column chromatography. Furthermore, adding hydrogen bromide yields bromophenols directly in a completely novel one-pot system which advances a truly practical and scalable approach to access highly substituted phenols (Scheme 1B).


Results and discussion

Previous reports on the use of hydrogen peroxide as oxidant for the *ipso*-hydroxylation were found to be of high interest because of the simplicity and clear green profile of H₂O₂.^{10,18g} Although several solvent systems have been reported with hydrogen peroxide, we decided to screen a range of solvents to ascertain reaction performance at ambient temperature and atmosphere with only one-minute reaction time. These reaction conditions underline our search for a highly efficient and truly practical transformation. In the solvent survey, conducted with the

UiT The Arctic University of Norway, Department of Chemistry, Chemical Synthesis and Analysis Group, N9037 Tromsø, Norway. E-mail: jorn.h.hansen@uit.no

† Electronic supplementary information (ESI) available: Procedures, analytical data and spectra is available online. See DOI: 10.1039/d0ra08580d






[a] Reaction Procedure: To a stirred solution of phenylboronic acid **1** (1 mmol) in EtOH (3 mL), was added H₂O₂ (30%, 3 equiv.) and stirred for 1 minute at room temperature.
 [b] Compound **2e** was purified by silica-gel flash column chromatography.

Scheme 2 Scope of *ipso*-hydroxylation chemistry.

one-minute protocol. Even *ortho*-substituted boronic acids work well in high to excellent chemical yields (83–96%). Notably, **2i** and **2p** were formed in excellent 95% and 96% yields, respectively, thus demonstrating very high tolerance to steric hindrance around the boronic acid moiety. Likewise, *meta*-substituted substrates were well tolerated across electronic diversity (80–92%), except of the *meta*-nitro substituted boronic acid **1e**, which afforded 60% yield of **2e**. This was the only entry that required flash column chromatography to obtain pure material. Furthermore, the chemistry is compatible with heterocycles as demonstrated with thiophene-2-boronic acid **1u** and indole-5-boronic acid **1v**. The former was isolated as the keto-form **2u**, which is a more stable tautomer of the corresponding aromatic hydroxy compound, in 80% yield. The 5-hydroxyindole product **2v** was isolated in 91% yield. To our delight, the polyaromatic pyrene boronic acid **1w** underwent a highly efficient transformation to **2w** in 98% yield. Thus, in addition to very short reaction time and practical conditions, the chemistry appears to be relatively independent of the electronic and steric nature of the substrate. The transformation under our conditions clearly has a broad scope and potential great synthetic utility in the formation of aromatic hydroxy compounds.




[a] Reaction procedure: To the stirred solution of phenylboronic acid **1** (5 g, 41 mmol) in EtOH (50 mL), was added H₂O₂ (30%, 12.5 mL, 3 equiv.) and stirred for 1 minute at ambient temperature. Isolated yield were reported.

Scheme 3 Gram-scale synthesis of phenols.

In order to further demonstrate the applicability of our reaction conditions, three of the *ipso*-hydroxylations were conducted at 5 gram-scale by simple scale-up of *p*-substituted systems. 85–90% isolated yields were obtained with electron-donating (methoxy), electron-withdrawing (cyano) and electron-neutral (H) examples shown in Scheme 3 (**2a**, **2b** and **2g**). This suggests that our conditions should be the method of choice for such a transformation conducted at larger scales for synthetic purposes.


The mechanism for the *ipso*-hydroxylation of boronic acids is presumed to be as outlined in Scheme 4.¹⁹ The boronic acid can form an adduct with hydrogen peroxide which can undergo rapid proton transfer (particularly in protic solvents) to afford intermediate **1a**-H₂O₂-2, set up for 1,2-aryl migration with departure of water, which gives the boronic ester readily amenable to hydrolysis to afford the phenol.

Treatment of phenols with HBr under oxidative conditions is known to yield bromophenols *via* electrophilic aromatic substitution.^{10f} To our delight, the addition of hydrogen bromide to the *ipso*-hydroxylation reaction from the very start yielded brominated phenols **3a–k** directly (Scheme 5), even though the reaction time was still kept at one minute. This direct one-pot protocol for generation of bromophenols from arylboronic acids is novel. The reaction likely proceeds by tandem *ipso*-hydroxylation-bromination, both processes which are facilitated by hydrogen peroxide. All the reactions involved must be rapid, since only a minute total reaction time afforded medium to excellent yields of the bromophenols (53–95% yield). The simple phenylboronic acid **1a** afforded the tribrominated phenol **3a** in excellent 95% yield. The electron-withdrawing *p*-cyano system **1b** gave only slightly diminished yield (89%) of **3b**.



Scheme 4 Hypothesized reaction mechanism for the *ipso*-hydroxylation of boronic acids.






[a] **Reaction Procedure:** To a stirred solution of phenylboronic acid **1** (1 mmol) in EtOH (3 mL), was added H_2O_2 (30%, 3 equiv.) followed by HBr (62%, 3 equiv.) and stirred for 1 minute at room temperature.

[b] *Ips*o brominated product **3k'** (38%) was obtained under the reaction conditions.

Scheme 5 Tandem *ipso*-hydroxylation-bromination.

The remaining *p*-substituted systems afforded medium to good yields of *o*-dibrominated products in 52–84% yields. *Ortho*- and *meta*-substituted systems afforded *ortho,para*-dibrominated products in 55–88% yields. The *meta*-brominated substrate **1j** afforded the tetrabromophenol **3j** in 52% yield. The *para*-acetamidoboronic acid **1k** afforded the *ortho*-dibrominated phenol **3k** in 54% yield, which represents a formal synthesis of bromo-functionalized pharmaceutical agent paracetamol. In this case, another product **3k'** was also isolated which turned out to be *para*-bromoacetamidobenzene in 38% yield. This represents a formal *ipso*-bromination of the boronic acid.²⁰ The tandem *ipso*-hydroxylation – bromination procedure appears to work quite well with a range of substrate and affords medium to excellent yields of bromophenols. This represents an attractive and direct route to this class of compounds due to its high efficiency and convenient procedure.

Our development of the one minute route to bromophenols, triggered us to investigate whether another reaction could be combined in the same pot in order to access further functionalized phenols. The presence of bromo-substituents would allow access to arylated phenols *via* cross-coupling chemistry. By generating bromophenols as described above, followed by



[a] **Reaction Procedure:** To the stirred solution of phenylboronic acid **1** (1 mmol) in EtOH (5 mL), H_2O_2 (30%, 3 mmol) and HBr (3 mmol) was added and stirred for 1-min. at ambient temperature, followed by adding the boronic acid (5 eq), Na_2CO_3 (5 eq) and $\text{Pd(PPh}_3)_4$ (6 mol%), flushed with nitrogen and irradiated with microwaves at 100 °C for 30 minutes. Isolated yields are reported.

[b] Na_2CO_3 (3 eq.) boronic acid (3 eq) and $\text{Pd(PPh}_3)_4$ (4 mol%) was used

Scheme 6 One-pot *ipso*-hydroxylation-bromination and Suzuki coupling to generate aryl-substituted phenols.



addition of base, 5 equivalents of a new boronic acid, tetrakis-palladiumtriphenylphosphine (6 mol%) and degassing with nitrogen, the Suzuki–Miyaura coupling products **4a–c** were generated in 20–67% yields (over 3 steps) upon microwave heating for 30 min (Scheme 6). **4a** was formed in diminished 20% yield using 3 equivalents of boronic acid and base with 4 mol% of Pd-catalyst loading. However, by increasing to 5 equivalents of boronic acid and base with 6 mol% of catalyst loading (2 mol% per coupling), the yields increased to 65% and 67% for **4b** and **4c**, respectively. Over 3 steps this is close to 90% average yield per step (unoptimized), thus, it constitutes an impressive one-pot procedure for generating such highly substituted phenols. Furthermore, **4b** is not symmetrical and demonstrates that this methodology can be applied to generate relatively complex phenols.

Conclusions

In summary, we have developed a very rapid and green synthesis of phenols from boronic acids *via ipso*-hydroxylation mediated by hydrogen peroxide in ethanol at ambient conditions. The method appears quite general and affords very good to excellent yields across a range of sterically and electronically diverse substituent patterns. By addition of hydrogen bromide to the reaction mixture, diverse bromophenols are available through a tandem hydroxylation/bromination process. The chemistry has been applied to develop a synthesis of highly substituted arylphenols *via* a one-pot *ipso*-hydroxylation/bromination/Suzuki–Miyaura coupling sequence. The results should be of broad interest to the chemical community and represent major advances in the synthesis of complex phenols.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors acknowledge generous funding for this project by the Research Council of Norway (Grant no. 275043 CasCat) and the Department of Chemistry at UiT The Arctic University of Norway.

References

- (a) J. H. P. Tyman, *Synthetic and Natural Phenols*, Elsevier, New York, 1996; (b) Z. Rappoport, *The Chemistry of Phenols*, Wiley-VCH, Weinheim, 2003; (c) S. Quideau, D. Deffieux, C. Douat-Casassus and L. Pouysegou, *Angew. Chem., Int. Ed.*, 2011, **50**, 586–621; (d) Y. Ji, P. Li, X. Zhang and L. Wang, *Org. Biomol. Chem.*, 2013, **11**, 4095–4101; (e) R. W. Owen, A. Giacosa, W. E. Hull, R. Haubner, B. Spiegel-hader and H. Bartsch, *Eur. J. Cancer*, 2000, **36**, 1235–1247.
- (a) K. W. Anderson, T. Ikawa, R. E. Tundel and S. L. Buchwald, *J. Am. Chem. Soc.*, 2006, **128**, 10694; (b) T. Schulz, C. Torborg, B. Schäffner, J. Huang, A. Zapf, R. Kadyrov, A. Börner and M. Beller, *Angew. Chem., Int. Ed.*, 2009, **48**, 918; (c) J. W. Han, J. Jung, Y.-M. Lee, W. Nam and S. Fukuzumi, *Chem. Sci.*, 2017, **8**, 7119; (d) G. Shan, X. Yang, L. Ma and Y. Rao, *Angew. Chem., Int. Ed.*, 2012, **51**, 13070; (e) A. Tlili, N. Xia, F. Monnier and M. Taillefer, *Angew. Chem., Int. Ed.*, 2009, **48**, 8725; (f) L. Yang, Z. Huang, G. Li, W. Zhang, R. Cao, C. Wang, J. Xiao and D. Xue, *Angew. Chem., Int. Ed.*, 2018, **57**, 1968.
- Z. Rappoport, *The Chemistry of Phenols*, Wiley-VCH, Weinheim, 2003.
- J. H. P. Tyman, *Synthetic and Natural Phenols*, Elsevier, New York, 1996.
- W. Wang, N. Li, H. Tang and Y. Ma X. Yang, *Mol. Catal.*, 2018, **453**, 113–120.
- V. M. Zakoshansky, *Pet. Chem.*, 2007, **47**, 273–284.
- J. P. Lambooy, *J. Am. Chem. Soc.*, 1950, **72**, 5327–5328.
- (a) J. Zheng, S. Lin, X. Zhu, B. Jiang, Z. Yang and Z. Pan, *Chem. Commun.*, 2012, **48**, 6235; (b) L. Wang, W. Zhang, D. Sheng Su, X. Meng and F.-S. Xiao, *Chem. Commun.*, 2012, **48**, 5476; (c) T. Begum, A. Gogoi, P. K. Gogoi and U. Bora, *Tetrahedron Lett.*, 2015, **56**, 95; (d) A. D. Chowdhury, S. M. Mobin, S. Mukherjee, S. Bhaduri and G. K. Lahiri, *Eur. J. Inorg. Chem.*, 2011, **2011**, 3232; (e) H. Yang, Y. Li, M. Jiang, J. Wang and H. Fu, *Chem.–Eur. J.*, 2011, **17**, 5652; (f) J. Xu, X. Wang, C. Shao, D. Su, G. Cheng and Y. Hu, *Org. Lett.*, 2010, **12**, 1964; (g) K. Inamoto, K. Nozawa, M. Yonemoto and Y. Kondo, *Chem. Commun.*, 2011, **47**, 11775; (h) N. Gogoi, P. K. Gogoi, G. Borah and U. Bora, *Tetrahedron Lett.*, 2016, **57**, 4050; (i) H. Yi and A. Lei, *Chem.–Eur. J.*, 2017, **23**, 10023–10027.
- (a) Y.-Q. Zou, J.-R. Chen, X.-P. Liu, L.-Q. Lu, R. L. Davis, K. A. Jørgensen and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2012, **51**, 784; (b) T. Toyao, N. Ueno, K. Miyahara, Y. Matsui, T.-H. Kim, Y. Horiuchi, H. Ikeda and M. Matsuoka, *Chem. Commun.*, 2015, **51**, 16103; (c) X. Yu and S. M. Cohen, *Chem. Commun.*, 2015, **51**, 9880; (d) J. A. Johnson, J. Luo, X. Zhang, Y.-S. Chen, M. D. Morton, E. Echeverría, F. E. Torres and J. Zhang, *ACS Catal.*, 2015, **5**, 5283; (e) A. K. Simlandy, B. Bhattacharyya, A. Pandey and S. Mukherjee, *ACS Catal.*, 2018, **8**, 5206; (f) H.-P. Liang, Q. Chen and B.-H. Han, *ACS Catal.*, 2018, **8**, 5313; (g) S. P. Pitre, C. D. McTiernan, H. Ismaili and J. C. Scaiano, *J. Am. Chem. Soc.*, 2013, **135**, 13286; (h) H.-Y. Xie, L.-S. Han, S. Huang, X. Lei, Y. Cheng, W. Zhao, H. Sun, X. Wen and Q.-L. Xu, *J. Org. Chem.*, 2017, **82**, 5236; (i) I. Kumar, R. Sharma, R. Kumar, R. Kumar and U. Sharma, *Adv. Synth. Catal.*, 2018, **360**, 2013; (j) I. K. Sideri, E. Voutyritsa and C. G. Kokotos, *Synlett*, 2018, **29**, 1324–1328.
- (a) A. Gogoi and U. Bora, *Tetrahedron Lett.*, 2013, **54**, 1821; (b) S. Gupta, P. Chaudhary, V. Srivastava and J. Kandasamy, *Tetrahedron Lett.*, 2016, **57**, 2506; (c) P. Shremoyee, M. Abhijit and R. M. Harunar, *Appl. Catal., A*, 2018, **562**, 58–66; (d) E. Saikia, S. Bora and B. Chetia, *RSC Adv.*, 2015, **5**, 102723; (e) J. Simon, S. Salzbrunn, G. K. S. Prakash, N. A. Petasis and G. A. Olah, *J. Org. Chem.*, 2001, **66**, 633–634; (f) G. K. S. Prakash, S. Chacko, C. Panja, T. E. Thomas, L. Gurung, G. Rasul, T. Mathew and G. A. Olah, *Adv. Synth. Catal.*, 2009, **351**, 1567–1574.



- 11 (a) G. A. Molander and L. N. J. Cavalcanti, *J. Org. Chem.*, 2011, **76**, 623; (b) J. J. Molloy, T. A. Clohessy, C. Irving, N. A. Anderson, G. C. Lloyd-Jones and A. J. B. Watson, *Chem. Sci.*, 2017, **8**, 1551.
- 12 C. Zhu, R. Wang and J. R. Falck, *Org. Lett.*, 2012, **14**, 3494.
- 13 N. Chatterjee and A. Goswami, *Tetrahedron Lett.*, 2015, **56**, 1524.
- 14 G. Cheng, X. Zeng and X. Cui, *Synthesis*, 2014, **46**, 295.
- 15 D.-S. Chen and J.-M. Huang, *Synlett*, 2013, **24**, 499.
- 16 (a) G. W. Kabalka, T. M. Shoup and N. M. Goudgaon, *Tetrahedron Lett.*, 1989, **30**, 1483; (b) X. Yang, X. Jiang, W. Wang, Q. Yang, Y. Ma and K. K. Wang, *RSC Adv.*, 2019, **9**, 34529.
- 17 I. Saikia, M. Hazarika, N. Hussian, M. R. Das and C. Tamuly, *Tetrahedron Lett.*, 2017, **58**, 4255–4259.
- 18 For recent examples, see: (a) W.-Z. Weng, H. Liang and B. Zhang, *Org. Lett.*, 2018, **20**, 4979–4983; (b) A. Gualandi, A. Savoini, R. Saporetti, P. Franchi, M. Lucarini and P. G. Cozzi, *Org. Chem. Front.*, 2018, **5**, 1573; (c) I. Muhammad, M. H. Balakrishnan, M. Sasidharan and S. Mannathan, *New J. Chem.*, 2019, **43**, 11065; (d) Y. Fang, R. Zhao, Y. Yao, Y. Liu, D. Chang, M. Yao and L. Shi, *Org. Biomol. Chem.*, 2019, **17**, 7558; (e) G. Latha, N. Devarajan, M. Karthik and P. Suresh, *Catal. Commun.*, 2020, **136**, 105911; (f) C. McCarthy, N. L. Garcia and J. M. Palomo, *ChemistrySelect*, 2020, **5**, 7492–7496; (g) L. Hao, G. Ding, D. A. Deming and Q. Zhang, *Eur. J. Org. Chem.*, 2019, 7307–7321; (h) X. Xu, H. Feng, H. Li and L. Huang, *Eur. J. Org. Chem.*, 2019, 3921–3928; (i) X. Yang, X. Jiang, W. Wang, Q. Yang, Y. Ma and K. Wang, *RSC Adv.*, 2019, **9**, 34529.
- 19 (a) E. J. Shin, S. R. Joo and S. H. Kim, *Tetrahedron Lett.*, 2019, **60**, 1509–1513; (b) E. Saikia, S. J. Bora and B. Chetia, *RSC Adv.*, 2015, **5**, 102723.
- 20 R. H. Szumigala, P. N. Devine, D. R. Gauthier and R. P. Volante, *J. Org. Chem.*, 2004, **69**, 566–569.

