RSC Advances



View Article Online

View Journal | View Issue

PAPER

Check for updates

Cite this: RSC Adv., 2020, 10, 39266

Received 13th October 2020 Accepted 16th October 2020

DOI: 10.1039/d0ra08726b

rsc.li/rsc-advances

Introduction

Crambescins are members of a class of guanidine alkaloids that were isolated from the marine sponges *Crambe crambe*,¹ *Monanchora pulchra*,² *Pseudaxinella reticulate*³ and *Monanchora* n. sp.⁴ To date, over 30 structurally related crambescins have been isolated.^{5,6} Crambescins are classified as three types and are all characterized by at least one chiral cyclic guanidine: type A corresponds to the bicyclic structures; type B to monocyclic ones and type C to spirocyclic frameworks (Fig. 1). Members of the crambescin class exhibit potent biological activity, including inhibition of binding of HIV gp 120 to human CD4,⁷ suppression of the multiplication of cancer cells,⁸ blocking of the voltage-gated sodium channels⁹ and potassium channels,¹⁰ and inhibition of the bacteria growth.¹¹

and Yongbiao Guo 💷*

For this family of natural products, a plethora of biological evaluations were done. Synthesis is necessary to secure the

access to the molecules as the sponges are not widely accessible and to anticipate an alternative production to preserve fragile marine ecosystems. Only three years after crambescin A–C were isolated, the racemic synthesis of crambescin A–C was reported by Snider.¹² More recently, a racemic synthesis of crambescin A was reproted by Thomas and co-workers employing a tethered Biginelli-like reaction between C-2/C-3 activated fatty acids and a central guanidinylated pyrrolinium.¹³ However, the asymmetric synthesis of crambescin A–C has not been reported. It was not until 2016 that the first asymmetric route to crambescin A–C derivatives (crambescin A–C carboxylic acid) was delineated by Nishikawa and co-workers.⁹ A *cis*-enyne (2, Scheme 1) was epoxidized using the Katsuki asymmetric epoxidation to give epoxide (3) and then product recrystallization allowed to reach >99% ee. Epoxide (3) was transformed to the chiral spirocyclic

Eight-step total synthesis of (+)-crambescin A⁺

(+)-Crambescin A belongs to the polycyclic guanidine natural product family and has been shown to

possess various medically important properties. The chiral bicyclic guanidine structure of (+)-crambescin

A presents a challenge for chemical synthesis. Here we implement a novel asymmetric Biginelli reaction

strategy to achieve the enantiospecific total synthesis of (+)-crambescin A in only 8 steps from the

Zhenhua Gao, Junchen Li, Yunyang Song, Xiaojing Bi, Xiangyan Meng*

abundant and inexpensive aliphatic aldehyde, urea and methyl 3-oxobutanoate.



Fig. 1 Structures of crambescin A type, B type and C type.

[†] Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ra08726b



Scheme 1 Nishikawa's total synthesis of (+)-crambescin A–C carboxylic acid.

(0)

State Key Laboratory of NBC Protection for Civilian Research Beijing, 102205, P. R. China. E-mail: van87120@126.com



guanidine structure (5) *via* bromocation-triggered cascade cyclization reaction. Then, subsequent functional group transformations from **6** resulted in the total synthesis of crambescin A carboxylic acid (16 steps from **2**, 0.9% total yield), crambescin B carboxylic acid (15 steps from **2**, 8.7% total yield) and crambescin C carboxylic acid (15 steps from **2**, 4.5% total yield), respectively. Although this achievement is significant, the practicality of this route is questionable because of lower overall yields (0.9–8.7%) and the requirement for a large number of linear synthetic steps (15–16 steps).

As we are interested in the unique structure of the crambescins, we planned to synthesize the crambescins by a novel strategy. Herein, we report an enantioselective total synthesis of (+)-crambescin A (1) based upon a strategy involving successive Biginelli reaction of aliphatic aldehyde.

As an approach to the synthesis of crambescin A (1), it seemed reasonable to couple the bicyclic guanidine carboxylic acid **8** with the side-chain alcohol **9** by means of an esterification at the final stage of the synthesis (Scheme 2). Compound **8** can be synthesized from the optically active intermediate **10** through four steps, including thionation, *S*-ethylation, guanidination and Boc protection. The intermediate **10** can be obtained from the key intermediate **11** by employing the C–C and C–N coupling/cyclization cascade reaction. The key intermediate **11** is a kind of 4-alkyl-3,4-dihydropyrimidin-2(1H)-ones (4alkyl-DHPMs), which can be synthesized by the enantioselective Biginelli reaction of urea, ethyl acetoacetate, and octanal.

Results and discussion

Recently, our group demonstrated that a chiral binaphthol phosphoric acid (**cat.1**, Table 1, entry 1) can successfully catalyze the asymmetric Biginelli reaction of aliphatic aldehydes, urea and β -ketoesters for the synthesis of optically active 4-alkyl-DHPMs.^{14,15} However, the reaction time was too long (at 25 °C for 10 days, then at 50 °C for 1 day). Moreover, the enantiose-lectivity (85% ee, Table 1, entry 1) was not satisfactory. So we commenced our synthetic study by screening more suitable reaction conditions for this particular reaction. Initially, a number of commonly used chiral phosphoric acids (**cat.2** and **cat.3a–e**), were examined in CHCl₃ at 25 °C (Table 1). Pleasingly, it was found that all of the catalysts could indeed promote the desired reaction giving the 4-alkyl-3,4-dihydropyrimidin-2(1H)-

Table 1 Optimization of the enantioselective Biginelli reaction conditions^a



Entry	Catalyst	Solvent	Time (d)	$\operatorname{Yield}^{b}(\%)$	ee ^c (%)
1	cat 1	CHCl	10 + 1	75	85
2	cat.2	CHCl	10 1	73	89
3	cat.3a	CHCl	5	26	46
4	cat.3b	CHCl ₃	5	70	72
5	cat.3c	CHCl ₃	5	39	0
6	cat.3d	CHCl ₃	5	49	15
7	cat.3e	CHCl ₃	5	44	58
8	cat.2	CHCl ₃	7	50	59
9	cat.2	CH_2Cl_2	7	60	90
10	cat.2	CCl_4	7	68	78
11	cat.2	Toluene	7	50	94
12	cat.2	ClCH ₂ CH ₂ Cl	7	73	91
13	cat.2	BrCH ₂ CH ₂ Cl	7	85	91
14	cat.2	Benzene	7	54	82
15	cat.2	Xylene	7	52	81
16^d	cat.2	BrCH ₂ CH ₂ Cl	7	85	91

^{*a*} Reactions conditions: **12** (0.2 mmol), **13** (1.5 mmol), **14** (0.24 mmol) and a catalyst (0.002 mmol) in solvent (2 mL) for 5–10 days (25 °C). ^{*b*} Yield was determined by HPLC analysis. ^{*c*} Determined by HPLC (Chiralcel AD-RH). ^{*d*} The reactions was run with **12** (0.1 mol), **13** (0.75 mol), **14** (0.12 mol) and **cat.2** (0.001 mol) in BrCH₂CH₂Cl (100 mL) at 25 °C for 7 days.

one **6** in variable yields and enantioselectivities with the exception of **cat.3c** (Table 1, entries 1 to 6). With **cat.2** as the best candidate (entry 2), we continued to study the effect of solvents and found that $BrCH_2CH_2Cl$ was the optimal choice (Table 1, entries 13 vs. 2, 8–12, 14–15). The synthetic application potentiality of this asymmetric Biginelli reaction of aliphatic aldehydes was demonstrated by the gram-scale transformation. A 0.1 mol–scale reaction of **12**, **13** and **14** was performed, affording 22.8 g of **11** in 85% yield with 91% ee. There was almost no change in either yield or enantioselectivity. Moreover, ee of **11** could be raised to 99% after a single recrystallization (88% crystallization yield).

Table 2 Optimization of the reaction conditions for the C–C and C–N coupling/cyclization cascade reaction of 6^a



Entry	Catalyst	Solvent	$\operatorname{Yield}^{b}(\%)$	Conversion (%)
1	<i>n</i> -BuLi (3.5)	1,2-Dibromoethane	10	_
2	<i>n</i> -BuLi (3.5)	1-Bromo-2-chloroethane	34	65
3	<i>n</i> -BuLi (4.0)	1-Bromo-2-chloroethane	38	61
4	<i>n</i> -BuLi (5.0)	1-Bromo-2-chloroethane	42	71
5	<i>n</i> -BuLi (6.0)	1-Bromo-2-chloroethane	40	64
6	LDA (5.0)	1-Bromo-2-chloroethane	38	58
7	LiHMDS (5.0)	1-Bromo-2-chloroethane	Trace	—

^{*a*} Reagents and conditions: **11** (0.5 mmol), base (1.75–3.0 mmol), dry THF (5 mL), -20 °C – r.t., 1 h, then 1,2-dibromoethane or 1-bromo-2-chloroethane (2.5 mmol), r.t. 3 h. ^{*b*} Isolated yield.



Scheme 3 Reagents and conditions: (a) $Et_3O^+BF_4^-$, NaHCO₃, dry CH_2Cl_2 , r.t., 1 h; (b) $NH_4^+EtCO_2^-$, 80 °C, 3 h; (c) Lawesson's reagent, xylene, 120 °C, 2 h, 95%.

With large quantities of optically active 4-alkyl-DHPM **11** in hand, we next examined the conversion of **11** into the bicyclic dihydropyrimidinone **10** through the C–C and C–N coupling/ cyclization cascade reaction. Initially, we envisaged that C–C and C–N coupling/cyclization cascade reaction of **11** with 1,2-dibromoethane by the Singh^{16,17} method would afford **10**.



Scheme 4 Reagents and conditions: (a) NaH, dry THF, 0 °C, then DMAP, (Boc)₂O, r.t.; (b) EtSLi, HMPA, r.t., then BOPCl, NEt₃, dry CH₂Cl₂, r.t.; (c) CF₃CO₂H, dry CH₂Cl₂.

However, the yield of bicyclic intermediate **10** was very low (10%). Thus, several bases and electrophiles for this conversion were further investigated, and the results are summarized in Table 2. The best yield (42%, 71% conv.) was obtained when 5.0 equivalents of *n*-BuLi and 5.0 equivalents of 1-bromo-2-chloroethane were used. Control experiments performed by chiral HPLC indicated that no erosion of the ee occurred during the cyclization process.

Compound **10** could be treated directly with ethylation reagent $(Et_3O^+BF_4^-)$, leading to ethoxypyrimidine **15**, which was then converted to bicyclic guanidine **16** *via* guanidination reaction with ammonium propionate. However, the total yield (53%) of two steps was not satisfactory owing to formation of the side product **10** in the guanidination step. So we decided to prevent its formation by substituting sulfur atom for oxygen atom to avoid the side reaction. Dihydropyrimidinthione **17** was generated in quantitative yield by treatment of **10** with Lawesson's reagent. By following the same procedure of **10**, we were pleased that the yield of **16** could be increased to 81% from **10** (Scheme 3).

The last three steps completed the total synthesis of crambescin A (Scheme 4). Treatment of the bicyclic guanidine **16** with (Boc)₂O in the presence of NaH and DMAP gave the fully protected bicyclic guanidine **19**. Then, demethylation of **19** with EtSLi and subsequent condensation with the guanidine alcohol **9** provided the fully protected crambescin A **20**. Finally, CF₃CO₂H promoted removal of the four Boc groups, furnishing the targeted crambescin A (only 8 steps, 10.7% total yield). All of the analytical data including ¹H, ¹³C, 2D NMR, and $[\alpha]_D$ value of the synthetic **1** were identical to those of the natural product (Tables S1 and S2[†]).¹¹

Conclusions

The first asymmetric total synthesis of (+)-crambescin A (1) has been accomplished *via* a short and practical route from urea,

Paper

ethyl acetoacetate, and octanal as the starting material. The key transformations include a chiral phosphoric acid-catalyzed Biginelli reaction and a C-C and C-N coupling/cyclization cascade reaction under strong alkali condition. This study should open up a new efficient route for synthesizing crambescins.

Experimental section

General information

Reagents and solvents were purchased from common commercial suppliers and were used without further purification. Column chromatography was generally performed on silica gel (200-300 mesh). Melting points were determined with a Büchi B-545 melting-point apparatus. 600 MHz ¹H NMR and 150 MHz ¹³C NMR spectra were recorded on Varian VMS-600 spectrometers, respectively. The chemical shifts are reported in ppm (δ scale) relative to internal tetramethylsilane, and coupling constants are reported in hertz (Hz). High-resolution mass spectra (HRMS) were obtained on Agilent 6502 Q-TOF HPLC and mass spectrometry.

Synthesis of 4-alkyl-DHMPs 11

A mixture of urea 12 (0.1 mol), catalyst cat.2 (0.001 mol), 14 (0.12 mol), 13 (0.75 mol) and BrCH₂CH₂Cl (100 mL) was stirred at 25 °C for 7 days. Then the resulting solution was concentrated under reduced pressure, and the orange residue was purified by column chromatography ([eluting with hexane/i-PrOH 15:1)] to afford 11 (22.8 g, 85%) as a white crystalline solid. Then the optically pure 11 were obtained in 88% yield with 99% ee by recrystallization from MeCN. m.p. 128-130 °C; ¹H NMR (600 MHz, CDCl₃): δ (ppm) 8.27 (s, 1H), 5.87 (s, 1H), 4.28 (dt, J =7.8 Hz, 3.6 Hz, 1H), 3.71 (s, 3H), 2.27 (s, 3H), 2.27 (s, 3H) 1.57-1.48 (m, 2H), 1.38–1.25 (m, 10H), 0.87 (t, J = 6.6 Hz, 3H). ¹³ C NMR (150 MHz, CDCl3): δ (ppm) 166.4, 154.6, 147.0, 101.4, 51.6, 51.1, 36.9, 31.8, 29.3, 29.2, 24.4, 22.6, 18.6, 14.1. HRMS (m/z) calcd for $C_{14}H_{25}N_2O_3 [M + H]^+$ 269.1860, found 269.1862; $[\alpha]_D^{25}$ = +140 (c = 0.30, CH₂Cl₂); The ee was determined by HPLC analysis (Chiralpak OD-RH, $CH_3CN/H_2O = 60/40$, v/v, 1.0 mL min⁻¹, 280 nm, $t_{\rm R}$ (minor) = 3.4 min, $t_{\rm R}$ (major) = 5.1 min), 99% ee.

Synthesis of 10

After stirring a solution of 11 (2 mmol) in dry THF (20 mL) under nitrogen atmosphere at -20 °C, 2.4 M nBuLi (4.2 mL, 10 mmol) was added dropwise at -20 °C. After the addition, reaction mixture was warmed to room temperature and stirred for additional 1 h. Then 1-bromo-2-chloroethane (12 mmol) was added dropwise using cannula at room temperature. Then the mixture was stirred at room temperature for 3 h. After which a saturated aqueous solution of NH₄Cl was introduced to terminate the reaction. The reaction was extracted with diethyl ether (3 \times 25 mL) treated in sequence with brine and water, dried over anhydrous Na₂SO₄ and filtered. After the excess solvent was evaporated, the resulting orange gum was purified by column chromatography on a silica gel column [eluting with

hexane/ethyl acetate (3:1)] to obtain 10 (247 mg, 42%) as a colorless oil. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 5.48 (s, 1H), 4.27 (dt, J = 7.2 Hz, 3.6 Hz, 1H), 3.71 (s, 3H), 3.72–3.65 (m, 2H), 3.20 (ddd, J = 18.0 Hz, 8.0 Hz, 4.2 Hz, 1H), 3.00 (dt, J = 18.0 Hz, 8.0 Hz, 1H), 2.06-1.91 (m, 2H), 1.55-1.48 (m, 2H), 1.38-1.24 (m, 10H), 0.86 (t, J = 7.2 Hz, 3H). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 166.2, 153.9, 153.1, 98.4, 51.9, 51.0, 46.6, 37.1, 31.9, 31.8, 29.2, 29.2, 24.4, 22.6, 21.4, 14.1. HRMS (m/z) calcd for $C_{16}H_{27}N_2O_3[M+H]^+$ 295.2016, found 295.2029; $[\alpha]_D^{25} = +146 (c$ = 0.2, CH_2Cl_2 ; the ee was determined by HPLC analysis (Chiralpak OD-H, $CH_3CN/H_2O = 60/40$, v/v, 1.0 mL min⁻¹, 280 nm, $t_{\rm R}$ (minor) = 5.5 min, $t_{\rm R}$ (major) = 6.9 min), 99% ee.

Synthesis of 17

A mixture of 10 (0.8 mmol) and Lawesson's reagent (0.96 mmol) in xylene was heated to 120 °C for 2 h. After the excess solvent was evaporated, the resulting orange gum was purified by column chromatography on a silica gel column [eluting with hexane/EtOAc (4:1)] to obtain 17 (236 mg, 95%) as a colorless oil. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 6.96 (s, 1H), 4.28 (dt, J =7.5, 3.6 Hz, 1H), 4.01 (ddd, *J* = 9.1, 5.5, 3.4 Hz, 2H), 3.73 (s, 3H), 3.30 (ddd, J = 18.2, 8.5, 3.6 Hz, 1H), 2.99 (dt, J = 18.4, 9.3 Hz, 1H), 2.06 (ddt, J = 12.9, 10.8, 4.0 Hz, 1H), 2.00–1.89 (m, 1H), 1.53 (dddd, J = 24.4, 14.1, 8.5, 3.9 Hz, 2H), 1.39–1.25 (m, 10H), 0.87 (t, J = 7.0 Hz, 3H). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 176.3, 165.9, 150.4, 100.4, 52.0, 51.9, 51.4, 36.9, 32.0, 31.8, 29.2, 29.1, 24.1, 22.6, 21.1, 14.1. HRMS (ESI) *m*/*z* calcd for C₁₆H₂₇N₂O₂S [M + H]⁺: 311.1793; found 311.1788. $[\alpha]_D^{25} = +240$ (c = 0.30, CH_2Cl_2).

Synthesis of 18

To a mixture of 17 (0.7 mmol) and NaHCO₃ (2.8 mmol) in dry CH_2Cl_2 (15 mL) was added dropwise $Et_3O^+BF_4^-$ (0.2 M CH_2Cl_2 solution, 7 mL, 1.4 mmol) under nitrogen atmosphere at room temperature. The resulting solution was stirred at room temperature for 0.5 h and then water (30 mL) was added at room temperature. The aqueous layer was extracted with CH_2Cl_2 (2 \times 20 mL). The combined organic layers were washed with brine (50 mL), dried over anhydrous Na₂SO₄ and filtered. After the excess solvent was evaporated, the resulting orange gum was purified by column chromatography on a silica gel column [eluting with hexane/ethyl acetate (6:1)] to obtain 18 (225 mg, 95%) as a colorless oil. ¹H NMR (600 MHz, CDCl₃): $\delta = 4.62$ (s, 1H), 3.72-3.69 (m, 1H), 3.68 (s, 3H), 3.51 (dt, J = 16.8, 8.5 Hz, 1H), 3.21 (ddd, J = 17.9, 8.4, 1.9 Hz, 1H), 3.14 (s, 1H), 2.96 (s, 1H), 2.85 (dt, *J* = 15.9, 9.3 Hz, 1H), 2.07–2.03 (m, 1H), 1.98–1.91 (m, 1H), 1.47–1.39 (m, 3H), 1.34-1.25 (m, 12H), 0.86 (t, J =7.2 Hz, 3H).¹³C NMR (150 MHz, CDCl₃): $\delta = 167.4$, 153.9, 150.5, 97.2, 55.9, 50.8, 48.4, 37.2, 31.9, 30.9, 29.7, 29.6, 29.3, 25.4, 24.7, 22.6, 22.0, 14.3, 14.1. HRMS: m/z calcd for $C_{18}H_{31}N_2O_2S$ [M + H]⁺: 339.2106; found: 339.2114.

Synthesis of 16

A mixture of **19** (0.5 mmol) and ammonium propionate (1.0 g) was stirred at 80 °C under nitrogen atmosphere for 3 h and then the mixture was concentrated under reduced pressure. The resulting orange gum was purified by column chromatography on a silica gel column [eluting with EtOAc/MeOH/TFA (100 : 10 : 1)] to obtain **16** (TFA salt, 183 mg, 90%) as a colorless oil. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 10.12 (s, 1H), 4.43 (s, 1H), 3.87(t, J = 8.4 Hz 1H), 3.75 (s, 3H), 3.69 (dd, J = 11.4 Hz, 8.4 Hz, 1H), 3.33 (ddd, J = 18.0 Hz, 8.4 Hz, 3.0 Hz, 1H), 2.94 (dt, J= 18.6 Hz, 9.6 Hz, 1H), 2.25–2.19 (m, 1H), 2.13–2.06 (m, 1H), 1.58–1.48 (m, 2H), 1.40–1.25 (m, 10H), 0.87 (t, J = 7.2 Hz, 3H). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 165.1, 152.0, 150.4, 102.3, 51.6, 50.1, 47.4, 36.7, 31.7, 30.7, 29.1, 29.0, 23.9, 22.6, 21.8, 14.0. HRMS (m/z) calcd for C₁₆H₂₈N₃O₂ [M + H]⁺ 294.2176, found 294.2177; [α]_D²⁵ = +50 (c = 0.30, MeOH, TFA salt).

Synthesis of 20

To a solution of 16 (0.4 mmol) in THF (5 mL) at 0 °C was added NaH (4 mmol). After stirring for 15 min, DMAP (0.2 mmol) and Boc₂O (2.4 mmol) were added and the solution was allowed to warm to RT. After stirring for 12 h, the solution was quenched with saturated NH₄Cl (5 mL) and then the solution was extracted with diethyl ether $(3 \times 30 \text{ mL})$ treated in sequence with brine and water, dried over anhydrous Na2SO4 and filtered. After the excess solvent was evaporated, the resulting orange gum was purified by column chromatography on a silica gel column [eluting with hexane/ethyl acetate (3:1)] to obtain crude 19 (162 mg, 82%, crude yield) as a colorless oil. Then, to crude 19 (0.328 mol) was added a solution of EtSLi in HMPA (3 mL of a 1.6 M solution). After stirring at 25 °C for 2 h, the reaction was quenched with a saturated solution of NH4Cl (0.3 mL) and diluted with Et2O (50 mL). After washing $H_2O(5 \times 30 \text{ mL})$, the combined aqueous layer was acidified to pH 3 with 0.1 N HCl and extracted with Et_2O (2 × 50 mL). The combined organic layers were then washed with 0.1 N HCl (3 \times 30 mL), dried over anhydrous Na₂SO₄ and filtered. After the excess solvent was evaporated, the crude carboxylic acid was obtained. A mixture of crude carboxylic acid, guanidine alcohol 9 (0.5 mmol), BOPCl (0.5 mmol), Et₃N (15 mmol) and dry CH₂Cl₂ (5 mL) was stirred at 25 °C for 15 h. Then the resulting solution was concentrated under reduced pressure, and the orange residue was purified by column chromatography [eluting with hexane/EtOAc (4:1)] to afford 20 (180 mg, 57%) as a colorless oil. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 11.50 (s, 1H), 8.37 (s, 1H), 5.21 (dd, *J* = 8.4 Hz, 4.2 Hz, 1H), 4.20–4.12 (m, 2H), 3.95 (dd, J = 12.0 Hz, 7.8 Hz, 1H), 3.73 (ddd, J = 12.0 Hz, 8.4 Hz, 4.2 Hz, 1H), 3.49–3.44 (m, 2H), 3.22 (ddd, *J* = 18.0 Hz, 8.4 Hz, 4.2 Hz, 1H), 2.91 (dt, J = 18.0 Hz, 9.0 Hz, 1H), 2.08–2.04 (m, 1H), 2.01–1.95 (m, 1H), 1.75-1.66 (m, 4H), 1.63-1.57 (m, 1H), 1.51 (s, 9H), 1.50 (s, 9H), 1.49 (s, 9H), 1.48 (s, 9H), 1.40-1.36 (m, 1H), 1.30-1.23 (m, 10H), 0.87 (t, J = 7.2 Hz, 3H). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 165.1, 158.9, 156.1, 153.3, 152.0, 150.8, 144.8, 103.0, 83.1(2), 79.7(2), 63.6, 52.9, 48.9, 33.6, 31.9, 31.2, 29.2, 29.1, 28.3(3), 28.2(3), 28.1(3), 28.0(3), 26.2, 25.8, 24.8, 22.6, 21.4, 14.1. HRMS (m/z) calcd for C₄₀H₆₉N₆O₁₀ [M + H]⁺ 793.5070, found 793.5084.

Synthesis of (+)-crambescin A

A mixture of urea **20** (0.2 mol), CF_3CO_2H (2.5 mol) and dry CH_2Cl_2 (7.5 mL) was stirred at 25 °C for 1 h. Then the resulting

solution was concentrated under reduced pressure, and the orange residue was purified by column chromatography [eluting with EtOAc/MeOH/TFA (100 : 10 : 1)] to afford (+)-crambescin A (113 mg, 91%, TFA salt) as a colorless oil. ¹H NMR (600 MHz, CD₃OD): δ (ppm) 4.40 (dd, J = 6.6 Hz, 4.8 Hz, 1H), 4.22 (td, J = 14.4 Hz, 6.6 Hz, 1H), 3.82 (td, J = 8.4 Hz, 1.8 Hz, 1H), 3.67 (dd, J = 16.2 Hz, 9.0 Hz, 1H), 3.35–3.31 (m, 1H), 3.23 (t, J = 7.2 Hz, 2H), 2.99(dt, J = 18.0 Hz, 9.0 Hz, 1H), 2.26–2.18 (m, 1H), 2.15–2.06 (m, 1H), 1.79–1.74 (m, 2H), 1.70–1.66 (m, 2H), 1.62–1.54 (m, 2H), 1.44–1.30 (m, 10H), 0.90 (t, J = 6.6 Hz, 3H). ¹³C NMR (150 MHz, CD₃OD): δ (ppm) 165.9, 158.5, 152.9, 152.6, 103.0, 64.9, 51.1, 48.7, 41.8, 37.2, 32.7, 31.7, 30.1, 30.0, 26.8, 26.4, 25.0, 23.5, 22.7, 14.2. HRMS (m/z) calcd for C₂₀H₃₇N₆O₂ [M + H]⁺ 393.2973, found 393.2987. [α]_D²⁵ = +13 (c = 0.1, MeOH, TFA salt).

Conflicts of interest

There are no conflicts to declare.

References

- 1 R. G. S. Berlinck, J. C. Braekman and D. Daloze, *Tetrahedron lett*, 1990, **31**, 6531–6534.
- 2 T. N. Makarieva, E. K. Ogurtsova and V. A. Denisenko, *Org. Lett.*, 2014, **16**, 4292–4295.
- 3 M. T. Jamison and T. F. Molinski, *J. Nat. Prod.*, 2015, **78**, 557–561.
- 4 A. El-Demerdash, C. Moriou and M. T. Martin, *J. Nat. Prod.*, 2016, **79**, 1929~1937.
- 5 R. G. S. Berlinck and S. Romminger, *Nat. Prod. Rep.*, 2016, 33, 456–490.
- 6 R. G. S. Berlinck, A. F. Bertonha, M. Takaki and J. P. G. Rodriguez, *Nat. Prod. Rep.*, 2017, **34**, 1264–1301.
- 7 A. D. Patil, N. V. Kumar and W. C. Kokke, J. Org. Chem., 1995, 60, 1182–1188.
- 8 M. Roel, J. A. Rubiolo and E. Ternon, *Mar. Drugs*, 2015, **13**, 4633–4653.
- 9 A. Nakazaki, Y. Nakane, Y. Ishikawa, M. Yostsu-Yamashia and T. Nishikawa, *Org. Biomol. Chem.*, 2016, **14**, 5304–5309.
- 10 V. Martín, C. Vale and S. Bondu S, *Chem. Res. Toxicol.*, 2013, 26, 169–178.
- 11 M. T. Jamison and T. F. Molinski, *J. Nat. Prod.*, 2015, **78**, 557–561.
- 12 B. B. Snider and Z. Shi, J. Org. Chem., 1993, 58, 3828-3839.
- S. B. L. Silva, F. Oberhänsli, M. A. Tribalat, G. Genta-Jouve, J. L. Teyssié, M. Y. Dechraoui-Bottein, J. F. Gallard, L. Evanno, E. Poupon and O. P. Thomas, *Angew. Chem.*, *Int. Ed.*, 2019, 58, 520–525.
- 14 Y. Guo, Z. Gao, X. Meng, Y. Huang, H. Zhong, H. Yu, X. Ding, H. Tang and C. Zou, *Synlett*, 2017, 28, 2041–2045.
- 15 Y. Guo, Z. Gao, C. Fan, J. Chen, J. Li, Y. Huang and C. Zou, *Synthesis*, 2018, **50**, 2394–2406.
- 16 K. Singh and S. Singh, J. Org. Chem., 2005, 70, 6114-6117.
- 17 K. Singh and S. Singh, Tetrahedron, 2008, 64, 11718-11723.