



Cite this: *RSC Adv.*, 2020, 10, 16971

DOI: 10.1039/d0ra90045a

rsc.li/rsc-advances

Correction: Surface phosphonation treatment shows dose-dependent enhancement of the bioactivity of polyetheretherketone

Lvhua Liu,^a Yanyan Zheng,^{*a} Qianyu Zhang,^b Lin Yu,^c Ziliang Hu^d and Ying Liu^{*b}

Correction for 'Surface phosphonation treatment shows dose-dependent enhancement of the bioactivity of polyetheretherketone' by Lvhua Liu *et al.*, *RSC Adv.*, 2019, 9, 30076–30086, DOI: 10.1039/C9RA05229A.

The authors regret that an incorrect version of Fig. 9 was included in the original article. The correct version of Fig. 9 is presented below.




Fig. 9 (a) Hard tissue sections of toluidine blue-fuchsin stained around the implant at 12 weeks post-implantation with the white arrows marking the direct bone contact between bone tissue and PEEK substrates. (b) Percentage of bone-implant contact ratios and (c) pull-out load between bone tissue and the PEEK samples after implantation for 12 weeks; $^*(p < 0.05)$ when compared to that of PEEK.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aSchool of Basic Medical Sciences, North Sichuan Medical College, Nanchong, China. E-mail: yanyzheng@163.com

^bDepartment of Stomatology, North Sichuan Medical College, Affiliated Hospital of North Sichuan Medical College, Nanchong, China. E-mail: ying_nsmc@hotmail.com

^cDepartment of Pharmacology, North Sichuan Medical College, Nanchong, China

^dDepartment of Preventive Medicine, North Sichuan Medical College, Nanchong, China

