RSC Advances



View Article Online

CORRECTION

Check for updates

Cite this: RSC Adv., 2020, 10, 16971

Correction: Surface phosphonation treatment shows dose-dependent enhancement of the bioactivity of polyetheretherketone

Lvhua Liu,^a Yanyan Zheng,^{*a} Qianyu Zhang,^b Lin Yu,^c Ziliang Hu^d and Ying Liu^{*b}

DOI: 10.1039/d0ra90045a

rsc.li/rsc-advances

Correction for 'Surface phosphonation treatment shows dose-dependent enhancement of the bioactivity of polyetheretherketone' by Lvhua Liu *et al.*, *RSC Adv.*, 2019, **9**, 30076–30086, DOI: 10.1039/C9RA05229A.

The authors regret that an incorrect version of Fig. 9 was included in the original article. The correct version of Fig. 9 is presented below.



Fig. 9 (a) Hard tissue sections of toluidine blue-fuchsine stained around the implant at 12 weeks post-implantation with the white arrows marking the direct bone contact between bone tissue and PEEK substrates. (b) Percentage of bone-implant contact ratios and (c) pull-out load between bone tissue and the PEEK samples after implantation for 12 weeks; *(p < 0.05) when compared to that of PEEK.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

"School of Basic Medical Sciences, North Sichuan Medical College, Nanchong, China. E-mail: yanyzheng@163.com

^bDepartment of Stomatology, North Sichuan Medical College, Affiliated Hospital of North Sichuan Medical College, Nanchong, China. E-mail: ying_nsmc@hotmail.com ^cDepartment of Pharmacology, North Sichuan Medical College, Nanchong, China

^dDepartment of Preventive Medicine, North Sichuan Medical College, Nanchong, China