

ORGANIC CHEMISTRY

FRONTIERS

RESEARCH ARTICLE

View Article Online
View Journal | View IssueCite this: *Org. Chem. Front.*, 2024,
11, 6609**Electrochemical diversification of cysteine derivatives and cysteine-containing peptides to phosphorothioates and sulfinates†**Longyu Xiao,^a Yifan Li,^{a,c} Jun Huang,^a Li Pan,^b Pan Wu^b and Yue Weng^{id} *^a

This study explores the application of modified cysteine in oligopeptides to overcome challenges in amino acid residue modification, with a focus on cysteine. Novel strategies for the functionalization of biomolecules are presented, with mechanistic investigations indicating the involvement of free radical processes, structural substitution, and nucleophilic displacement. Notably, the gram-scale synthesis of modified cysteine is achieved with high yields, highlighting its potential as a novel anti-fungus agent in biomedical and agrochemical research.

Received 15th August 2024,
Accepted 27th September 2024

DOI: 10.1039/d4qo01502a

rsc.li/frontiers-organic

Introduction

In the fields of chemical biology, biomedicine, and pesticide science, mild chemical reactions have gradually become key tools for modifying biomolecules.^{1–3} These reactions typically exhibit high efficiency, strong specificity, and mild conditions, allowing them to proceed in complex biological environments without disrupting the natural structure and function of biomolecules.^{4,5} For example, click chemistry reactions such as CuAAC (copper-catalyzed azide–alkyne cycloaddition)⁶ and SPAAC (strain-promoted azide–alkyne cycloaddition)⁷ have been widely used for the labeling and modification of macromolecules such as proteins and nucleic acids.^{8,9} These methods not only enhance the targeting and stability of drugs but also provide new means for studying the functional mechanisms of biomolecules.¹⁰ Modification of amino acid residues plays a crucial role in regulating protein functions and biological processes.¹¹ Through chemical modifications, the structure and activity of proteins can be precisely controlled to investigate their functional mechanisms.¹² Common modifications include phosphorylation,¹³ acetylation,¹⁴ methylation,¹⁵ and ubiquitination,¹⁶ which can affect protein stability, localization, interactions, and

enzymatic activity. Currently, mild chemical reactions, such as bio-orthogonal chemistry reactions,¹⁷ have been widely applied to the specific modification of amino acid residues.¹⁸ These reactions can proceed in complex biological environments without disrupting the natural structure of proteins.¹⁹ For example, the use of 4-nitrophenyl phosphate allows for the specific modification of serine or threonine residues.²⁰ These studies not only provide new tools for protein engineering but also offer new perspectives for disease treatment, such as the development of novel drugs²¹ that target and modify pathological proteins. The modification of amino acid residues²² has become an important research direction in chemical biology and drug development.²³

The modification of cysteine²⁴ holds significant importance in protein function regulation and biomedical research. Cysteine contains a reactive thiol group (–SH) that can participate in various crucial biochemical reactions.^{25,26} By modifying cysteine, the structure, stability, and function of proteins could be regulated.^{27,28} For instance, the formation of disulfide bonds involving cysteine plays a key role in protein folding and stability.^{29,30} Additionally, cysteine modifications can regulate enzyme activity,^{31,32} signal transduction, and the cell cycle.³³ Currently, mild chemical reactions are widely used for the specific modification of cysteine,^{34,35} employing reagents such as maleimide and iodoacetamide. These methods not only provide powerful tools for studying protein function and interactions but also show great potential in developing novel therapeutic approaches, such as designing targeted anticancer drugs through cysteine modification (Fig. 1).

In summary, cysteine modification plays an indispensable role in modern biochemistry and drug development. In recent years, electrochemical methods have made remarkable progress in the field of biomolecule modification. These methods utilize the efficiency and precision of electrochemical reactions

^aHubei Key Laboratory of Precision Manufacturing for Small-molecular Active Pharmaceutical Ingredients, School of Chemistry and Chemical Engineering, Hubei University, Wuhan, 430062, P. R. China. E-mail: wengyue@hubu.edu.cn

^bState Key Laboratory of Biocatalysis and Enzyme Engineering, School of Life Sciences, Hubei University, Wuhan, 430062, P. R. China

^cThe Key Laboratory for Chemical Biology of Fujian Province, The MOE Key Laboratory of Spectrochemical Analysis & Instrumentation, and Department of Chemical Biology, College of Chemistry and Chemical Engineering, Xiamen University, Xiamen 361005, China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4qo01502a>




Fig. 1 Electrocatalytic S–P and S(O)–O bonding reaction.

to achieve specific modifications of biomolecules such as peptides and proteins under mild conditions. For instance, electrochemical redox reactions can selectively modify serine and tyrosine residues containing hydroxyl groups. Such modifications not only regulate the function and structure of biomolecules but also introduce functional tags for detection and imaging. The advancement of these technologies provides powerful tools for biomolecule functional research and drug development, pushing the frontiers of chemical biology, biomedicine, and pesticide science. Therefore, electrochemical strategies and bioconjugation processes have been designed to achieve highly selective and efficient modification of cysteine residues.

Results and discussion


Here, two novel strategies are presented for the bioconjugation of cysteine-containing biomolecules, employing electrochemical synthetic chemistry under simple, mild, and environmentally friendly conditions. This straightforward method involves the anodic oxidation of cysteine to generate free radicals, followed by cross-coupling with diethyl phosphite or methanol to form S–P or S(O)–O bonds, resulting in the formation of phosphorothioates and sulfonates. Next, the optimal reaction conditions for Boc-Cys-OMe (**1a**) with phosphite (**2a**) were explored. As shown in Table 1, using platinum electrodes for both the anode and cathode, with ${}^n\text{Bu}_4\text{NI}$ as the electrolyte, and conducting the reaction at a current of 12 mA for 1.5 hours, the amino acid-modified product **3a** was obtained with an 87% yield (Table 1, entry 1). The effect of current intensity was also investigated, revealing that maintaining a steady current is crucial for the reaction (Table 1, entries 2 and 3). The impact of different electrolytes on the yield was then examined (Table 1, entries 4–7), revealing that ${}^n\text{Bu}_4\text{NBr}$ reduced the yield, while ${}^n\text{Bu}_4\text{NBF}_4$ led to a significant decline. Neither KI nor the absence of an electrolyte resulted in any reaction. Choosing an appropriate solvent is also critical. Using methanol or dichloromethane significantly lowered the yield (Table 1, entries 8 and 9). Among the electrodes tested, the dual platinum electrode system provided the best results (Table 1, entries 10–12). The effect of the reaction atmosphere is also noteworthy; yields dropped significantly under nitrogen or oxygen (Table 1, entries 13 and 14). Interestingly, TLC ana-

Table 1 Optimization of the reaction conditions^a

Entry	Variation from the standard conditions ^a	Yield (%)
1	None	87
2	9 mA instead of 12 mA	74
3	15 mA instead of 12 mA	48
4	${}^n\text{Bu}_4\text{NBF}_4$ instead of ${}^n\text{Bu}_4\text{NI}$	16
5	${}^n\text{Bu}_4\text{NBr}$ instead of ${}^n\text{Bu}_4\text{NI}$	72
6	KI instead of ${}^n\text{Bu}_4\text{NI}$	N.D
7	Without ${}^n\text{Bu}_4\text{NI}$	N.R
8	CH_3OH instead of CH_3CN	Trace
9	CH_2Cl_2 instead of CH_3CN	53
10	C(+), Pt(–) as the electrodes	64
11	C(+), C(–) as the electrodes	51
12	Pt(+), Ni(–) as the electrodes	73
13	Under N_2	43
14	Under O_2	34
15	No electricity	N.R

^a Reaction conditions: undivided cell, Pt anode, Pt cathode, **1a** (0.2 mmol), **2a** (0.4 mmol), ${}^n\text{Bu}_4\text{NI}$ (0.2 mmol), MeCN (8 mL), air, room temperature, 12 mA, 1.5 h. Yield of isolated products. N.D = not detected, N.R = not reaction.

lysis indicated the formation of an “unexpected” compound when methanol was used as the solvent (Table 1, entry 8). The initially low yield was attributed to competing side reactions.




Scheme 1 Substrate scope of alcohols and phosphites. ^a Reaction conditions: undivided cell, Pt anode, Pt cathode, **1a** (0.2 mmol), phosphite ester (0.4 mmol), ${}^n\text{Bu}_4\text{NI}$ (0.2 mmol), MeCN (8 mL), air, room temperature, 12 mA, 1.5 h. ^b React 1 h. ^c Reaction condition: undivided cell, Pt anode, Pt cathode, **1a** (0.2 mmol), alcohol (5.0 mmol), ${}^n\text{Bu}_4\text{NBF}_4$ (0.2 mmol), CH_2Cl_2 (6 mL), H_2O (10 μL), air, room temperature, 8 mA, 4.0 h. Yield of isolated products.

Upon reacting Boc-Cys-OMe with methanol, an unforeseen product was obtained with a yield of 78% (Table S1,† entry 7). The optimization of the reaction conditions for Boc-Cys-OMe and methanol is summarized in Table S1.†


The applicability of the reaction with various phosphites or alcohols was subsequently explored. It was observed that phosphites with shorter carbon chains in the ester group led to phosphorothioates with higher yields (Scheme 1, **3a–3b**). When using phosphites with longer carbon chains, the yields decreased (Scheme 1, **3c–3d**). However, when the ester group contained a relatively stable conjugated structure, the target product was obtained with higher yields (Scheme 1, **3e**). The effect of electron-withdrawing groups on the reaction was also investigated. It was found that product **3f** was obtained in low yields when bis(2,2,2-trifluoroethyl) phosphonate was utilized in this protocol (Scheme 1, **3f**). Additionally, pentaacetyl-5-thio-D-glucose, containing a thiol group, was reacted with diethyl phosphite, resulting in a considerable yield (Scheme 1, **3g**). However, when alcohols reacted with Boc-Cys-OMe, the spatial structure had a significant impact on the reaction. When the carbon atoms in the main chain of the alcohol exceeded two, the alcohol could not react with Boc-Cys-OMe. Nevertheless, when using alcohols with shorter carbon chains, the sulfinates were obtained with moderate to high yields

(Scheme 1, **6a–6c**). It is noteworthy that all the electrochemical reactions had excellent isolated yields, with no residual self-coupled cystine.


Furthermore, to investigate the selectivity of Boc-Cys-OMe labeling and its tolerance towards polypeptide labeling, various dipeptides containing Boc-Cys-OMe were introduced into the reaction system. These dipeptides can serve as protein components present in various tissues and cells. Subsequently, diethyl phosphite or methanol was used as a coupling reagent to evaluate the selectivity and tolerance of the Boc-Cys-OMe reaction on dipeptides. Excitingly, these dipeptides met the requirements for modifying the target molecules onto the peptide chains (Scheme 2, **4a–4j**, and **7a–7f**). Surprisingly, dipeptides formed with relatively stable amino acids such as glycine, alanine, valine, leucine, and isoleucine, resulted in the target products with relatively low yields (Scheme 2, **4a**, **4c**, **4d**, **4i**, **7a**, **7c**, **7d**). In contrast, dipeptides formed with amino acids containing reactive groups, such as phenylalanine (containing a phenyl group), lysine (containing an amino group), glutamic acid (containing a carboxyl group), methionine (containing a methylthio group), and tryptophan (containing an indole structure), exhibited better selectivity and group tolerance (Scheme 2, **4b**, **4e**, **4f**, **4g**, **4h**, **4j**,



Scheme 2 Substrate scope of dipeptides. ^a Reaction conditions: undivided cell, Pt anode, Pt cathode, **3** (0.2 mmol), diethyl phosphite (0.4 mmol), $^t\text{Bu}_4\text{NI}$ (0.2 mmol), MeCN (8 mL), air, rt, 12 mA, 1.5 h. ^b Reaction conditions: undivided cell, Pt anode, Pt cathode, **3** (0.2 mmol), methyl alcohol (5.0 mmol), $^t\text{Bu}_4\text{NBF}_4$ (0.2 mmol), CH_2Cl_2 (6 mL), H_2O (10 μL), air, rt, 8 mA, 1.5–8.0 h. Yield of isolated products.



Scheme 3 Electrochemical synthesis of phosphorothioates and sulfinates of endogenous peptides.




Scheme 4 Gram-scale reaction.


7b, 7e, 7f). Electrocatalytic bioconjugation often exhibits better selectivity and tolerance to reactive groups, especially the bioconjugation of dipeptides containing reactive groups with high

selectivity and group tolerance is of great significance for electrochemical bioconjugation.

To synthesize biomolecules, particularly polypeptides, *via* electrochemical methods, the applicability of electrocatalytic synthesis to polypeptides must be assessed. Building on the electrochemical synthesis of dipeptides, further research was initiated, beginning with the electrolysis of annexin inhibitory peptides with exposed amino and hydroxyl groups, cell adhesion peptides with exposed carboxyl, hydroxyl, and imidazole groups, and HPV16 E7-derived peptides with exposed hydroxyl groups. These peptides contain cysteine at the N-terminus or C-terminus and have certain biological activities. It is worth noting that some of these endogenous peptides containing cysteine successfully reacted to produce phosphorothioates and sulfinates (Scheme 3, 8a–8d). The successful labeling of these endogenous peptides in a short period of time indicates that the reaction has good site selectivity and functional group tolerance.



Scheme 5 (a) Anti-fungus experiment; (b) optical density of treated fungal patches at 600 nm (OD₆₀₀).



Scheme 6 Control reaction.

To demonstrate the industrial application value of this reaction, conducted gram-scale reactions: these included the reactions of Boc-Cys-OMe with diethyl phosphite and Boc-Cys-OMe with methanol (Scheme 4). A total of 1.48 grams of the target product **3a** was obtained with an 80% yield, and 1.10 grams of the target product **6a** was produced with a 78% yield. This experiment highlights the great potential of cysteine modification method in subsequent anti-fungus research and new drug synthesis.

The antifungal efficacy of compounds **1a** and **3a** against the pathogenic fungus *Magnaporthe grisea* was also investigated (Scheme 5). To ensure the activity and consistency of the experimental materials (Scheme 5a), the target strain was first activated and cultured in LB liquid medium for 48 hours to reach the logarithmic growth phase. The activated *Magnaporthe grisea* strains were then inoculated into 100 mL of molten potato dextrose agar (PDA) medium, ensuring uniform distribution through thorough mixing. Using standardized 96-well plates as the experimental platform, three parallel samples were prepared for each group to enhance data reliability and reproducibility. Each well was filled with 200 μL of PDA solid medium containing *Magnaporthe grisea*. To assess the inhibitory effects of the compounds on fungal

growth, 20 μL of the antibacterial solutions of **1a** and **3a** were introduced into specific wells, while negative control groups using methanol and pure water were included to rule out solvent effects. All treated plates were incubated at a constant temperature of 28 $^{\circ}\text{C}$ for 3–5 days to fully observe the potential effects of each anti-fungus substance. Then, the optical density at 600 nm (OD_{600}) of each sample was determined (Scheme 5b). The experimental results show that Boc-Cys-OMe promotes the growth of *Magnaporthe grisea*, while the electrochemical reaction product **3a** demonstrates a notable inhibitory effect on the fungus. These findings have potential implications for the development of bioactive molecules with fungicidal properties in agricultural chemistry.

To gain deeper insight into the electrochemical reaction mechanism, a series of control experiments were designed. Under the established conditions, Boc-Cys-OMe was allowed to react with diethyl phosphite for 30 minutes, yielding the intermediate cystine **E** with a 78% yield, along with a 6% yield of the target product. The reaction was then continued for an additional hour, during which intermediate **E** reacted with diethyl phosphite, producing the product **3a** with a 73% yield. This preliminary analysis suggests that the reaction proceeds by first forming cystine from Boc-Cys-OMe, which sub-




Fig. 2 (1) Cyclic voltammograms of reactants and the mixtures in 0.1 M $^n\text{Bu}_4\text{NI}/\text{CH}_3\text{CN}$ using a glassy carbon working electrode (diameter, 3 mm), oxidation peak (a), reduction peak (b). Pt electrode as counter electrode; Ag/AgCl as reference electrode, at 100 mV s^{-1} scan rate. Boc-Cys-OMe (8 mmol L^{-1}), diethyl phosphite (16 mmol L^{-1}). (2) Electron paramagnetic resonance (EPR) experiments of Boc-Cys-OMe and $^n\text{Bu}_4\text{NI}$. (3) Proposed mechanism.

sequently reacts with diethyl phosphite to yield the product **3a** (Scheme 6A). When the radical scavenger TEMPO was introduced into the reaction system under the same conditions, a significant reduction in yield was observed, indicating the involvement of radical intermediates in the reaction.

Furthermore, Boc-Cys-OMe was reacted with methanol for 150 minutes, again producing the cystine intermediate with a high yield, along with another intermediate, **5aa**. This observation explains the reduced yields of products **3a** and **6a** when Boc-Cys-OMe, methanol, and diethyl phosphite are present in the same system. Similarly, when intermediate **E** was reacted with methanol for 4 hours, the product **6a** was obtained with a 93% yield. Isotope labeling experiments were also conducted, confirming that the oxygen in the S=O bond originates from water (Scheme 6B).

Based on controlled experiments, a more in-depth investigation of the reaction mechanism was conducted using cyclic voltammetry. As shown in Fig. 2, ${}^{137}\text{Bu}_4\text{NI}$ is oxidized at 0.481 V and 1.062 V, and reduced at -1.102 V (Fig. 2 (**1a**), red curve). This corresponds to the oxidation of I^- in ${}^{137}\text{Bu}_4\text{NI}$ to iodine radicals, which subsequently convert to I_2 , followed by a disproportionation reaction to form I^- and I^+ . Boc-Cys-OMe is oxidized at 1.308 V (Fig. 2 (**1a**), blue curve), corresponding to the formation of Boc-Cys-OMe radicals. A mixture of Boc-Cys-OMe and diethyl phosphite exhibits a pronounced oxidation peak at 1.287 V, positioned between the oxidation peaks of ${}^{137}\text{Bu}_4\text{NI}$ and cysteine, indicating chemical interactions among the three compounds. To further confirm the presence of radical intermediates in the reaction, electron paramagnetic resonance (EPR) experiments were conducted (Fig. 2 (2)). The results revealed signals corresponding to the radicals of ${}^{137}\text{Bu}_4\text{NI}$ and Boc-Cys-OMe.

Based on the aforementioned study and previous reports,^{36–38} a possible mechanism for the phosphorylation of cysteine is presented (Fig. 2 (3)). Firstly, I^- in ${}^{137}\text{Bu}_4\text{NI}$ is oxidized to iodine radical at the anode. Iodine radical dimerizes to form I_2 , which disproportionates to yield I^- and I^+ . I^+ then undergoes electrophilic substitution with diethyl phosphite (**2a**) to produce diethyl iodophosphate (**I**). Simultaneously, cysteine is oxidized at the anode to generate cysteine radical (**II**). The cysteine radical dimerizes to form cystine, which is subsequently reduced at the cathode to form cystine radical anion (**III**). These cystine radical anion then convert back to cysteine radical and cysteine anion (**IV**). The cysteine anion then undergoes nucleophilic reaction with diethyl iodophosphate (**I**), ultimately yielding **3a**.

Conclusions

In conclusion, a mild and efficient method for the modification and labeling of cysteine and cysteine-containing peptides *via* electrocatalysis has been developed. This approach offers excellent selectivity and functional group tolerance while maintaining operational simplicity. Importantly, it avoids the use of metals, strong acids, or bases, making it environmen-

tally friendly. Through a combination of control experiments, isotope labeling, cyclic voltammetry, and electron paramagnetic resonance (EPR) studies, a plausible reaction mechanism has been proposed. This green modification strategy significantly broadens the scope of cysteine modifications. Additionally, the modified products demonstrated notable antifungal activity against the pathogenic fungus *Magnaporthe grisea*. Looking ahead, this methodology is expected to contribute to advancements in antifungal and agrochemical research.

Author contributions

There are no conflicts to declare.

Data availability

All relevant data are within the manuscript and its ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Mr Heming Qi College of Resources and Environmental Science, South-Central University for Nationalities for the help of EPR experiment. National Natural Science Foundation of China (Grant No. 22301069, 31900051); National Natural Science Foundation of China (Grant No. 22271085).

References

- 1 Y.-C. Song, S. Ingram, R. E. Arbon, D. O. Topping, D. R. Glowacki and J. P. Reid, Transient cavity dynamics and divergence from the Stokes–Einstein equation in organic aerosol, *Chem. Sci.*, 2020, **11**, 2999–3006.
- 2 C. Wan, R. Sun, W. Xia, H. Jiang, B.-X. Chen, P.-C. Kuo, W.-R. Zhang, G. Yang, D. Li, C.-W. Chiang and Y. Weng, Electrochemical Bioconjugation of Tryptophan Residues: A Strategy for Peptide Modification, *Org. Lett.*, 2024, **26**, 5447–5452.
- 3 D. Alvarez-Dorta, C. Thobie-Gautier, M. Croyal, M. Bouzelha, M. Mével, D. Deniaud, M. Boujtita and S. G. Gouin, Electrochemically Promoted Tyrosine-Click-Chemistry for Protein Labeling, *J. Am. Chem. Soc.*, 2018, **140**, 17120–17126.
- 4 P. P. Singh, P. K. Singh and V. Srivastava, Visible light metallaphotoredox catalysis in the late-stage functionalization of pharmaceutically potent compounds, *Org. Chem. Front.*, 2023, **10**, 216–236.

- 5 Y. Weng, C. Song, C.-W. Chiang and A. Lei, Single electron transfer-based peptide/protein bioconjugations driven by biocompatible energy input, *Commun. Chem.*, 2020, **3**, 171.
- 6 M. Z. C. Hatit, L. F. Reichenbach, J. M. Tobin, F. Vilela, G. A. Burley and A. J. B. Watson, A flow platform for degradation-free CuAAC bioconjugation, *Nat. Commun.*, 2018, **9**, 4021.
- 7 Y. Xie, T. L. Lopez-Silva and J. P. Schneider, Hydrophilic Azide-Containing Amino Acid to Enhance the Solubility of Peptides for SPAAC Reactions, *Org. Lett.*, 2022, **24**, 7378–7382.
- 8 C. S. Swenson, A. Velusamy, H. S. Argueta-Gonzalez and J. M. Heemstra, Bilingual Peptide Nucleic Acids: Encoding the Languages of Nucleic Acids and Proteins in a Single Self-Assembling Biopolymer, *J. Am. Chem. Soc.*, 2019, **141**, 19038–19047.
- 9 C. Loynd, S. J. Singha Roy, V. J. Ovalle, S. E. Canarelli, A. Mondal, D. Jewel, E. D. Ficaretta, E. Weerapana and A. Chatterjee, Electrochemical labelling of hydroxyindoles with chemoselectivity for site-specific protein bioconjugation, *Nat. Chem.*, 2024, **16**, 389–397.
- 10 S. Jiang, L. Xiao, L. Pan, Q. Huang, F. Huo, M. Gao, C. Lu, P. Wu and Y. Weng, Electro-induced O–S bonding reaction targeting biological macromolecules, *Org. Chem. Front.*, 2024, **11**, 1090–1096.
- 11 R. Bag and N. K. Sharma, Pd-catalyzed site-selective C(sp²)–H chalcogenation of amino acids and peptides using a picolinamide auxiliary, *Org. Chem. Front.*, 2023, **10**, 1252–1262.
- 12 C.-F. Lee, R. Bai, K.-C. Liu, Z.-W. Chen, A. Gurjar and S. S. Badsara, Cs₂CO₃-Mediated synthetic strategy for iprobenfos derivatives via thiophilic addition of H-phosphites on in situ generated thioaldehydes, *ARKIVOC*, 2024, **2**, 202312055.
- 13 B. Huang, Y. Liu, H. Yao and Y. Zhao, NMR-based investigation into protein phosphorylation, *Int. J. Biol. Macromol.*, 2020, **145**, 53–63.
- 14 J. Baeza, M. J. Smallegan and J. M. Denu, Mechanisms and Dynamics of Protein Acetylation in Mitochondria, *Trends Biochem. Sci.*, 2016, **41**, 231–244.
- 15 M. E. Jakobsson, Enzymology and significance of protein histidine methylation, *J. Biol. Chem.*, 2021, **297**, 101130.
- 16 W. Gui, G. A. Davidson and Z. Zhuang, Chemical methods for protein site-specific ubiquitination, *RSC Chem. Biol.*, 2021, **2**, 450–467.
- 17 J. Li and P. R. Chen, Development and application of bond cleavage reactions in bioorthogonal chemistry, *Nat. Chem. Biol.*, 2016, **12**, 129–137.
- 18 Y. Zheng, W. Lu, T. Ma and S. Huang, Recent advances in photochemical and electrochemical strategies for the synthesis of sulfonyl fluorides, *Org. Chem. Front.*, 2024, **11**, 217–235.
- 19 Z. Jiang, N. Liu, D. Hu, G. Dong, Z. Miao, J. Yao, H. He, Y. Jiang, W. Zhang, Y. Wang and C. Sheng, The discovery of novel antifungal scaffolds by structural simplification of the natural product sampangine, *Chem. Commun.*, 2015, **51**, 14648–14651.
- 20 S. Mukherjee, Y.-H. Hao and K. Orth, A newly discovered post-translational modification – the acetylation of serine and threonine residues, *Trends Biochem. Sci.*, 2007, **32**, 210–216.
- 21 B. Permanne, A. Sand, S. Ousson, M. Nény, J. Hantson, R. Schubert, C. Wiessner, A. Quattropiani and D. Beher, O-GlcNAcase Inhibitor ASN90 is a Multimodal Drug Candidate for Tau and α -Synuclein Proteinopathies, *ACS Chem. Neurosci.*, 2022, **13**, 1296–1314.
- 22 M. N. Maas, J. C. J. Hintzen and J. Mecinović, Probing lysine posttranslational modifications by unnatural amino acids, *Chem. Commun.*, 2022, **58**, 7216–7231.
- 23 Z.-W. Chen, A. Pratheepkumar, R. Bai, Y. Hu, S. S. Badsara, K.-W. Huang and C.-F. Lee, Cesium carbonate-catalyzed synthesis of phosphorothioates via S-phosphination of thioketones, *Chem. Commun.*, 2022, **58**, 11001–11004.
- 24 S. A. Byrne, M. J. Bedding, L. Corcilius, D. J. Ford, Y. Zhong, C. Franck, M. Larance, J. P. Mackay and R. J. Payne, Late-stage modification of peptides and proteins at cysteine with diaryliodonium salts, *Chem. Sci.*, 2021, **12**, 14159–14166.
- 25 K. Jiang, Y. Liu, Y. Yan, S. Wang, L. Liu and W. Yang, Combined chain- and step-growth dispersion polymerization toward PSt particles with soft, clickable patches, *Polym. Chem.*, 2017, **8**, 1404–1416.
- 26 M. H. Asim, I. Nazir, A. Jalil, F. Laffleur, B. Matuszczak and A. Bernkop-Schnürch, Per-6-Thiolated Cyclodextrins: A Novel Type of Permeation Enhancing Excipients for BCS Class IV Drugs, *ACS Appl. Mater.*, 2020, **12**, 7942–7950.
- 27 P. C. Roehm and J. M. Berg, Selectivity of Methylation of Metal-Bound Cysteines and Its Consequences, *J. Am. Chem. Soc.*, 1998, **120**, 13083–13087.
- 28 H. Seki, S. J. Walsh, J. D. Bargh, J. S. Parker, J. Carroll and D. R. Spring, Rapid and robust cysteine bioconjugation with vinylheteroarenes, *Chem. Sci.*, 2021, **12**, 9060–9068.
- 29 R. Chang, M. Gruebele and D. E. Leckband, Protein Folding Stability and Kinetics in Alginate Hydrogels, *Biomacromolecules*, 2023, **24**, 5245–5254.
- 30 N. G. Jayaprakash and A. Surolia, Role of glycosylation in nucleating protein folding and stability, *Biochem. J.*, 2017, **474**, 2333–2347.
- 31 Y. Zhang, Y. Deng, C. Wang, L. Li, L. Xu, Y. Yu and X. Su, Probing and regulating the activity of cellular enzymes by using DNA tetrahedron nanostructures, *Chem. Sci.*, 2019, **10**, 5959–5966.
- 32 T. Kurakawa, N. Ueda, M. Maekawa, K. Kobayashi, M. Kojima, Y. Nagato, H. Sakakibara and J. Kyojuka, Direct control of shoot meristem activity by a cytokinin-activating enzyme, *Nature*, 2007, **445**, 652–655.
- 33 E. V. Vinogradova, C. Zhang, A. M. Spokoiny, B. L. Pentelute and S. L. Buchwald, Organometallic palladium reagents for cysteine bioconjugation, *Nature*, 2015, **526**, 687–691.
- 34 J. Yu, X. Yang, Y. Sun and Z. Yin, Highly Reactive and Tracelessly Cleavable Cysteine-Specific Modification of Proteins via 4-Substituted Cyclopentenone, *Angew. Chem., Int. Ed.*, 2018, **57**, 11598–11602.

- 35 Y.-C. Liu and C.-F. Lee, N-Chlorosuccinimide-promoted synthesis of thiophosphates from thiols and phosphonates under mild conditions, *Green Chem.*, 2014, **16**, 357–364.
- 36 C.-Y. Li, Y.-C. Liu, Y.-X. Li, D. M. Reddy and C.-F. Lee, Electrochemical Dehydrogenative Phosphorylation of Thiols, *Org. Lett.*, 2019, **21**, 7833–7836.
- 37 L. Deng, Y. Wang, H. Mei, Y. Pan and J. Han, Electrochemical Dehydrogenative Phosphorylation of Alcohols for the Synthesis of Organophosphinates, *J. Org. Chem.*, 2019, **84**, 949–956.
- 38 F. Xu, Y.-J. Li, C. Huang and H.-C. Xu, Ruthenium-Catalyzed Electrochemical Dehydrogenative Alkyne Annulation, *ACS Catal.*, 2018, **8**, 3820–3824.