



Cite this: *Chem. Sci.*, 2020, **11**, 1542

All publication charges for this article have been paid for by the Royal Society of Chemistry

## Borane-induced ring closure reaction of oligomethylene-linked bis-allenes†

Xin Tao, Karel Škoch, Constantin G. Daniliuc, Gerald Kehr and Gerhard Erker\*

The trimethylene-linked bis-allene **3a** reacts with Piers' borane [HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>] by a hydroboration/allylboration sequence to generate the cyclization product **5a**. Its pyridine adduct was isolated and characterized by X-ray diffraction. Compound **5a** undergoes a typical frustrated Lewis pair 1,2-P/B alkene addition reaction with PPh<sub>3</sub> to give the heterobicyclic bridged olefinic zwitterionic product **9a**. The tetramethylene-linked bis-allene **3b** and its phenylene annulated analogue **3c** react with HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> to give the analogous seven-membered ring products **5b,c** under mild conditions. The cyclization product **5a** undergoes a series of sequential allylboration reactions with two equivalents of allene followed by ring-closure to give the four-component coupling product **12a**. It undergoes FLP addition to an exo-methylene group upon treatment with PPh<sub>3</sub>. Compound **12a** is oxidatively converted to the boron-free alcohol.

Received 3rd August 2019  
Accepted 17th December 2019


DOI: 10.1039/c9sc03870a

rsc.li/chemical-science

## Introduction

Allenes are important building blocks in organic synthesis.<sup>1</sup> They show interesting and useful stereochemical properties.<sup>2</sup> We had recently shown that the borane [HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>]<sup>3</sup> catalyses the cyclotrimerization of allene as well as a small series of mono-alkyl-substituted allenes **1** to selectively give the respective 1,3,5-trimethylene cyclohexanes **2** as single isomers under metal-free conditions (Scheme 1).<sup>1,4,5</sup>

There is a rich cyclization chemistry of bis-allenes reported in the literature (see Chart 1). Systems **I** (mostly with X = NTs, less frequently CR<sub>2</sub> or O) were reported to rearrange to **II**



Scheme 1 HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>-catalyzed cyclotrimerization of alkyl-substituted allenes.

Organisch-Chemisches Institut, Westfälische Wilhelms-Universität Münster, Corrensstraße 40, 48149 Münster, Germany. E-mail: erker@uni-muenster.de

† Electronic supplementary information (ESI) available: Additional experimental details, further spectral and crystallographic data. CCDC deposition numbers are 1922906–1922913 and 1957134. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc03870a

thermally induced.<sup>6a</sup> They added R<sub>3</sub>Si-SnBu<sub>3</sub>/or -GeR<sub>3</sub> reagents Pd(0) catalysed or radical induced to give the products **III**.<sup>6b,c</sup> With H<sub>2</sub>NR nucleophiles cyclization to medium-sized rings (*e.g.* **IV**) was reported.<sup>6d</sup> The products **V** and **VI** of internal [2 + 2] cycloaddition were formed under the influence of Au(I)<sup>6e</sup> or Pd(0)<sup>6a</sup> catalysis, respectively. In some cases coupling between two bis-allenes occurred to give hetero-steroidal frameworks.<sup>6f,g</sup> It should be noted that the vast majority of these reaction is metal catalysed.

This posed the question what the favoured reaction pathway would be if we treated *e.g.* oligomethylene-linked bis-allenes with HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>,<sup>7</sup> *i.e.* under metal-free conditions. We have now performed these reactions starting from two examples of that bis-allene family. It turned out that a different cyclization type prevailed under these conditions. The outcome of these reactions will be presented and discussed below.




Chart 1 Examples of cyclization reactions of bis-allenes.



## Results and discussion

### Borane-induced ring-closure reaction of the bis-allenes

The allenes **3a** and **3b** (Scheme 2) were prepared by a variant of the Crabbé reaction as described by Ma *et al.*<sup>8</sup> Copper(I) induced treatment of 1,6-heptadiyne with *para*-formaldehyde as C<sub>1</sub>-building block and dicyclohexylamine gave the trimethylene-linked bis-allene **3a<sup>6h</sup>** (40% isolated). The analogous reaction starting from 1,7-octadiyne gave the tetramethylene-linked bis-allene **3b<sup>6f</sup>** (63% isolated, see the ESI† for details). The phenylene containing bis-allene **3c** was synthesised analogously.

We reacted compound **3a** with one molar equivalent of Piers' borane [HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>]. The rapid reaction (r.t., minutes, in CD<sub>2</sub>Cl<sub>2</sub>) gave the cyclized product **5a**. It was not isolated but characterized by spectroscopy [<sup>1</sup>H NMR: δ 5.55 (olefinic =CH-) (C<sub>6</sub>-H, see Fig. 1 for the unsystematic atom numbering scheme), δ 5.52, 5.03, (-CH=CH<sub>2</sub> substituent); <sup>11</sup>B NMR: δ 69.3; Δδ<sup>19</sup>F<sub>m,p</sub> = 13.1 ppm]. Compound **5a** was then generated *in situ* on a preparative scale and trapped by subsequent addition of pyridine. We isolated the pyridine adduct **6a** as a white crystalline solid in 73% yield. The X-ray crystal structure analysis of compound **6a** showed the newly formed cyclohexene core that has a vinyl group attached in the allylic position and it bears the -CH<sub>2</sub>B(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>(pyridine) substituent adjacent to it at the olefinic ring carbon atom C4 (Fig. 1). In solution (CD<sub>2</sub>Cl<sub>2</sub>) we monitored the typical <sup>1</sup>H/<sup>13</sup>C NMR features of the vinyl group at carbon atom C3 and the central doubly substituted cyclohexene core. The <sup>1</sup>H NMR features of the coordinated pyridine moiety show up at δ 8.67, 8.11 and 7.65 (<sup>11</sup>B: δ -0.6). Due to the chiral centre (ring carbon C3) the C<sub>6</sub>F<sub>5</sub> groups at boron are diastereotopic and give rise to a 1 : 1 set of the respective *o,p,m*-C<sub>6</sub>F<sub>5</sub> <sup>19</sup>F NMR signals.





Fig. 1 A view of the molecular structure of the borane pyridine adduct **6a**. Selected bond lengths (Å) and angles (°): B1–N31 1.613(3), B1–C5 1.651(3), C1–C2 1.322(4), C4–C6 1.337(3), B1–C5–C4 117.7(2).

We assume that the reaction started with hydroboration of the terminal =CH<sub>2</sub> unit of one allenyl group by the HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> reagent. This resulted in the formation of the functionalized borane **4** which contained a reactive allylborane unit opposite to a reactive allene unit. This situation was set up for an internal allylboration reaction of the allene, which directly opened a pathway to the observed cyclization product **5a**. Addition of the pyridine Lewis base to the strongly Lewis acidic B(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> group gave **6a** (Scheme 2).

We reacted the tetramethylene-linked bis-allene **3b** with HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> and found that the seven-membered cyclization product **5b** was formed analogously. Treatment of the *in situ* generated borane **5b** with pyridine gave the respective pyridine adduct **6b**, which we isolated crystalline in 39% yield. Compounds **5b** and **6b** were characterized by spectroscopy (see the ESI† for details); product **6b** was characterized by C, H, N elemental analysis and by an X-ray crystal structure analysis (Fig. 2). It shows the newly formed seven-membered ring in a typical cycloheptene boat-like conformation.<sup>9</sup> The -CH<sub>2</sub>B(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> moiety is attached at the sp<sup>2</sup>-carbon atom C4 (the boron atom bears a pair of C<sub>6</sub>F<sub>5</sub> groups and the coordinated



Scheme 2 HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>-induced ring closure of the bis-allenes **3**.




Fig. 2 Molecular structure of compound **6b**. Selected bond lengths (Å) and angles (°): B1–N31 1.627(5), B1–C5 1.645(6), C1–C2 1.285(8), C4–C6 1.340(6), B1–C5–C4 119.9(3).



pyridine ligand in a pseudo-tetrahedral geometry). The vinyl substituent is bonded to carbon atom C3.


The reaction of the bis-allene **3a** responded to the stoichiometry of the  $\text{HB}(\text{C}_6\text{F}_5)_2$  reagent since the primary ring-closure product **5a** contained a reactive pendant vinyl group. Therefore, the reaction of **3a** with two molar equivalents of Piers' borane gave the bis-borane product **7a**. Its  $^{11}\text{B}$  NMR spectrum showed two signals ( $\delta$  73.3 and  $\delta$  69.4) that are attributed to the pair of Lewis acidic planar-tricoordinate boron centres. Consequently, we observed two sets of  $^{19}\text{F}$  NMR resonances of the pairs of the  $\text{C}_6\text{F}_5$  substituents at the two  $\text{B}(\text{C}_6\text{F}_5)_2$  groups. The *in situ* generated bis-borane **7a** was then treated with two molar equivalents of pyridine to give the respective double pyridine adduct **8a** (isolated as a white solid in 72% yield). It was characterized by C, H, N elemental analysis, by X-ray diffraction (the structure is shown in the ESI†) and by spectroscopy. Due to the chiral centre (C3) the pair of  $\text{C}_6\text{F}_5$  groups at each  $\text{B}(\text{C}_6\text{F}_5)_2$ -(pyridine) moiety are diastereotopic and, consequently, we observed four sets of *o,m,p*- $\text{C}_6\text{F}_5$   $^{19}\text{F}$  NMR signals of compound **8a**. We observed two sets of pyridine  $^1\text{H}/^{13}\text{C}$  NMR resonances and located the single olefinic  $^1\text{H}$  NMR signal (C6–H) at  $\delta$  4.35 (t,  $J_{\text{HH}} = 3.3$  Hz, 1H).

We performed the reaction of the phenylene bridged bis-allene system **3c** in a similar way. The reaction of **3c** with one molar equiv. of  $\text{HB}(\text{C}_6\text{F}_5)_2$  was carried out in  $\text{CD}_2\text{Cl}_2$  at r.t. and the almost instantaneously *in situ* generated product was characterized by NMR spectroscopy [ $^1\text{H}$ :  $\delta$  5.48/5.13/5.08 (vinyl substituent), 5.72 (ring-CH=),  $^{11}\text{B}$ :  $\delta$  71.9,  $^{19}\text{F}$ :  $\Delta\delta^{19}\text{F}_{m,p} = 13.2$ ]. The latter heteroatom NMR signals are typical for the presence of strongly Lewis acidic tricoordinate boron with this substituent pattern. Compound **5c** was treated with pyridine and the respective pyridine/borane Lewis adduct **6c** was isolated in 37% yield after workup involving pentane extraction. The NMR spectra now show a  $^{11}\text{B}$  NMR resonance in the typical range of tetracoordinate boron ( $\delta$  –0.5) and the  $\text{C}_6\text{F}_5$  substituents at boron are diastereotopic (see the ESI† for further details).

### Subsequent FLP ring-closure reactions

The compounds **5** each contain a sterically encumbered, strongly Lewis acidic borane functionality and in its vicinity an accessible reactive vinyl group. We used this for carrying out a typical frustrated Lewis pair<sup>10</sup> reaction, namely a 1,2-borane/phosphane addition to the  $\text{C}=\text{C}$  double bond.<sup>11</sup>

We reacted the *in situ* generated cyclization product **5a** with triphenylphosphane at room temperature. The reaction was practically instantaneous under these typical conditions and we isolated the P/B addition product **9a** to the internal vinyl group as a white powder in 76% yield (Scheme 3). Single crystals of compound **9a** that were suited for the X-ray crystal structure analysis were obtained at room temperature from a solution in dichloromethane that was layered with pentane. It showed that 1,2-phosphane/borane addition to the vinyl substituent had occurred. The internal  $-\text{B}(\text{C}_6\text{F}_5)_2$  Lewis acid had been added to the  $=\text{CH}_2$  terminus of the alkene and the external  $\text{PPh}_3$  nucleophile to the  $-\text{CH}=\text{C}$  carbon atom. The resulting zwitterionic heterobicyclo[4.4.0]decene type system features



Scheme 3 FLP reaction of the boranes **5** with  $\text{PPh}_3$ .

a bridgehead  $\text{C}=\text{C}$  double bond (C4–C6). There is a borate system inside the heterocyclic six-membered ring that was formed in the FLP addition reaction. Consequently, the  $-\text{PPh}_3^+$  phosphonium substituent is found attached at the same ring at carbon atom C2. We have isolated a single diastereoisomer of **9a** from this reaction; it features the hydrogen atoms at carbon atoms C2 and C3 oriented *trans* to each other at the heterobicyclic framework (Fig. 3). In solution compound **9a** shows the NMR heteronuclear resonances at  $\delta$  –12.3 ( $^{11}\text{B}$ ) and  $\delta$  27.9 ( $^{31}\text{P}$ ). A diastereotopic pair of  $\text{C}_6\text{F}_5$  substituents is bonded at boron, giving rise to two separate sets of  $^{19}\text{F}$  NMR resonances. The  $^1\text{H}$  NMR [P]–CH– signal is found at  $\delta$  3.64 and the single olefinic  $=\text{CH}$ – signal at  $\delta$  5.17.

The seven-membered ring compounds **5b** and **5c** react analogously with  $\text{PPh}_3$ . Compound **5b** was *in situ* generated and triphenylphosphane was added. We isolated the product **9b** in 61% yield by crystallization. It was characterized by C, H, elemental analysis, by spectroscopy and by X-ray diffraction. The molecular structure is similar to that of **9a**. It also contains a *trans*-relationship of the C3–H and C2–H hydrogen atoms of the heterobicyclic ring system. The hetero-NMR signals occur at




Fig. 3 Molecular structure of the P/B FLP addition product **9a**. Selected bond lengths (Å) and angles ( $^\circ$ ): B1–C1 1.647(2), B1–C5 1.640(2), P1–C2 1.856(2), C1–C2 1.549(2), C4–C6 1.333(2), C1–B1–C5 105.7(1), B1–C1–C2 114.6(1), C1–C2–C3 112.4(1), C2–C3–C4 109.2(1), C3–C4–C5 114.9(1), C4–C5–B1 108.9(1), P1–C2–C1 109.3(1), P1–C2–C3 111.9(1), P1–C2–C3–C4 174.9(1), P1–C2–C1–B1 175.9(1).




$\delta$   $-13.3$  ( $^{11}\text{B}$ ) and  $\delta$   $30.5$  ( $^{31}\text{P}$ ). Compound **9c** was prepared analogously from **5c** (see Scheme 3). It was isolated as a white solid in 77% yield after workup and characterized by NMR spectroscopy and by an X-ray crystal structure analysis. For further details of the characterization of the compounds **9b** and **9c** including their depicted molecular structures see the ESI.†

### Subsequent cyclooligomerization reaction with allene

Internal allene allylboration<sup>12</sup> represents the important step of the ring-closure reaction sequence starting from **3** to form the products **5**. The compounds **5** themselves each contain an allylboration functionality which might show the respective reactivity towards added allene reagents. Therefore, we exposed the cyclization product **5a** to an excess of the parent allene  $\text{H}_2\text{C}=\text{C}=\text{CH}_2$ . An NMR experiment revealed a close to complete conversion to the new product **12a** within 24 h at room temperature. We carried out this reaction on a preparative scale under analogous conditions. Workup involving crystallization from pentane at  $-35$  °C (3 d) gave the crystalline product **12a**, which we isolated in 37% yield. Compound **12a** was characterized by C, H elemental analysis, by spectroscopy and by an X-ray crystal structure analysis. This showed (Fig. 4) that the endocyclic allylboration moiety of the starting material **5a** had reacted with two molar equivalents of  $\text{H}_2\text{C}=\text{C}=\text{CH}_2$  to give the functionalized decalin derivative **12a**. Apparently, compound **5a** had undergone an allylboration reaction with allene to generate the intermediate **10a**. This itself represents an “elongated” allylboration system, that subsequently took up another allene equivalent to give **11a**. The intermediate **11a** could in principle have reacted with further allene, but instead its allylboration “found” the remaining exo-methylene group at the six-membered core with which it underwent a favoured intramolecular allylboration reaction<sup>4</sup> to directly give the observed product **12a** (Scheme 4).

The X-ray crystal structure analysis of compound **12a** shows the newly formed *trans*-decalin framework that was formed by the consecutive C–C coupling between **5a** and two molar equivalents of allene. The ring carbon atom C3 bears the



Scheme 4 Formation of compound **12a** and allene by sequential allylboration reactions.

remaining vinyl substituent; carbon atoms C10 and C12 are both part of the pair of exo-methylene groups 1,3-positioned in the second ring. The  $-\text{CH}_2\text{B}(\text{C}_6\text{F}_5)_2$  group is attached at the bridgehead carbon (C4 in Fig. 4).

In solution ( $\text{CD}_2\text{Cl}_2$ ) compound **12a** shows the typical NMR features of the vinyl substituent. The pair of  $=\text{CH}_2$  exo-methylene groups shows a total of four  $^1\text{H}$  NMR resonances ( $\delta$  4.85/4.61 and  $\delta$  4.68/4.42). The  $-\text{CH}_2[\text{B}]$  substituent shows the  $^1\text{H}$  NMR signals of a pair of diastereotopic hydrogen atoms (AB system at  $\delta$  2.30/2.15;  $^{13}\text{C}$ :  $\delta$  36.7). The corresponding  $^{11}\text{B}$  NMR feature is at  $\delta$  76.4, *i.e.* in a typical range of a strongly Lewis acidic tri-coordinated boron atom in this substituent situation.<sup>13</sup> Consequently, we observed three  $^{19}\text{F}$  NMR signals of the pair of the  $\text{C}_6\text{F}_5$  substituents at boron with a large  $\Delta\delta^{19}\text{F}_{m,p} = 11.5$  ppm chemical shift separation.

### Some typical reactions of the borane **12a**

Compound **12a** is a reactive borane and it contains C=C double bond functionalities. Therefore, it should be suitable to undergo typical FLP addition to one of the olefinic units in the presence of an external phosphane nucleophile.<sup>11</sup> We, consequently, reacted the *in situ* generated borane **12a** with triphenylphosphane in dichloromethane. The reaction with  $\text{PPh}_3$  was instantaneous. The volatiles were removed and the residue was washed with pentane to give the P/B addition product **13a**, which we isolated as a white solid in 56% yield (see Scheme 5). Compound **13a** was characterized by C, H-elemental analysis, by spectroscopy and by X-ray diffraction (Fig. 5). It showed that a P/B FLP addition had taken place at the proximal  $\text{C}=\text{CH}_2$  moiety (C12=C15 in compound **12a**, see Fig. 4) by using the adjacent pendent internal borane and the external phosphane. Compound **13a** is the isomer that was formed by borane addition to the  $\text{C}=\text{CH}_2$  terminus and, consequently, phosphane addition to the  $\text{sp}^2$ -ring carbon atom. This resulted in the formation of a heterocyclic six-membered ring-system that had become 1,3-attached at the “lower” six-membered decalin ring of compound **12a**. The phosphonium  $\text{PPh}_3^+$  moiety is found attached at the new bridgehead atom C12 (Fig. 5).





Fig. 4 Molecular structure of compound **12a**. Selected bond lengths (Å) and angles (°): B1–C5 1.554(3), C1–C2 1.313(3), C4–C5 1.557(3), C4–C6 1.566(3), C10–C14 1.324(3), C12–C15 1.324(3), B1–C5–C4 123.2(2), C5–C4–C6 109.0(2).





Scheme 5 Reaction of compound **12a** with PPh<sub>3</sub> and oxidative replacement of the boryl group.




Fig. 5 A view of the P/B FLP alkene addition product **13a**. Selected bond lengths (Å) and angles (°): B1–C5 1.640(6), B1–C15 1.667(6), P1–C12 1.858(4), C1–C2 1.302(6), C4–C6 1.565(5), C10–C14 1.328(6), C12–C15 1.555(5), C5–B1–C15 110.3(3), C12–C15–B1 114.3(3), P1–C12–C15–B1 –129.8(3).

Compound **13a** shows a typical borate <sup>11</sup>B NMR resonance at  $\delta$  –15.1 in solution (CD<sub>2</sub>Cl<sub>2</sub>, 273 K) and a phosphonium <sup>31</sup>P NMR signal at  $\delta$  31.1. It shows the <sup>19</sup>F NMR features of a pair of diastereotopic C<sub>6</sub>F<sub>5</sub> groups at the boron atom (for further details of the NMR characterization of compound **13a** see the ESI†).

Compound **13a** is the P/B FLP addition product that has been formed under kinetic control. When we stored the CD<sub>2</sub>Cl<sub>2</sub> solution of compound **13a** for 7 days at room temperature the resulting NMR spectra showed the formation of an equilibrium mixture of **13a** (ca. 20 mol%), the starting material **12a** (plus PPh<sub>3</sub>, ca. 8 mol%) and the new compound **14a** (ca. 65 mol%) (plus some minor contaminants). The major product **14a**, apparently formed under thermodynamic control, was prepared similarly on a preparative scale (24 h, r.t., CH<sub>2</sub>Cl<sub>2</sub> layered with pentane) and crystallized from the mixture. Crystalline compound **14a** was isolated in 60% yield and the product was characterized by C, H-elemental analysis, by spectroscopy and by X-ray diffraction.

The X-ray crystal structure analysis of **14a** shows the presence of a five-membered boratacycle that is 1,3-annulated to the

“lower” six-membered ring of the *trans*-decalin framework. It was apparently formed by a 1,2-P/B FLP addition to the C12=C15 carbon–carbon double bond of the starting material **12a**, similar as we had seen it in the formation of its isomer **13a**, only that in this case PPh<sub>3</sub> addition had taken place at the =CH<sub>2</sub> terminus of the exo-methylene group concurrent with borane addition to its adjacent doubly substituted sp<sup>2</sup>-carbon atom. The structure of the resulting P/B zwitterion **14a** is depicted in Fig. 6.

The <sup>1</sup>H NMR spectrum of compound **14a** (in CD<sub>2</sub>Cl<sub>2</sub>, at 299 K) shows the P-coupled system of the exocyclic –CH<sub>2</sub>–[P] moiety at  $\delta$  4.18/2.70 and the resonances of the endocyclic –CH<sub>2</sub>–[B] group at  $\delta$  1.32/0.35. The heteroatom NMR signals occur at  $\delta$  –6.8 (<sup>11</sup>B) and 19.9 (<sup>31</sup>P), respectively and we observed two sets of <sup>19</sup>F NMR signals of the pair of diastereotopic C<sub>6</sub>F<sub>5</sub> groups at boron (for further details see the ESI†).

We eventually converted the borane-induced multi-component cyclization product **12a** to a boron-free derivative.<sup>14</sup> This was carried out in the usual way of oxidative deborylation as it is done in conventional hydroboration chemistry.<sup>15</sup> Treatment of the strongly electrophilic –B(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> borane **12a** with NaOH/H<sub>2</sub>O<sub>2</sub> gave the alcohol **15a** that we isolated as a white solid in 42% yield after workup (see the ESI† for its characterization by NMR spectroscopy).

We briefly investigated the reaction of the bis-allylic ether **16**<sup>cc,hj</sup> with HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>. The reaction was carried out in CD<sub>2</sub>Cl<sub>2</sub> solution at r.t. The products of the reaction were not isolated but directly identified *in situ* generated from the solution. We subsequently added a total of three molar equivalents of HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> to eventually achieve a complete conversion of compound **16** with a clean product formation. The NMR analysis (for details see the ESI†) revealed that the by far predominant reaction was ether cleavage. This gave the (C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>B–O–CH<sub>2</sub>–CH=C=CH<sub>2</sub> cleavage product **17** (see Scheme 6) and butadiene (**18**) as primary products. The latter was then subsequently converted by added HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> to the bis-hydroboration product **19**. The boryl ether **17** also was not stable under the reaction conditions, probably due to subsequent ether cleavage with additional HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> (see the ESI† for details). We also investigated briefly the reaction of the





Fig. 6 A projection of the molecular structure of compound **14a**. Selected bond lengths (Å) and angles (°): B1–C5 1.657(4), B1–C15 1.695(3), P1–C15 1.823(2), C1–C2 1.301(13), C4–C6 1.556(3), C10–C14 1.325(4), C12–C15 1.540(3), C5–B1–C12 99.9(2), C12–C15–B1 112.1(2), B1–C12–C15–P1 179.1(2).





Scheme 6 Reaction of the bis-allenic ether **16** with  $\text{HB}(\text{C}_6\text{F}_5)_2$ .

respective bis-allenic *N*-tosyl amine with  $\text{HB}(\text{C}_6\text{F}_5)_2$ , but that gave a complicated mixture of as yet unidentified products.

## Conclusions

With this study we have found a new variant of our borane induced carbon-carbon coupling reactions between allene building blocks. In this case the reaction starts as it is commonly observed in our systems by 1,2-[B]-H addition<sup>16</sup> to a terminal allene  $=\text{CH}_2$  group by the strongly electrophilic  $\text{HB}(\text{C}_6\text{F}_5)_2$  hydroboration reagent to probably generate a reactive allylborane intermediate *in situ*, which is set for undergoing rapid intramolecular ring-closure with the pendant second allenyl moiety to generate the products **5a** to **5c**, respectively. These are then obviously protected by their special geometry from undergoing further intermolecular allylborane coupling under the applied reaction conditions, so that the reaction stopped at the functionalized six- or seven-membered ring products. The compounds **5** are, however, in principle still active allylborane reagents. This we could show by the rapid reaction of the example **5a** with the parent allene  $\text{H}_2\text{C}=\text{C}=\text{CH}_2$ . Two equivalents of allene were consumed in a sequence of consecutive intramolecular allylborane reactions, followed by a final intramolecular allylborane ring-closure reaction to give the four-component coupling product **12a**. This in turn was oxidatively converted to the boron-free product **15a**. These metal-free reactions are markedly different from the common metal catalysed bis-allenic cyclization reactions reported in the literature (see Chart 1 and the respective references). We will see how the products of our metal free cyclization reactions and their follow-up products (and related systems) might become easily available useful reagents for further external C-C coupling reactions using either of the newly generated functionalities.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- (a) A. S. K. Hashmi, *Angew. Chem., Int. Ed.*, 2000, **39**, 3590; (b) *Modern Allene Chemistry*, ed. N. Krause, and A. S. K. Hashmi, Wiley-VCH, Weinheim, Germany, 2004; (c) S. Ma, *Chem. Rev.*,

2005, **105**, 2829; (d) N. Krause and C. Winter, *Chem. Rev.*, 2011, **111**, 1994; (e) S. Yu and S. Ma, *Angew. Chem., Int. Ed.*, 2012, **51**, 3074; (f) W. Yang and A. S. K. Hashmi, *Chem. Soc. Rev.*, 2014, **43**, 2941.

- C. Gropp, N. Trapp and F. Diederich, *Angew. Chem., Int. Ed.*, 2016, **55**, 14444.
- (a) D. J. Parks, R. E. H. Spence and W. E. Piers, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 809; (b) D. J. Parks, W. E. Piers and G. P. A. Yap, *Organometallics*, 1998, **17**, 5492; (c) M. Hoshi, K. Shirakawa and M. Okimoto, *Tetrahedron Lett.*, 2007, **48**, 8475; (d) A. Schnurr, K. Samigullin, J. M. Breunig, M. Bolte, H.-W. Lerner and M. Wagner, *Organometallics*, 2011, **30**, 2838; (e) J. Zhang, S. Park and S. Chang, *Angew. Chem., Int. Ed.*, 2017, **56**, 13757.
- (a) X. Tao, G. Kehr, C. G. Daniliuc and G. Erker, *Angew. Chem., Int. Ed.*, 2017, **56**, 1376; (b) See also: X. Tao, C. Wölke, C. G. Daniliuc, G. Kehr and G. Erker, *Chem. Sci.*, 2019, **10**, 2478.
- Arylallenes react differently, see: X. Tao, C. G. Daniliuc, D. Dittrich, G. Kehr and G. Erker, *Angew. Chem., Int. Ed.*, 2018, **57**, 13922.
- (a) X. Jiang, X. Cheng and S. Ma, *Angew. Chem., Int. Ed.*, 2006, **45**, 8009; (b) S.-K. Kang, T.-G. Baik, A. N. Kulak, Y.-H. Ha, Y. Lim and J. Park, *J. Am. Chem. Soc.*, 2000, **122**, 11529; (c) Y.-T. Hong, S.-K. Yoon, S.-K. Kang and C.-M. Yu, *Eur. J. Org. Chem.*, 2004, 4628; (d) J. Cheng, X. Jiang and S. Ma, *Org. Lett.*, 2011, **13**, 5200; (e) S. M. Kim, J. H. Park, Y. K. Kang and Y. K. Chung, *Angew. Chem., Int. Ed.*, 2009, **48**, 4532; (f) S. Ma, P. Lu, L. Lu, H. Hou, J. Wei, Q. He, Z. Gu, X. Jiang and X. Jin, *Angew. Chem., Int. Ed.*, 2005, **44**, 5275; (g) S. Ma and L. Lu, *Chem.-Asian J.*, 2007, **2**, 199; (h) X. Lian and S. Ma, *Chem.-Eur. J.*, 2010, **16**, 7960; (i) V. A. D'yakonov, G. N. Kadikova, L. M. Khalilov and U. M. Dzhemilev, *Russ. J. Org. Chem.*, 2013, **49**, 1139; (j) S.-K. Kang, Y.-H. Ha, D.-H. Kim, Y. Lim and J. Jung, *Chem. Commun.*, 2001, 1306.
- For examples of thermally induced reactions of bis-allenes, see: (a) L. Skattebol and S. Solomon, *J. Am. Chem. Soc.*, 1965, **87**, 4506; (b) W. R. Roth, M. Heiber and G. Erker, *Angew. Chem., Int. Ed. Engl.*, 1973, **12**, 504; (c) S. Sakai, *J. Phys. Chem. A*, 2006, **110**, 9443.
- (a) J. Kuang and S. Ma, *J. Org. Chem.*, 2009, **74**, 1763; for the original Crabbé reaction see *e.g.*; (b) P. Crabbé, H. Fillion, D. André and J.-L. Luche, *J. Chem. Soc., Chem. Commun.*, 1979, 859; see also; (c) S. Kitagaki, M. Komitsu and C. Mukai, *Synlett*, 2011, 1129.
- Basic Organic Stereochemistry*, ed. E. L. Eliel, S. H. Wilen and M. P. Doyle, John Wiley & Sons, Inc., New York, 2001.
- (a) D. W. Stephan and G. Erker, *Angew. Chem., Int. Ed.*, 2010, **49**, 46; (b) D. W. Stephan and G. Erker, *Angew. Chem., Int. Ed.*, 2015, **54**, 6400.
- (a) J. S. J. McCahill, G. C. Welch and D. W. Stephan, *Angew. Chem., Int. Ed.*, 2007, **46**, 4968; (b) J. B. Sortais, T. Voss, G. Kehr, R. Fröhlich and G. Erker, *Chem. Commun.*, 2009, 7417; (c) T. Voss, J.-B. Sortais, R. Fröhlich, G. Kehr and G. Erker, *Organometallics*, 2011, **30**, 584; (d) X. X. Zhao and D. W. Stephan, *J. Am. Chem. Soc.*, 2011, **133**, 12448; (e)



- G. Ménard, L. Tran, J. S. J. McCahill, A. J. Lough and D. W. Stephan, *Organometallics*, 2013, **32**, 6759.
- 12 (a) B. M. Mikhailov and Y. N. Bubnov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1964, **13**, 1874; (b) *Organoboron Compounds in Organic Synthesis*, ed. B. M. Mikhailov and Y. N. Bubnov, Harwood, Chur, 1984; (c) Y. N. Bubnov, *Pure Appl. Chem.*, 1987, **59**, 895; (d) R. W. Hoffmann, *Pure Appl. Chem.*, 1988, **60**, 123; (e) Y. Yamamoto and N. Asao, *Chem. Rev.*, 1993, **93**, 2207; (f) S. Y. Erdyakov, M. E. Gurskii, A. V. Ignatenko and Y. N. Bubnov, *Mendeleev Commun.*, 2004, **14**, 242; (g) M. E. Gurskii, S. Y. Erdyakov, T. V. Potapova and Y. N. Bubnov, *Russ. Chem. Bull.*, 2008, **57**, 802.
- 13 For a comparison, the  $^{11}\text{B}$  NMR signal of  $\text{B}(\text{C}_6\text{F}_5)_3$  is  $\delta$  59.5. (a) G. Massey and A. J. Park, *J. Organomet. Chem.*, 1964, **2**, 245; (b) A. G. Massey and A. J. Park, *J. Organomet. Chem.*, 1966, **5**, 218.
- 14 We had previously shown that alkenyl- $\text{B}(\text{C}_6\text{F}_5)_2$  derivatives undergo typical borane reactions, e.g. Pd-catalyzed cross-coupling with iodobenzene: (a) C. Chen, G. Kehr, R. Fröhlich and G. Erker, *J. Am. Chem. Soc.*, 2010, **132**, 13594; (b) C. Chen, T. Voss, R. Fröhlich, G. Kehr and G. Erker, *Org. Lett.*, 2011, **13**, 62.
- 15 (a) H. C. Brown and B. C. Rao, *J. Am. Chem. Soc.*, 1956, **78**, 5694; (b) H. C. Brown and B. C. Rao, *J. Org. Chem.*, 1957, **22**, 1136; (c) H. C. Brown and G. Zweifel, *J. Am. Chem. Soc.*, 1959, **81**, 247.
- 16 For an example of a rare alternative 1,1-hydroboration reaction, see: A. Ueno, J. Yu, X. Tao, C. G. Daniliuc, G. Kehr and G. Erker, *Organometallics*, 2018, **37**, 2665.

