Chemical Science

EDGE ARTICLE



View Article Online View Journal | View Issue

Check for updates

Cite this: Chem. Sci., 2020, 11, 3028

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 2nd December 2019 Accepted 15th February 2020

DOI: 10.1039/c9sc06102a

rsc.li/chemical-science

Introduction

The recent surge of interest in molecular carbon nanoscience is driven by the pursuit of atomic precision in larger nanostructures (e.g. graphene nanoribbons and carbon nanotubes)¹ and by the extraordinary properties displayed by molecular nanocarbons themselves.² Large polycyclic aromatic hydrocarbons (PAHs) represent a transition in the continuum between molecules and nanostructures. These structurally diverse compounds exhibit promise for a range of applications due to their exceptional photophysical and electronic properties, and their ability to organize into functional assemblies via π stacking.3 Since most carbon nanostructures are comprised of fused rings, PAHs are their natural synthetic building blocks. For example, the polymerization of PAH monomers may offer the best approach for growing atomically precise graphene nanoribbons⁴ and the synthesis of carbon nanotubes and fullerenes from belt- and bowl-shaped PAH precursors should eventually be feasible.5 PAHs are also attractive as components

Site-selective [2 + 2 + n] cycloadditions for rapid, scalable access to alkynylated polycyclic aromatic hydrocarbons[†]

Gavin R. Kiel, 🝺 Harrison M. Bergman ២ and T. Don Tilley D *

Polycyclic aromatic hydrocarbons (PAHs) are attractive synthetic building blocks for more complex conjugated nanocarbons, but their use for this purpose requires appreciable quantities of a PAH with reactive functional groups. Despite tremendous recent advances, most synthetic methods cannot satisfy these demands. Here we present a general and scalable [2 + 2 + n] (n = 1 or 2) cycloaddition strategy to access PAHs that are decorated with synthetically versatile alkynyl groups and its application to seven structurally diverse PAH ring systems (thirteen new alkynylated PAHs in total). The critical discovery is the site-selectivity of an Ir-catalyzed [2 + 2 + 2] cycloaddition, which preferentially cyclizes tethered diyne units with preservation of other (peripheral) alkynyl groups. The potential for generalization of the siteselectivity to other [2 + 2 + n] reactions is demonstrated by identification of a Cp₂Zr-mediated [2 + 2 + n]1]/metallacycle transfer sequence for synthesis of an alkynylated, selenophene-annulated PAH. The new PAHs are excellent synthons for macrocyclic conjugated nanocarbons. As a proof of concept, four were subjected to alkyne metathesis catalysis to afford large, PAH-containing arylene ethylene macrocycles, which possess a range of cavity sizes reaching well into the nanometer regime. Notably, these highyielding macrocyclizations establish that synthetically convenient pentynyl groups can be effective for metathesis since the 4-octyne byproduct is sequestered by 5 Å MS. Most importantly, this work is a demonstration of how site-selective reactions can be harnessed to rapidly build up structural complexity in a practical, scalable fashion.

> of many other types of functional materials, including polymers,⁶ macrocycles,⁷ and metal and covalent organic frameworks.⁸

> The synthesis of PAHs is challenging since it requires regioselective fusions of many rings. Recently, much progress has been made to address this challenge,^{1,2a,9} and metalmediated cycloadditions have proven particularly valuable due to their efficacy in the formation of aromatic rings.¹⁰ This research group is focused on development of a unified approach to PAH synthesis (Scheme 1a), based on divergent [2 + 2 + n] cycloadditions of biaryl-tethered diynes or dinitriles (**A**) to give highly functionalized PAHs (**B**).¹¹ Importantly, since only one conformational degree of freedom is frozen out with each cyclization, the minimal loss of entropy contributes to excellent yields^{11a,b} and enables the use of more challenging dinitrile^{11c} substrates.

> For a PAH to be a useful synthon, it must be available in appreciable quantities *and* be suitably functionalized, but this set of requirements has proven difficult to achieve. Given their high carbon content and synthetic versatility, alkynyl substituents are attractive for the elaboration of PAHs into more complex functional materials.¹² Such substituents are especially useful for macrocyclization reactions to give arylene ethynylene macrocycles (AEMs),¹³ cycloarylenes,¹⁴ and other exotic

Department of Chemistry, University of California, Berkeley, Berkeley, California, 94720, USA. E-mail: tdtilley@berkeley.edu

[†] Electronic supplementary information (ESI) available. See DOI: 10.1039/c9sc06102a



macrocycles.15 These and related macrocyclic nanocarbons5c,7,16 represent a broad structural class of tremendous interest for their internal cavities and unique spatial arrangements of carbon-rich building blocks.

Alkynyl groups are usually installed onto PAHs from the corresponding halide (via Sonogashira coupling) or carbonyl (via nucleophilic attack), but these precursors are limited in number given the established regiochemical preferences for halogenation or oxidation. Other widely employed methods to access carbonyl-containing precursors involve Friedel-Craftstype cyclizations or Diels-Alder/oxidation sequences, but these also offer limited structural diversity. Perhaps most importantly, these routes add synthetic steps and usually involve harsh conditions. Thus, a more direct entry to alkynylated PAHs that bypasses traditional intermediates would be highly desirable. Relevant previous work by the Matano¹⁷ and Tanaka^{15b} groups (Scheme 1b) demonstrated use of [2 + 2 + n] reactions to produce alkynylated synthetic intermediates C and D en route to oligomeric (non-PAH) products. Importantly, in both cases, one of the alkynes on the starting material was protected by a bulky silyl group, which minimizes the many possible deleterious cycloaddition side reactions.18 The efficiency described above for cycloadditions of biaryl-tethered diynes (A) suggested that these transformations might be site-selective19 and select for tethered divne units (i.e. with preservation of peripheral, nontethered alkynyl groups), as shown in Scheme 1c. This would alleviate the need to protect one of the alkynes, which would greatly simplify the syntheses of oligo(alkynyl)arylene precursors (1) and provide direct access to valuable alkynylated PAHs (2).

Here we describe development of the strategy of Scheme 1c, which was found to be possible with both Ir-catalyzed [2 + 2 + 2]and Cp_2Zr -mediated [2 + 2 + 1] cycloadditions. This previously unrecognized site-selectivity gives rapid access to alkynylated PAHs, which is demonstrated with thirteen examples, representing seven different ring systems (Schemes 2-4). The [2 +2+1 variant also offers the possibility of a divergent annulation with heterocyclic rings. Due to the streamlined precursor syntheses and generally high yields, multigram quantities of alkynylated PAHs are easily accessed, which is critical for their intended use as synthetic intermediates. As a proof of concept, four of the new PAHs were exploited as synthons for macrocyclic nanocarbons via alkyne metathesis reactions of their alkynyl groups (Scheme 5). Three of these structurally complex macrocycles are derived from previously unknown PAH ring systems. More importantly, they possess a range of internal cavity sizes reaching well into the nanometer regime.

Results and discussion

The new strategy was initially envisioned as a solution to difficulties encountered in the synthesis of alkynylated triphenylenes such as 2a and 2b (Scheme 2) using classical ring fusion techniques (e.g. the Scholl reaction²⁰). Notably, there was no literature precedent for, or suitable synthetic precursors to, the deceptively simple 2,7-dialkynyl-3,6-dihalo triphenylene substitution pattern. Since the starting dialkynyl dihalides 4a and 4b are available in only two steps and can be easily homocoupled to yield tetra(alkynyl)biphenylene precursors 1a and 1b,^{11a} this strategy would provide streamlined access to triphenylenes 2a and 2b if the subsequent site-selective [2 + 2 +2] reaction was successful. The most important step in the synthesis of precursors 1a and 1b is the highly selective in situ desymmetrization via Li/Br exchange that precedes the Cumediated oxidative homocoupling. This desymmetrization also streamlines the synthesis of all other oligo(alkynyl)arylene substrates in this manuscript (e.g. 1c-1l in Scheme 3; see ESI for more details[†]).

The [2 + 2 + 2] pre-catalyst Ir(COD)(dppe)Cl, which can be isolated²¹ or generated in situ^{22a,b} from commercially available materials, proved to be highly effective for synthesis of model pentynylated triphenylene 2a. This compound was isolated in a remarkable 89% yield under optimized conditions. The preservation of the pentynyl groups in 2a was verified by ¹H and ¹³C NMR spectroscopies (one propargylic methylene triplet and



Scheme 2 Model system for site-selective [2 + 2 + 2] reaction.



*¹H NMR yields; **Synthesized by protodebromination of **2f** (for **2g**) or **2h** (for **2i**)

Scheme 3 Generalization of the site-selective [2 + 2 + 2] reaction to more complex PAHs.

two quaternary alkynyl resonances, respectively) and highresolution mass spectrometry. The choice of Ir(COD)(dppe)Clas pre-catalyst was informed both by its exceptional efficiency in [2+2+2] reactions involving biphenyl-tethered diynes^{11b} and by Takeuchi's demonstration of its ability to furnish alkynylated benzene products^{22a,b} (in [2+2+2] reactions involving a 1,3diyne reactant).

Since the peripheral pentynyl groups (shown in red) in the starting material (1a) and product (2a) are virtually identical, it

was anticipated that competing intermolecular cycloadditions involving these groups would necessitate extensive screening and optimization efforts. In contrast, the yield of this transformation was fairly insensitive (82-93%) to wide ranges of values for several critical parameters, including concentration (10-300 mM), monoyne equivalents (1.0-5.7), catalyst loading (1–10%), and reaction time (1–10 h), as discussed further in the ESI (Tables S1-S3[†]). Under the optimized conditions, there was a notable decrease in yield (61%) for triphenylene 2b, which has less sterically demanding propynyl groups. The major side product was dimeric 5 (8-18% by ¹H NMR, depending on concentration), resulting from a cycloaddition between the product (2b) and starting material (1b). Dilution from to 10 mM (from 100 mM) did not improve the yield. Notably, the yield of 2b increased to 84% with a larger excess of monoyne (10 equiv.), and side product 5 was not observed. Since bulkier pentynyl groups provided higher yields in the model system, they were used for the remaining examples below.

The site-selective [2 + 2 + 2] reaction proved to be a general means to access alkynylated PAHs, as demonstrated by the isolation of 2c-2l (Scheme 3). Precursors 1c-1l were synthesized in a modular fashion via standard C-C bond-forming reactions (see Schemes S1-S3 for details[†]), all of which were streamlined by Li-mediated desymmetrization reactions as mentioned above. From the same benzene, naphthalene, and phenanthrene building blocks, a range of different shapes, sizes, and alkynyl group substitution patterns were obtained via onefold, twofold, or threefold cycloadditions. Notably, the decrease in yields in the absence of the sterically demanding Br groups ortho to the alkynes (for 2d and 2j relative to 2c and 2k) was relatively small. This suggests that ortho substituents play a smaller role than the substituent on the alkynyl group in shutting down undesirable intermolecular side reactions. Also, the size of the PAH building blocks does not appear to play a large role. This can be seen by the comparable yields for 2a (89%), 2e (80%), and 2f (85%), which were derived from cycloadditions of biphenylene (1a), binaphthalene (1e), and biphenanthrene (1f) precursors, respectively. Interestingly, the generally good-to-excellent yields do not correlate with the number of required cycloadditions. For example, a twofold cycloaddition produced 2j in 70% yield, whereas a threefold cycloaddition produced 2l in 74% yield. The substitution patterns of the new PAHs would be difficult or impossible to access using previous synthetic methodology, even for the deceptively simple triphenylene systems 2a-d. Represented among these examples are the first known tribenzo [a,h,o] pentaphene (2f-g), tribenzo[a,h,o]phenanthro[2,3-q]hexaphene (2j**k**), and pentabenzo [a,c,j,q,s] heptaphene (21) ring systems. Perhaps most importantly, since high dilution is not necessary (divne concentration = 100 mM), the new strategy is highly scalable. For example, 2e, 2f, and 2h were isolated on 2.0, 3.4, and 1.3 gram scales, respectively.

This strategy was subsequently generalized to include a formal [2 + 2 + 1] reaction, as shown in Scheme 4. Heating an equimolar mixture of **1m** and Cp₂Zr(pyr)(Me₃SiC \equiv CSiMe₃)²³ (an isolable source of "Cp₂Zr") in hexanes, isolation of the crude zirconacyclopentadiene intermediate (**2m**) *via* filtration, and



Scheme 4 Extension of the strategy to include a Cp_2Zr -mediated formal [2 + 2 + 1] reaction.

treatment of a CH₂Cl₂ solution of 2m with SeCl₂(bipy)^{11b} afforded selenophene-annulated dibenzo[*a*,*o*]pentaphene **2n** in 41% isolated yield. The silvl groups on 1m were chosen since they often lead to reversible [2 + 2 + 1] cycloadditions involving zirconocene,^{14,24} and 2m was expected to be the thermodynamic product (entropy should favor monomeric 2m over the oligomeric products that would result from intermolecular cycloadditions¹⁴). Indeed, the site-selectivity of this reaction appears to occur under thermodynamic control. Thus, when 1m was treated with 1.0 equiv. of $Cp_2Zr(pyr)(Me_3SiC \equiv CSiMe_3)$ (in benzene- d_6 due to solubility concerns), 2m was observed in only 20% yield after 10 min at 22 °C, along with starting 1m (30%), a complex mixture of unidentified products (likely oligomers), and no remaining $Cp_2Zr(pyr)(Me_3SiC \equiv CSiMe_3)$. Temperatures up to 90 °C did not lead to notable increases in 2m, but after 18 h at 105 °C the yield of 2m increased to 45%.²⁵ Importantly, 2m is expected to be a versatile synthon for the installation of a range of other heterocycles using well-developed zirconacycle transfer chemistry.26

The angular disposition of the alkynyl groups in PAHs 2cd and 2f-I makes them attractive for macrocyclization *via* alkyne metathesis, which often provides AEMs in a high yielding, scalable fashion.²⁷ PAHs are intriguing building blocks for more complex AEMs, as has been demonstrated on a number of occasions.²⁸ However, for all but the smallest PAHs (phenanthrene and carbazole), such AEMs have been produced by routes that are difficult to scale or generalize, probably due to difficulties associated with accessing appropriately functionalized PAH precursors. The site-selective [2 + 2 + 2] cycloaddition presented above directly addresses those difficulties.

For the macrocyclization reactions of Scheme 5, the Fürstnertype metathesis catalyst²⁹ "EtC=Mo(OSiPh₃)₃" (Mo-1) was employed. In analogy to the approach taken by Moore,³⁰ we found that Mo-1 can be conveniently generated *in situ* by treatment of easily-prepared EtC=Mo(OC(CH₃)(CF₃)₂)₃(DME) (Mo-2)³¹ with 6 equiv. of Ph₃SiOH (see ESI for details on this ligand exchange†). Triphenylene ethynylene macrocycle **3-Trim**_a was isolated in excellent yield (88%) after treatment of **2d** with a 5% loading of Mo-1 in the presence of 5 Å molecular sieves²⁹ (16 h at 80 °C in



Scheme 5 New PAHs as synthons for large arylene ethynylene macrocycles.

toluene). The identity of **3-Trim**_a was unambiguously determined by ¹H and ¹³C NMR spectroscopy (the former showed the absence of propargylic methylene resonances and the latter showed only one quaternary alkynyl resonance) and MALDI-TOF. As is often the case for alkyne metathesis macrocyclizations,^{27c,d} minimal purification was required (elution of the reaction mixture through a plug of silica gel followed by precipitation with MeOH), and high dilution was not necessary. Attempts at the macrocyclization of the analogous, Br-substituted **2c** yielded a complex mixture that contained only small amounts of macrocycle, even at higher (20%) catalyst loading.

Remarkably, the synthesis of **3-Trim**_a demonstrates that 5 Å MS are effective for sequestration of the byproduct 4-octyne, which is imperative for the high yield. Fürstner's 2010 discovery²⁹ that alkyne metathesis reactions can be driven by sequestration of (smaller) 2-butyne has played a critical role in the continued rise of this synthetic methodology.³² As a result, the use of propynyl groups has become ubiquitous since the easily-sequestered 2-butyne byproduct is produced, but the installation of propynyl groups requires either gaseous propyne (for Sonogashira coupling) or expensive propynyl magnesium bromide solution (for Kumada coupling). The high-yielding synthesis of **3-Trim**_a (and the other macrocycles presented below) demonstrates that pentynyl groups may provide a cheaper, more convenient alternative to propynyl groups.

Since the presence of bulky Br substituents appears to inhibit macrocyclization with Mo-1 as catalyst, attention was focused on "Br-free" 2g, 2i, 2j, and 2l. Due to the larger sizes of these PAHs, the solvent was switched to o-dichlorobenzene (o-DCB) to limit the possibility of kinetic traps via precipitation of oligomeric or polymeric intermediates. Subjection of 2j, 2g, and 2l to the slightly modified metathesis conditions afforded the elliptical macrocycle 3-Dim and large, triangular macrocycles 3-Trim_b and 3-Trim_c in good to excellent isolated yields. While the yield of the largest of these macrocycles (3-Trim_c) improved on dilution (from 43% at 25 mM to 74% at 9 mM), this is still quite concentrated for a macrocyclization reaction. Since no welldefined side products were observed in any case, purification was again streamlined. Unfortunately, subjection of 2i to the conditions of Scheme 5 produced an intractable mixture composed primarily (~95% by mass) of insoluble material that could not be characterized. It is important to note that, except for the modest dilution in the case of 3-Trim_c, we have not attempted to optimize any parameter in these reactions.

In general, the new PAHs (2a-n) and macrocycles (3) are highly soluble in chlorinated solvents (>100 mg mL⁻¹) and insoluble in saturated hydrocarbons (e.g. hexanes) and more polar solvents (e.g. acetone and EtOH). The exception is 3-Dim, which is only moderately soluble in chloroform (15-20 mg mL^{-1}). The solution behavior of the new AEMs (3) and their PAH precursors (2d, 2j, 2g, and 2l) were investigated by variable concentration ¹H NMR spectroscopy. The chemical shifts in chloroform-d of all except 3-Dim and 3-Trim_b are independent of concentration (see Fig. S74-S81[†]), which suggests that the high solubility results from limited aggregation via π -stacking. This is likely a result of the alkyl groups in the bay positions, which induce non-planarity. In contrast, despite this alkyl group substitution, the ¹H NMR spectra of 3-Dim and 3-Trim_b in chloroform-d solution exhibit significantly broadened aromatic resonances with chemical shifts that depend upon concentration, which suggests aggregation via π -stacking.

The UV/vis absorption and emission spectra of AEMs (3) and their monomeric precursors (2) display the typical features of large, highly benzenoid PAH systems,^{3a} including high molar absorptivities (up to $5.0 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$, for **3-Trim**_c), relatively large optical HOMO–LUMO gaps (between 2.8 to 3.3 eV), and fluorescence (see Table S4 and Fig. S82–S86†). Despite the large increases in the sizes of the π -systems as a result of macrocyclization of 2 to form 3, the expected red shifts in absorption maxima range from small (21 nm for **3-Dim**) to modest (54 nm for **3-Trim**_a). More strikingly, very small decreases in optical HOMO–LUMO gap ($\leq 0.15 \text{ eV}$) and emission maxima ($\leq 12 \text{ nm}$) are observed. Thus, to a large extent, the new PAH-containing AEMs retain the properties of their PAH precursors.

Conclusions

The results presented above demonstrate how the newly discovered site-selectivity of an Ir-catalyzed [2 + 2 + 2] cycloaddition enables streamlined construction of complex nanocarbons. Here, the [2 + 2 + 2] reaction was used in concert with an orthogonal alkyne metathesis macrocyclization to provide scalable, two-step access to large, PAH-containing AEMs. Several opportunities are available for the introduction of structural and functional diversity, which might be systematically exploited to perturb self-assembly, photophysical, or electronic properties of a specific AEM, or to access entirely new AEMs. It is important to note, however, that the targeted AEMs represent only a proof of concept; the intermediate alkynylated PAHs **2a–n** are valuable synthons for a range of other conjugated nanocarbons, including expanded helicenes, cycloarenes, and other macrocyclic nanocarbons. Such applications are the focus of continuing efforts in our laboratory.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was funded by the National Science Foundation under Grant No. CHE-1708210. We thank UC Berkeley's NMR facility for resources provided and the staff for their assistance. Instruments in the NMR facility are supported in part by NIH S10OD024998. We thank Stephen von Kugelgen and Felix R. Fischer for initial samples of **Mo-2**, and Stephen von Kugelgen for helpful discussions.

Notes and references

- 1 (*a*) K. Müllen, Graphene as a Target for Polymer Synthesis, in *Hierarchical Macromolecular Structures: 60 Years after the Staudinger Nobel Prize II*, ed. V. Percec, Advances in Polymer Science; Springer International Publishing, 2013, pp. 61–92; (*b*) Y. Segawa, H. Ito and K. Itami, Structurally Uniform and Atomically Precise Carbon Nanostructures, *Nat. Rev. Mater.*, 2016, 1(1), 15002.
- 2 (a) Polycyclic Arenes and Heteroarenes: Synthesis, Properties, and Applications, ed. Q. Miao, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2015; (b) Carbon-Rich Compounds: From Molecules to Materials, ed. M. M. Haley and R. R. Tykwinski, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2006; (c) J. Wu, W. Pisula and K. Müllen, Graphenes as Potential Material for Electronics, Chem. Rev., 2007, 107(3), 718-747.
- 3 (a) R. Rieger and K. Müllen, Forever Young: Polycyclic Aromatic Hydrocarbons as Model Cases for Structural and Optical Studies, *J. Phys. Org. Chem.*, 2010, **23**(4), 315–325; (b) Z. Sun, Q. Ye, C. Chi and J. Wu, Low Band Gap Polycyclic Hydrocarbons: From Closed-Shell near Infrared Dyes and Semiconductors to Open-Shell Radicals, *Chem. Soc. Rev.*, 2012, **41**, 7857–7889; (c) Z. Chen, A. Lohr, C. R. Saha-Möller and F. Würthner, Self-Assembled π -Stacks of Functional Dyes in Solution: Structural and Thermodynamic Features, *Chem. Soc. Rev.*, 2009, **38**(2), 564–584.
- 4 (a) P. Ruffieux, S. Wang, B. Yang, C. Sánchez-Sánchez, J. Liu,
 T. Dienel, L. Talirz, P. Shinde, C. A. Pignedoli, D. Passerone,
 T. Dumslaff, X. Feng, K. Müllen and R. Fasel, On-Surface

Synthesis of Graphene Nanoribbons with Zigzag Edge Topology, *Nature*, 2016, **531**(7595), 489–492; (*b*) Y. Fogel, L. Zhi, A. Rouhanipour, D. Andrienko, H. J. Räder and K. Müllen, Graphitic Nanoribbons with Dibenzo[e,l]Pyrene Repeat Units: Synthesis and Self-Assembly, *Macromolecules*, 2009, **42**(18), 6878–6884; (*c*) D. J. Rizzo, G. Veber, T. Cao, C. Bronner, T. Chen, F. Zhao, H. Rodriguez, S. G. Louie, M. F. Crommie and F. R. Fischer, Topological Band Engineering of Graphene Nanoribbons, *Nature*, 2018, **560**(7717), 204–208.

- 5 (a) E. H. Fort, P. M. Donovan and L. T. Scott, Diels-Alder Reactivity of Polycyclic Aromatic Hydrocarbon Bay Regions: Implications for Metal-Free Growth of Single-Chirality Carbon Nanotubes, J. Am. Chem. Soc., 2009, 131(44), 16006–16007; (b) R. Jasti and C. R. Bertozzi, Progress and Challenges for the Bottom-up Synthesis of Carbon Nanotubes with Discrete Chirality, Chem. Phys. Lett., 2010, 494(1), 1–7; (c) H. Omachi, T. Nakayama, E. Takahashi, Y. Segawa and K. Itami, Initiation of Carbon Nanotube Growth by Well-Defined Carbon Nanorings, Nat. Chem., 2013, 5(7), 572–576.
- 6 S. R. Peurifoy, J. C. Russell, T. J. Sisto, Y. Yang, X. Roy and C. Nuckolls, Designing Three-Dimensional Architectures for High-Performance Electron Accepting Pseudocapacitors, *J. Am. Chem. Soc.*, 2018, **140**(35), 10960– 10964.
- 7 (a) M. Iyoda, J. Yamakawa and M. J. Rahman, Conjugated Macrocycles: Concepts and Applications, *Angew. Chem., Int. Ed.*, 2011, 50(45), 10522–10553; (b) P. G. Ghasemabadi, T. Yao and G. J. Bodwell, Cyclophanes Containing Large Polycyclic Aromatic Hydrocarbons, *Chem. Soc. Rev.*, 2015, 44, 6494–6518.
- 8 (a) L. D. Tran, J. Ma, A. G. Wong-Foy and A. J. Matzger, A Coordination Perylene-Based Microporous Polymer Interacts Selectively with Electron-Poor Aromatics, Chem.-Eur. J., 2016, 22(16), 5509-5513; (b) W. B. Fellows, A. M. Rice, D. E. Williams, E. A. Dolgopolova, A. K. Vannucci, P. J. Pellechia, M. D. Smith, J. A. Krause and N. B. Shustova, Redox-Active Corannulene Buckybowls in a Crystalline Hybrid Scaffold, Angew. Chem., Int. Ed., 2016, 55(6), 2195–2199; (c) R. Dong, Z. Zhang, D. C. Tranca, S. Zhou, M. Wang, P. Adler, Z. Liao, F. Liu, Y. Sun, W. Shi, Z. Zhang, E. Zschech, S. C. B. Mannsfeld, C. Felser and Feng, A Coronene-Based Semiconducting Two-Х. Dimensional Metal-Organic Framework with Ferromagnetic Behavior, Nat. Commun., 2018, 9(1), 1-9; (d) J.-S. Qin, S. Yuan, L. Zhang, B. Li, D.-Y. Du, N. Huang, W. Guan, H. F. Drake, J. Pang, Y.-Q. Lan, A. Alsalme and H.-C. Zhou, Creating Well-Defined Hexabenzocoronene in Zirconium Metal-Organic Framework by Postsynthetic Annulation, J. Am. Chem. Soc., 2019, 141(5), 2054-2060; (e) S. Suginome, H. Sato, A. Hori, A. Mishima, Y. Harada, S. Kusaka, R. Matsuda, J. Pirillo, Y. Hijikata and T. Aida, One-Step Synthesis of an Adaptive Nanographene MOF: Adsorbed Gas-Dependent Geometrical Diversity, J. Am. Chem. Soc., 2019, 141(39), 15649-15655.

- 9 (a) A. Narita, X.-Y. Wang, X. Feng and K. Müllen, New Advances in Nanographene Chemistry, *Chem. Soc. Rev.*, 2015, 44, 6616–6643; (b) M. Stępień, E. Gońka, M. Żyła and N. Sprutta, Heterocyclic Nanographenes and Other Polycyclic Heteroaromatic Compounds: Synthetic Routes, Properties, and Applications, *Chem. Rev.*, 2017, 117(4), 3479–3716; (c) H. Ito, Y. Segawa, K. Murakami and K. Itami, Polycyclic Arene Synthesis by Annulative π-Extension, *J. Am. Chem. Soc.*, 2019, 141(1), 3–10.
- 10 General metal-mediated cycloaddition reviews: (a)K. P. C. Vollhardt, Cobalt-Mediated [2 + 2 + 2]-Cycloadditions: A Maturing Synthetic Strategy [New Synthetic Methods (43)], Angew. Chem., Int. Ed. Engl., 1984, 23(8), 539-556; (b) Transition-Metal-Mediated Aromatic Ring Construction, ed. K. Tanaka, John Wiley & Sons, Inc., Hoboken, NJ, 2013, pp. 1–320; Selected PAH applications: (c) T. Jin, J. Zhao, N. Asao and Y. Yamamoto, Metal-Catalyzed Annulation Reactions for π -Conjugated Polycycles, *Chem.-Eur. J.*, 2014, 20(13), 3554-3576; (d) S. Han, A. D. Bond, R. L. Disch, D. Holmes, J. M. Schulman, S. J. Teat, K. P. C. Vollhardt and G. D. Whitener, Total Syntheses and Structures of Angular [6]- and [7]Phenylene: The First Helical Phenylenes (Heliphenes), Angew. Chem., Int. Ed., 2002, 41(17), 3223-3227; (e) F. Teplý, I. G. Stará, I. Starý, A. Kollárovič, D. Šaman, L. Rulíšek and P. Fiedler, Synthesis of [5]-, [6]-, and [7]Helicene via Ni(0)- or Co(I)-Catalyzed Isomerization of Aromatic Cis, Dienetriynes, J. Am. Chem. Soc., 2002, 124(31), 9175-9180; (f) Y.-T. Wu, T. Hayama, K. K. Baldridge, A. Linden and Synthesis of Fluoranthenes J. S. Siegel, and Indenocorannulenes: Elucidation of Chiral Stereoisomers on the Basis of Static Molecular Bowls, J. Am. Chem. Soc., 2006, 128(21), 6870-6884; (g) D. Pérez, D. Peña and E. Guitián, Aryne Cycloaddition Reactions in the Synthesis of Large Polycyclic Aromatic Compounds, Eur. Chem., 2013, 2013(27), 5981-6013; (h) J. Org. T. Takahashi, M. Kitamura, B. Shen and K. Nakajima, Straightforward Method for Synthesis of Highly Alkyl-Substituted Naphthacene and Pentacene Derivatives by Homologation, J. Am. Chem. Soc., 2000, 122(51), 12876-12877; (i) Y. Sawada, S. Furumi, A. Takai, M. Takeuchi, Noguchi and K. Tanaka, Rhodium-Catalyzed K. Enantioselective Synthesis, Crystal Structures, and Photophysical Properties of Helically Chiral 1,1'-Bitriphenylenes, J. Am. Chem. Soc., 2012, 134(9), 4080-4083.
- 11 (a) G. R. Kiel, M. S. Ziegler and T. D. Tilley, Zirconacyclopentadiene-Annulated Polycyclic Aromatic Hydrocarbons, Angew. Chem., Int. Ed., 2017, 56(17), 4839– 4844; (b) G. R. Kiel, S. C. Patel, P. W. Smith, D. S. Levine and T. D. Tilley, Expanded Helicenes: A General Synthetic Strategy and Remarkable Supramolecular and Solid-State Behavior, J. Am. Chem. Soc., 2017, 139(51), 18456–18459; (c) G. R. Kiel, A. E. Samkian, A. Nicolay, R. J. Witzke and T. D. Tilley, Titanocene-Mediated Dinitrile Coupling: A Divergent Route to Nitrogen-Containing Polycyclic

Aromatic Hydrocarbons, J. Am. Chem. Soc., 2018, 140(7), 2450–2454.

- 12 F. Diederich, P. J. Stang and R. R. Tykwinski, *Acetylene Chemistry: Chemistry, Biology and Material Science*, John Wiley & Sons, 2006.
- 13 (a) D. Zhao and J. S. Moore, Shape-Persistent Arylene Ethynylene Macrocycles: Syntheses and Supramolecular Chemistry, Chem. Commun., 2003, 807-818; (b) S. Höger, Shape-Persistent Macrocycles: From Molecules to Materials, Chem.-Eur. J., 2004, 10(6), 1320-1329; (c) A. L. Sadowy and R. R. Tykwinski, Shape-Persistent Macrocycles Based on Acetylenic Scaffolding, in Modern Supramolecular Chemistry: Strategies for Macrocycle Synthesis, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, 2008, pp. 185-231; (d) M. K. Smith and O. Š. Miljanić, Arylene Ethynylene Macrocycles: From Molecular Hosts to Components of High-Performance Supramolecular Architectures, Org. Biomol. Chem., 2015, 13(29), 7841-7845.
- 14 V. H. Gessner, J. F. Tannaci, A. D. Miller and T. D. Tilley, Assembly of Macrocycles by Zirconocene-Mediated, Reversible Carbon–Carbon Bond Formation, *Acc. Chem. Res.*, 2011, 44(6), 435–446.
- 15 (a) J. K. Sprafke, D. V. Kondratuk, M. Wykes, A. L. Thompson, M. Hoffmann, R. Drevinskas, W.-H. Chen, C. K. Yong, J. Kärnbratt, J. E. Bullock, M. Malfois, M. R. Wasielewski, B. Albinsson, L. M. Herz, D. Zigmantas, D. Beljonne and H. L. Anderson, Belt-Shaped π-Systems: Relating Geometry to Electronic Structure in a Six-Porphyrin Nanoring, *J. Am. Chem. Soc.*, 2011, 133(43), 17262–17273; (b) N. Hayase, H. Sugiyama, H. Uekusa, Y. Shibata and K. Tanaka, Rhodium-Catalyzed Synthesis, Crystal Structures, and Photophysical Properties of [6]Cycloparaphenylene Tetracarboxylates, *Org. Lett.*, 2019, 21(11), 3895–3899.
- 16 (a) M. R. Golder and R. Jasti, Syntheses of the Smallest Carbon Nanohoops and the Emergence of Unique Physical Phenomena, Acc. Chem. Res., 2015, 48(3), 557-566; (b) Y. Segawa, A. Yagi, K. Matsui and K. Itami, Design and Synthesis of Carbon Nanotube Segments, Angew. Chem., Int. Ed., 2016, 55(17), 5136-5158; (c) E. Kayahara, T. Kouyama, T. Kato, H. Takaya, N. Yasuda and S. Yamago, Isolation and Characterization of the Cycloparaphenylene Radical Cation and Dication, Angew. Chem., Int. Ed., 2013, 52(51), 13722-13726; (d) Y. Xu and M. von Delius, The Supramolecular Chemistry of Strained Carbon Nanohoops, Angew. Chem., Int. Ed., 2020, 59(2), 559-573; (e) H. Miyoshi, S. Nobusue, A. Shimizu and Y. Tobe, Non-Alternant Non-Benzenoid Kekulenes: The Birth of a New Kekulene Family, Chem. Soc. Rev., 2015, 44, 6560-6577; (f) J. C. Buttrick and B. T. King, Kekulenes, Cycloarenes, and Heterocycloarenes: Addressing Electronic Structure and Aromaticity through Experiments and Calculations, Chem. Soc. Rev., 2017, 46, 7-20; (g) H. Gregolińska, M. Majewski, P. J. Chmielewski, J. Gregoliński, A. Chien, J. Zhou, Y.-L. Wu, Y. J. Bae, M. R. Wasielewski, P. M. Zimmerman and M. Stepień, Fully Conjugated [4]Chrysaorene. Redox-Coupled Anion Binding in a Tetraradicaloid Macrocycle, J.

Am. Chem. Soc., 2018, 140(43), 14474-14480; (h) C. Liu, M. E. Sandoval-Salinas, Y. Hong, T. Y. Gopalakrishna, H. Phan, N. Aratani, T. S. Herng, J. Ding, H. Yamada, Kim, D. Casanova and J. Wu, Macrocyclic D. Polyradicaloids with Unusual Super-Ring Structure and Global Aromaticity, Chem, 2018, 4(7), 1586-1595; (i) M. Iyoda and H. Shimizu, Multifunctional π -Expanded Oligothiophene Macrocycles, Chem. Soc. Rev., 2015, 44(18), 6411-6424; (j) S. Lee, C.-H. Chen and A. H. Flood, A Pentagonal Cyanostar Macrocycle with Cyanostilbene CH Donors Binds Anions and Forms Dialkylphosphate [3] Rotaxanes, Nat. Chem., 2013, 5(8), 704-710; (k)T. Schneebeli, M. Frasconi, Z. Liu, Y. Wu, S. D. M. Gardner, N. L. Strutt, C. Cheng, R. Carmieli, M. R. Wasielewski and J. F. Stoddart, Electron Sharing and Anion- π Recognition in Molecular Triangular Prisms, Angew. Chem., Int. Ed., 2013, 52(49), 13100-13104; (l) Z. Liu, G. Liu, Y. Wu, D. Cao, J. Sun, S. T. Schneebeli, M. S. Nassar, C. A. Mirkin and J. F. Stoddart, Assembly of Supramolecular Nanotubes from Molecular Triangles and 1,2-Dihalohydrocarbons, J. Am. Chem. Soc., 2014, 136(42), 16651-16660; (m) M. Ball, Y. Zhong, B. Fowler, B. Zhang, P. Li, G. Etkin, D. W. Paley, J. Decatur, A. K. Dalsania, H. Li, S. Xiao, F. Ng, M. L. Steigerwald and C. Nuckolls, Macrocyclization in the Design of Organic N-Type Electronic Materials, J. Am. Chem. Soc., 2016, 138(39), 12861-12867.

- 17 Y. Matano, M. Nakashima and H. Imahori, A Convenient Method for the Synthesis of α -Ethynylphospholes and Modulation of Their π -Conjugated Systems, *Angew. Chem.*, *Int. Ed.*, 2009, **48**(22), 4002–4005.
- 18 Note that Marder subsequently developed Rh- and Irmediated cyclizations on substrates similar to Matano's, but with aryl groups in place of the silyls. Here, the aryl groups may have played a similar role in shutting down competing intermolecular cycloaddition reactions. Rh: (a) A. Steffen, M. G. Tay, A. S. Batsanov, J. A. K. Howard, A. Beeby, K. Q. Vuong, X.-Z. Sun, M. W. George and T. B. Marder, 2,5-Bis(p-R-Arylethynyl) Rhodacyclopentadienes Show Intense Fluorescence: Denying the Presence of a Heavy Atom, Angew. Chem., Int. Ed., 2010, 49, 2349-2353; Ir: (b) A. Steffen, K. Costuas, A. Boucekkine, M.-H. Thibault, A. Beeby, A. S. Batsanov, Charaf-Eddin, D. Jacquemin, J.-F. Halet A. and Т. B. Marder, Fluorescence in Rhodaand Iridacyclopentadienes Neglecting the Spin-Orbit Coupling of the Heavy Atom: The Ligand Dominates, Inorg. Chem., 2014, 53, 7055-7069.
- 19 Z. Huang and G. Dong, Site-Selectivity Control in Organic Reactions: A Quest To Differentiate Reactivity among the Same Kind of Functional Groups, *Acc. Chem. Res.*, 2017, 50(3), 465–471.
- 20 M. Kivala, D. Wu, X. Feng, C. Li and K. Müllen, Cyclodehydrogenation in the Synthesis of Graphene-Type Molecules, in *Materials Science and Technology*, Wiley-VCH Verlag GmbH & Co. KGaA, 2006.

- 21 T. Makino, Y. Yamamoto and K. Itoh, Synthesis and Structure of Novel Iridium(I) Complexes Containing η^4 -1,6-Diene and Diphosphine Ligands: Remarkable Effect of Ligand Bite Angle upon Ligand Dissociation, Organometallics, 2004, 23(8), 1730–1737.
- 22 (*a*) S. Kezuka, S. Tanaka, T. Ohe, Y. Nakaya and R. Takeuchi, Iridium Complex-Catalyzed [2 + 2 + 2] Cycloaddition of α, ω -Diynes with Monoynes and Monoenes, *J. Org. Chem.*, 2006, **71**(2), 543–552; (*b*) R. Takeuchi and Y. Nakaya, Iridium Complex-Catalyzed Highly Selective Cross [2 + 2 + 2]Cycloaddition of Two Different Monoynes: 2:1 Coupling versus 1:2 Coupling, *Org. Lett.*, 2003, 5(20), 3659–3662.
- 23 (a) U. Rosenthal, A. Ohff, W. Baumann, A. Tillack, H. Görls, V. V. Burlakov and V. B. Shur, Struktur, Eigenschaften und NMR-spektroskopische Charakterisierung von Cp₂Zr(Pyridin)(Me₃SiC≡CSiMe₃), Z. Anorg. Allg. Chem., 1995, 621(1), 77–83; (b) J. R. Nitschke, S. Zürcher and T. D. Tilley, New Zirconocene-Coupling Route to Large, Functionalized Macrocycles, J. Am. Chem. Soc., 2000, 122(42), 10345–10352.
- 24 G. Erker and R. Zwettler, Reactions of in situ Generated (η₂-Phenyltrimethylsilylacetylene)Zirconocene, *J. Organomet. Chem.*, 1991, **409**(1), 179–188.
- 25 The low yield may result from kinetic traps such as oxidative addition of the C–Br bond to "Cp₂Zr".
- 26 (a) P. J. Fagan, W. A. Nugent and J. C. Calabrese, Metallacycle Transfer from Zirconium to Main Group Elements: A Versatile Synthesis of Heterocycles, J. Am. Chem. Soc., 1994, 116(5), 1880-1889; (b) T. Takahashi, R. Hara, Y. Nishihara and М. Kotora, Copper-Mediated Coupling of Zirconacyclopentadienes with Aromatic Dihalo Compounds. Formation of Fused Aromatic Rings, J. Am. Chem. Soc., 1996, 118(21), 5154-5155; (c) T. Takahashi, Y. Li, P. Stepnicka, M. Kitamura, Y. Liu, K. Nakajima and M. Kotora, Coupling Reaction of Zirconacyclopentadienes with Dihalonaphthalenes and Dihalopyridines: A New Procedure for the Preparation of Substituted Anthracenes, Quinolines, and Isoquinolines, J. Am. Chem. Soc., 2002, 124(4), 576-582; (d) X. Yan and C. Xi, Conversion of Zirconacyclopentadienes into Metalloles: Fagan-Nugent Reaction and Beyond, Acc. Chem. Res., 2015, 48(4), 935-946.
- 27 (a) P.-H. Ge, W. Fu, W. A. Herrmann, E. Herdtweck, C. Campana, R. D. Adams and U. H. F. Bunz, Structural Characterization of Cyclohexameric Metaа Phenyleneethynylene Made by Alkyne Metathesis with In Situ Catalysts, Angew. Chem., Int. Ed., 2000, 39(20), 3607-3610; (b) O. Š. Miljanić, K. P. C. Vollhardt and G. D. Whitener, An Alkyne Metathesis-Based Route to Ortho-Dehydrobenzannulenes, Synlett, 2003, 1, 29-34; (c) W. Zhang and J. S. Moore, Arylene Ethynylene Macrocycles Prepared by Precipitation-Driven Alkyne Metathesis, J. Am. Chem. Soc., 2004, 126(40), 12796; (d) C. Yu, Y. Jin and W. Zhang, Shape-Persistent Macrocycles through Dynamic Covalent Reactions, in Dynamic Covalent Chemistry, John Wiley & Sons, Ltd, 2017, pp. 121-163.
- 28 (a) S. Höger, X. H. Cheng, A.-D. Ramminger, V. Enkelmann, A. Rapp, M. Mondeshki and I. Schnell, Discotic Liquid

Crystals with an Inverted Structure, Angew. Chem., Int. Ed., 2005, 44(18), 2801-2805; (b) W.-J. Liu, Y. Zhou, O.-F. Zhou, Y. Ma and J. Pei, Shape-Persistent Elliptic Macrocycles Composed of Polycyclic Aromatic Hydrocarbons: Synthesis and Photophysical Properties, Org. Lett., 2008, 10(11), 2123-2126; (c) J. M. W. Chan, J. R. Tischler, S. E. Kooi, V. Bulović and T. M. Swager, Synthesis of J-Aggregating Dibenz[a,j]Anthracene-Based Macrocycles, J. Am. Chem. Soc., 2009, 131(15), 5659-5666; (d) L. Zhang, H. Gopee, D. L. Hughes and A. N. Cammidge, Antiaromatic Twinned Triphenylene Discotics Showing Nematic Phases and 2-Dimensional π -Overlap in the Solid State, *Chem. Commun.*, 2010, 46, 4255-4257; (e) T. Chen, G.-B. Pan, H. Wettach, M. Fritzsche, S. Höger, L.-J. Wan, H.-B. Yang, B. H. Northrop and P. J. Stang, 2D Assembly of Metallacycles on HOPG by Shape-Persistent Macrocycle Templates, J. Am. Chem. Soc., 2010, 132(4), 1328-1333; (f) A. D. Finke, D. E. Gross, A. Han and J. S. Moore, Engineering Solid-State Morphologies in Carbazole-Ethynylene Macrocycles, J. Am. Chem. Soc., 2011, 133(35), 14063–14070; (g) G. Venkataramana, P. Dongare, L. N. Dawe, D. W. Thompson, Y. Zhao and G. J. Bodwell, 1,8-Pyrenylene-Ethynylene Macrocycles, Org. Lett., 2011, 13(9), 2240-2243; (h) S. Toyota, Construction of Novel Molecular Architectures from Anthracene Units and Acetylene Linkers, Pure Appl. Chem., 2012, 84(4), 917-929; (i) Z. He, X. Xu, X. Zheng, T. Ming and Q. Miao, Conjugated Macrocycles of Phenanthrene: A New Segment of [6,6]-Carbon Nanotube and Solution-Processed Organic Semiconductors, Chem. Sci., 2013, 4(12), 4525-4531; (j) H. Yang, Y. Du, S. Wan, G. D. Trahan, Y. Jin and W. Zhang, Mesoporous 2D Covalent Organic Frameworks Based on Shape-Persistent Arylene-Ethynylene Macrocycles, Chem. Sci., 2015, 6(7), 4049-4053; (k) Y. Takaki, R. Ozawa, T. Kajitani, T. Fukushima, M. Mitsui and K. Kobayashi, Synthesis and Self-Assembly of Cyclic 2,7-Anthrylene Ethynylene 1,3-Phenylene Ethynylene Trimer with a Planar Conformation, Chem.-Eur. J., 2016, 22(47), 16760-16764.

- 29 J. Heppekausen, R. Stade, R. Goddard and A. Fürstner, Practical New Silyloxy-Based Alkyne Metathesis Catalysts with Optimized Activity and Selectivity Profiles, *J. Am. Chem. Soc.*, 2010, **132**(32), 11045–11057.
- 30 S. Lee, A. Yang, T. P. Moneypenny and J. S. Moore, Kinetically Trapped Tetrahedral Cages via Alkyne Metathesis, *J. Am. Chem. Soc.*, 2016, **138**(7), 2182–2185.
- 31 R. L. Gdula and M. J. A. Johnson, Highly Active Molybdenum–Alkylidyne Catalysts for Alkyne Metathesis: Synthesis from the Nitrides by Metathesis with Alkynes, *J. Am. Chem. Soc.*, 2006, **128**(30), 9614–9615.
- 32 (a) A. Fürstner, Alkyne Metathesis on the Rise, Angew. Chem., Int. Ed., 2013, 52, 2794–2819; (b) M. Ortiz, C. Yu, Y. Jin and W. Zhang, Poly(Aryleneethynylene)s: Properties, Applications and Synthesis Through Alkyne Metathesis, Top. Curr. Chem., 2017, 375(4), 69; (c) H. Ehrhorn and M. Tamm, Well-Defined Alkyne Metathesis Catalysts: Developments and Recent Applications, Chem.–Eur. J., 2019, 25(13), 3190–3208.