

Cite this: *Chem. Sci.*, 2020, 11, 3068

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 16th December 2019

Accepted 18th February 2020

DOI: 10.1039/c9sc06342k

rsc.li/chemical-science

Chiral *N,N'*-dioxide/Mg(OTf)<sub>2</sub> complex-catalyzed asymmetric [2,3]-rearrangement of *in situ* generated ammonium salts†Qianchi Lin,<sup>ID</sup> Bowen Hu,<sup>ID</sup> Xi Xu,<sup>ID</sup> Shunxi Dong,<sup>ID</sup> Xiaohua Liu<sup>ID</sup>\* and Xiaoming Feng<sup>ID</sup>\*Catalytic enantioselective [2,3]-rearrangements of *in situ* generated ammonium ylides from glycine pyrazoleamides and allyl bromides were achieved by employing a chiral *N,N'*-dioxide/Mg<sup>II</sup> complex as the catalyst. This protocol provided a facile and efficient synthesis route to a series of *anti*- $\alpha$ -amino acid derivatives in good yields with high stereoselectivities. Moreover, a possible catalytic cycle was proposed to illustrate the reaction process and the origin of stereoselectivity.

## Introduction

[2,3]-Rearrangements have been regarded as a class of synthetically powerful organic transformations due to their inherently high efficiency.<sup>1</sup> In particular, [2,3]-rearrangement of ammonium ylides has been extensively investigated for rapid construction of valuable nitrogen-containing molecules.<sup>2</sup> It is highly attractive to disclose asymmetric versions of such intriguing rearrangement,<sup>3–5</sup> but only a few examples concerning the catalytic enantioselective [2,3]-rearrangement of ammonium ylides have been reported to date.<sup>5</sup> In 2014, Smith and co-workers demonstrated the first example of chiral isothiurea-catalyzed [2,3]-rearrangement of allylic ammonium ylides to gain optically enriched *syn*-configured  $\alpha$ -amino acid derivatives (Scheme 1a).<sup>5a</sup> In 2017, they developed an elegant tandem *in situ* protocol utilizing Pd/chiral isothiurea relay catalysis, which provides a direct method for the synthesis of *syn*- $\alpha$ -amino acid derivatives from *N,N*-disubstituted glycine aryl esters and allylic phosphates (Scheme 1b).<sup>5c</sup> Recently, the group of Song reported an interesting study on an isothiurea catalyzed asymmetric [2,3]-rearrangement reaction of propargyl ammonium salts, which allows access to optically active allenyl  $\alpha$ -amino amides.<sup>5f</sup> Despite these impressive advances, there is still room for further development. For instance, chiral isothiurea is a unique catalyst with which *syn*- $\alpha$ -amino acid derivatives were preferentially afforded in current reports.<sup>5a–e</sup> Given the wide and versatile use of  $\alpha$ -amino acids in organic


synthesis and pharmaceutical chemistry,<sup>6</sup> it is highly desirable to search for new catalytic systems for [2,3]-rearrangement of ammonium ylides in terms of the catalyst, substrate scope and the method of ammonium ylide formation, as well as stereo-divergence of products.<sup>7</sup>

Inspired by the achievements in enantioselective [2,3]-rearrangement<sup>1–5,8</sup> and our ongoing interest in synthesis of unnatural  $\alpha$ -amino acid derivatives,<sup>9</sup> we envisaged that chiral *N,N'*-dioxide-metal complex catalysts<sup>10</sup> developed by our group would be potentially useful in promoting asymmetric [2,3]-rearrangement of allylic ammonium ylides upon rationally introducing pyrazoleamide groups.<sup>11</sup> As depicted in Scheme 1c, pyrazoleamide ammonium salts generated from glycine



Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China. E-mail: liuxh@scu.edu.cn; xmfeng@scu.edu.cn

† Electronic supplementary information (ESI) available: [<sup>1</sup>H, <sup>13</sup>C{<sup>1</sup>H}] and <sup>19</sup>F{<sup>1</sup>H} NMR, HPLC spectra, and CD spectra (PDF). X-ray crystallographic data for **4u** (CIF). CCDC 1960932. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc06342k

(a) Isothiurea-Catalyzed Enantioselective Catalytic [2,3]-Rearrangement (Smith, 2014)



(b) Enantioselective Pd/Isothiurea Relay Catalysis (Smith, 2017)

(c) Lewis Acid-Enabled Asymmetric [2,3]-Rearrangement with *anti*-selectivity (This work)

## Features:

- First example of Lewis acid-catalyzed [2,3]-rearrangement of ammonium salts
- *Anti*-selectivity, good to excellent enantioselectivity under strong background reaction
- One-pot procedure, easy operation

Scheme 1 [2,3]-Rearrangements of allylic ammonium ylides.




pyrazoleamides **1** and allyl bromides **2** were selected as the precursors of [2,3]-rearrangement. It was thought that such allylic ammonium salts could be activated by bidentate coordination with the chiral *N,N'*-dioxide–metal complex and then subjected to deprotonation with the assistance of an external base to afford ammonium ylides, which subsequently undergo [2,3]-rearrangement to deliver non-racemic  $\alpha$ -amino acid derivatives. There are some difficulties associated with the catalytic asymmetric [2,3]-rearrangement, such as the compatibility of all reactants with the catalyst<sup>5a,f</sup> and the background reaction in the presence of the external base.<sup>12</sup> Herein, we wish to disclose our effort toward one-pot asymmetric [2,3]-rearrangement of *in situ* formed allylic ammonium ylides. Chiral *N,N'*-dioxide/Mg(OTf)<sub>2</sub> (ref. 13) was found to promote the diastereo- and enantioselective rearrangement efficiently, and various *anti*- $\alpha$ -amino acid derivatives<sup>14</sup> were readily obtained in good yields with high stereoselectivities (up to 95% yield, >19 : 1 *anti* : *syn* and 98.5 : 1.5 er) from easily available glycine pyrazoleamides and allyl bromides.

## Results and discussion

In the initial study, *N,N*-dimethylglycine pyrazoleamide (**1a**) and cinnamyl bromide (**2a**) were selected as the model

substrates to optimize the reaction conditions. The preliminary study indicated that the tandem ammonium salt formation/[2,3]-rearrangement took place well in the presence of an external base, and the desired product **3a** was isolated in 91% yield with a 1 : 1 *anti* : *syn* ratio by using diisopropylamine as the base (see page 9 in the ESI† for more details). This result showed the difficulty in achieving the asymmetric version of such one-pot transformation. For the purpose of determination of the er value, compound **3a** was converted to **4a** quantitatively with MeOH at 60 °C for further analysis. Next, different metal salts coordinated with chiral *N,N'*-dioxide **L-PrMe<sub>2</sub>** were examined in CH<sub>2</sub>Cl<sub>2</sub> at 30 °C (Table 1, entries 1–4). Mg(OTf)<sub>2</sub> performed better than other metal salts, giving the desired rearranged product **3a** in 91% yield, 1.8 : 1 *anti* : *syn*, and 67.5 : 32.5 er for the major diastereomer (Table 1, entry 3). The complex of Mg(NTf)<sub>2</sub> provided a comparable result (Table 1, entry 4). Subsequently, the amide moiety in the ligand was evaluated, and it was found that sterically bulky 1-adamantyl amine derived **L-PrAd** afforded better stereoselectivities (Table 1, entry 7 *vs.* entries 5 and 6). Further investigations on the chiral backbone in the ligand showed that the *L*-ramipril-derived **L-RaAd** was superior to the *L*-proline-derived **L-PrAd** and (*S*)-pipercolic acid-derived **L-PiAd** in terms of enantioselectivity (91% yield, 3 : 1 *anti* : *syn*, and 81 : 19 er; entries 7–9).

Table 1 Optimization of the reaction conditions<sup>a</sup>



Entry	Metal salt	Ligand	Yield of <b>3a</b> <sup>b</sup> (%)	<i>anti</i> : <i>syn</i> of <b>3a</b> <sup>c</sup>	er of <b>4a</b> <sup>d</sup>
1	Sc(OTf) <sub>3</sub>	<b>L-PrMe<sub>2</sub></b>	77	1 : 1	race/race
2	Ni(OTf) <sub>2</sub>	<b>L-PrMe<sub>2</sub></b>	90	1.1 : 1	54 : 46/57.5 : 42.5
3	Mg(OTf) <sub>2</sub>	<b>L-PrMe<sub>2</sub></b>	91	1.8 : 1	67.5 : 32.5/67.5 : 32.5
4	Mg(NTf) <sub>2</sub>	<b>L-PrMe<sub>2</sub></b>	93	1.6 : 1	65.5 : 34.5/64.5 : 35.5
5	Mg(OTf) <sub>2</sub>	<b>L-PrEt<sub>2</sub></b>	99	3 : 1	61 : 39/80 : 20
6	Mg(OTf) <sub>2</sub>	<b>L-Pr<i>t</i>Bu</b>	90	2 : 1	56.5 : 43.5/52.5 : 47.5
7	Mg(OTf) <sub>2</sub>	<b>L-PrAd</b>	98	5 : 1	75.5 : 24.5/57 : 43
8	Mg(OTf) <sub>2</sub>	<b>L-PiAd</b>	94	2 : 1	52.5 : 47.5/race
9	Mg(OTf) <sub>2</sub>	<b>L-RaAd</b>	91	3 : 1	81 : 19/54.5 : 45.5
10 <sup>e</sup>	Mg(OTf) <sub>2</sub>	<b>L-RaAd</b>	99	13 : 1	93.5 : 6.5
11 <sup>e,f</sup>	Mg(OTf) <sub>2</sub>	<b>L-RaAd</b>	82	12 : 1	95.5 : 4.5
12 <sup>e,f,g</sup>	Mg(OTf) <sub>2</sub>	<b>L-RaAd</b>	94	>19 : 1	97 : 3

<sup>a</sup> Unless otherwise noted, all reactions were carried out with **1a** (0.10 mmol), **2a** (0.10 mmol), *i*Pr<sub>2</sub>NH (0.15 mmol) and metal salt/ligand (1 : 1, 10 mol%) in CH<sub>2</sub>Cl<sub>2</sub> (1.0 mL) at 30 °C for 14 h. <sup>b</sup> Isolated yield of **3a**. <sup>c</sup> Determined by <sup>1</sup>H NMR. <sup>d</sup> Determined by HPLC on a chiral stationary phase. <sup>e</sup> Carried out in MeCN. The metal salt/ligand was pretreated in CH<sub>2</sub>Cl<sub>2</sub>. <sup>f</sup> –20 °C for 24 h. <sup>g</sup> NaBAR<sub>4</sub><sup>F</sup> {NaB[3,5-(F<sub>3</sub>C)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>]<sub>4</sub>} (20 mol%) was added. The metal salt/ligand/NaBAR<sub>4</sub><sup>F</sup> was pretreated in EtOAc instead of CH<sub>2</sub>Cl<sub>2</sub>. Tf = trifluoromethanesulfonyl.



Other reaction parameters were investigated and the solvent was proven to play a significant role in the reaction. With MeCN as the solvent, the amino acid derivative **3a** was produced in 99% yield with 13 : 1 *anti* : *syn* and 93.5 : 6.5 er for the major diastereomer (entry 10). The enantioselectivity could be further enhanced when the reaction was performed at decreased temperature ( $-20\text{ }^{\circ}\text{C}$ )<sup>15</sup> while a slightly lower yield was obtained (82% yield, 12 : 1 *anti* : *syn*, 95.5 : 4.5 er; entry 11). To our delight, the addition of NaBAR<sub>4</sub><sup>F</sup> as an additive and preparation of the catalyst in EtOAc produced optimized results (entry 12; 94% yield, >19 : 1 *anti* : *syn*, 97 : 3 er).

Moreover, when the product with a low *anti* : *syn* ratio was subjected to the reaction conditions, it was found that both *anti* : *syn* ratios and er values did not change. This result implied that an ultimately high *anti* : *syn* ratio was obtained during the [2,3]-rearrangement rather than epimerization of the product during the reaction (for further details, see ESI,† page 11).

With the optimized reaction conditions in hand, the substrate scope of [2,3]-rearrangement was screened (Table 2). Varying *N*-substituents in the glycine pyrazoleamides had no effect on the *anti* : *syn* ratio of the reaction (>19 : 1 in all cases).

Table 2 Substrate scope for the [2,3]-rearrangement<sup>a</sup>

<b>3a</b> , 24 h, 94% yield >19:1 <i>anti</i> : <i>syn</i> , 97:3 er <sup>c</sup>	<b>3b</b> , 168 h, 64% yield >19:1 <i>anti</i> : <i>syn</i> , 95:5 er <sup>c</sup>	<b>3c</b> , 168 h, 71% yield >19:1 <i>anti</i> : <i>syn</i> , 94.5:5.5 er <sup>c</sup>	<b>3d</b> , 72 h, 65% yield >19:1 <i>anti</i> : <i>syn</i> , 90.5:9.5 er <sup>c</sup>	<b>3e</b> , 168 h, 27% yield >19:1 <i>anti</i> : <i>syn</i> , 92:8 er	<b>3f</b> , 144 h, 60% yield 1:2 <i>anti</i> : <i>syn</i> , 65.5:34.5/54.5:45.5 er <sup>c</sup>
<b>3g</b> , 24 h, 88% yield >19:1 <i>anti</i> : <i>syn</i> , 95:5 er <sup>c</sup>	<b>3h</b> , 56 h, 63% yield <sup>b</sup> 6:1 <i>anti</i> : <i>syn</i> , 94:6 er	<b>3i</b> , 56 h, 75% yield <sup>b</sup> 7:1 <i>anti</i> : <i>syn</i> , 94.5:5.5 er	<b>3j</b> , 56 h, 57% yield <sup>b</sup> 8:1 <i>anti</i> : <i>syn</i> , 90.5:9.5 er	<b>3k</b> , 120 h, 62% yield <sup>b</sup> 5:1 <i>anti</i> : <i>syn</i> , 95.5:4.5 er	<b>3l</b> , 120 h, 70% yield <sup>b</sup> 4:1 <i>anti</i> : <i>syn</i> , 96:4 er
<b>3m</b> , 56 h, 57% yield <sup>b</sup> 4:1 <i>anti</i> : <i>syn</i> , 92:8 er	<b>3n</b> , 144 h, 54% yield, 3:7 <i>anti</i> : <i>syn</i> , 71:29/53:47 er <sup>c</sup>	<b>3o</b> , 72 h, 83% yield >19:1 <i>anti</i> : <i>syn</i> , 96:4 er <sup>c</sup>	<b>3p</b> , 48 h, 85% yield 14:1 <i>anti</i> : <i>syn</i> , 92.5:7.5 er <sup>c</sup>	<b>3q</b> , 56 h, 85% yield <sup>b</sup> >19:1 <i>anti</i> : <i>syn</i> , 96.5:3.5 er	<b>3r</b> , 24 h, 91% yield <sup>b</sup> >19:1 <i>anti</i> : <i>syn</i> , 98:2 er
<b>3s</b> , 168 h, 60% yield >19:1 <i>anti</i> : <i>syn</i> , 90:10 er	<b>3t</b> , 96 h, 82% yield <sup>b</sup> >19:1 <i>anti</i> : <i>syn</i> , 97:3 er	<b>3u</b> , 24 h, 95% yield >19:1 <i>anti</i> : <i>syn</i> , 98.5:1.5 er <sup>c</sup>	<b>4u</b> , 92% yield >19:1 <i>anti</i> : <i>syn</i> , 98.5:1.5 er	<b>(2<i>R</i>,3<i>S</i>)-4u</b> (CCDC: 1960932)	<b>3v</b> , 168 h, trace

<sup>a</sup> All reactions were carried out with L-RaAd/Mg(OTf)<sub>2</sub>/NaBAR<sub>4</sub><sup>F</sup> (1 : 1 : 2, 10 mol%), **1** (0.2 mmol), **2** (0.2 mmol), and iPr<sub>2</sub>NH (0.3 mmol) in MeCN (2.0 mL) at  $-20\text{ }^{\circ}\text{C}$ . Isolated total yield of product **3**. The *anti* : *syn* ratio was determined by <sup>1</sup>H NMR analysis. The er value was determined by HPLC on a chiral stationary phase. <sup>b</sup> Isolated yield of the major diastereomer **3**. <sup>c</sup> er value of product **4**.





## Conclusions

We have successfully developed the first Lewis acid catalyzed asymmetric [2,3]-rearrangement of quaternary ammonium ylides formed *in situ* from glycine pyrazoleamides and allyl bromides. The  $N,N'$ -dioxide/Mg(OTf)<sub>2</sub> catalytic system benefited the rearrangement process efficiently, providing diverse chiral *anti*- $\alpha$ -amino acid derivatives in good yields with high stereoselectivities (up to 95% yield, >19 : 1 *anti* : *syn* and 98.5 : 1.5 er). Besides, the potential use of the current method was illustrated by gram-scale synthesis and further transformations of products. A possible catalytic cycle along with the working mode was proposed to elucidate the reaction process and chiral induction. Further investigations on other reactions enabled by chiral  $N,N'$ -dioxide-metal complex catalysts are ongoing in our laboratory.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We appreciate the National Natural Science Foundation of China (No. 21625205 and 21772127) for financial support.

## Notes and references

- For selected books and reviews on [2,3]-rearrangements, see: (a) A.-H. Li, L.-X. Dai and V. K. Aggarwal, *Chem. Rev.*, 1997, **97**, 2341; (b) A. P. A. Arboré, D. J. Cane-Honeysett, I. Coldham and M. L. Middleton, *Synlett*, 2000, **2000**, 236; (c) J. B. Sweeney, *Chem. Soc. Rev.*, 2009, **38**, 1027; (d) *Nitrogen, Oxygen and Sulfur Ylide Chemistry: A Practical Approach in Chemistry*, ed. J. S. Clark, Oxford University Press, New York, 2002; (e) H. L. Bao and U. K. Tambar, [2,3]-Rearrangements of Ammonium Zwitterions, in *Molecular Rearrangements in Organic Synthesis*, ed. C. M. Rojas, John Wiley & Sons, Inc., Hoboken, N.J., USA, 2015, pp 459–496; (f) A. C. Jones, J. A. May, R. Sarpong and B. M. Stoltz, *Angew. Chem., Int. Ed.*, 2014, **53**, 2556; (g) T. H. West, S. S. M. Spoehrle, K. Kasten, J. E. Taylor and A. D. Smith, *ACS Catal.*, 2015, **5**, 7446; (h) Z. Sheng, Z. K. Zhang, C. H. Chu, Y. Zhang and J. B. Wang, *Tetrahedron*, 2017, **73**, 4011; (i) H. Wu, Q. Wang and J. P. Zhu, *Eur. J. Org. Chem.*, 2019, **2019**, 1964.
- For recent examples of [2,3]-rearrangement of ammonium ylides to synthesize useful natural products or active molecules, see: (a) J. A. Vanecko, H. Wan and F. G. West, *Tetrahedron*, 2006, **62**, 1043; (b) H. Cho, H. Jeon, J. E. Shin, S. Lee, S. Park and S. Kim, *Chem.–Eur. J.*, 2019, **25**, 2447; (c) H. Cho, J. Jung, J. Kim, S. Park and S. Kim, *Asian J. Org. Chem.*, 2019, **8**, 1010; (d) H. Jeon, H. Cho and S. Kim, *Org. Lett.*, 2019, **21**, 1121.
- For selected examples of asymmetric [2,3]-rearrangement of ammonium ylides with chiral auxiliaries or substrate control, see: (a) K. Hiroi and K. Nakazawa, *Chem. Lett.*, 1980, 1077; (b) J. A. Workman, N. P. Garrido, J. Sançon, E. Roberts, H. P. Wessel and J. B. Sweeney, *J. Am. Chem. Soc.*, 2005, **127**, 1066; (c) T.-S. Zhu and M.-H. Xu, *Chem. Commun.*, 2012, **48**, 7274; (d) E. Tayama, N. Naganuma, H. Iwamoto and E. Hasegawa, *Chem. Commun.*, 2014, **50**, 6860.
- For selected Lewis acid mediated asymmetric [2,3]-rearrangements of ammonium ylides, see: (a) J. Blid and P. Somfai, *Tetrahedron Lett.*, 2003, **44**, 3159; (b) J. Blid, P. Brandt and P. Somfai, *J. Org. Chem.*, 2004, **69**, 3043; (c) J. Blid, O. Panknin, P. Tuzina and P. Somfai, *J. Org. Chem.*, 2007, **72**, 1294.
- (a) T. H. West, D. S. B. Daniels, A. M. Z. Slawin and A. D. Smith, *J. Am. Chem. Soc.*, 2014, **136**, 4476; (b) T. H. West, D. M. Walden, J. E. Taylor, A. C. Brueckner, R. C. Johnston, P. H.-Y. Cheong, G. C. Lloyd-Jones and A. D. Smith, *J. Am. Chem. Soc.*, 2017, **139**, 4366; (c) T. H. West, S. S. M. Spoehrle and A. D. Smith, *Tetrahedron*, 2017, **73**, 4138; (d) K. Kasten, A. M. Z. Slawin and A. D. Smith, *Org. Lett.*, 2017, **19**, 5182; (e) S. S. M. Spoehrle, T. H. West, J. E. Taylor, A. M. Z. Slawin and A. D. Smith, *J. Am. Chem. Soc.*, 2017, **139**, 11895; (f) L. Zhang, Z.-J. Zhang, J.-Y. Xiao and J. Song, *Org. Lett.*, 2018, **20**, 5519.
- For selected reviews, see: (a) E. R. Jarvo and S. J. Miller, *Tetrahedron*, 2002, **58**, 2481; (b) G. Prabhu, N. Narendra, Basavaprabhu, V. Panduranga and V. V. Sureshbabu, *RSC Adv.*, 2015, **5**, 48331; (c) S.-M. Paek, M. Jeong, J. Jo, Y. M. Heo, Y. T. Han and H. Yun, *Molecules*, 2016, **21**, 951; (d) P. Singh, S. L. K. Manda, K. Samanta and G. Panda, *Tetrahedron*, 2017, **73**, 1911; (e) E. D. Carpio, L. Hernández, C. Ciangherotti, V. V. Coa, L. Jiménez, V. Lubes and G. Lubes, *Coord. Chem. Rev.*, 2018, **372**, 117.
- For selected reviews, see: (a) Y. Ohfuné and T. Shinada, *Eur. J. Org. Chem.*, 2005, 5127; (b) H. Vogt and S. Bräse, *Org. Biomol. Chem.*, 2007, **5**, 406; (c) K. Bera and I. N. N. Namboothiri, *Asian J. Org. Chem.*, 2014, **3**, 1234; (d) Y. B. Wang, X. H. Song, J. Wang, H. Moriwaki, V. A. Soloshonok and H. Liu, *Amino Acids*, 2017, **49**, 1487.
- For selected reviews, see: (a) M. P. Doyle and D. C. Forbes, *Chem. Rev.*, 1998, **98**, 911; (b) A. Moyano, N. El-Hamdouni and A. Atlamsani, *Chem.–Eur. J.*, 2010, **16**, 5260. For selected examples of enantioselective [2,3]-rearrangement in our laboratory, see: (c) X. Xu, J. L. Zhang, S. X. Dong, L. L. Lin, X. B. Lin, X. H. Liu and X. M. Feng, *Angew. Chem., Int. Ed.*, 2018, **57**, 8734; (d) X. B. Lin, Y. Tang, W. Yang, F. Tan, L. L. Lin, X. H. Liu and X. M. Feng, *J. Am. Chem. Soc.*, 2018, **140**, 3299; (e) X. B. Lin, W. Yang, W. K. Yang, X. H. Liu and X. M. Feng, *Angew. Chem., Int. Ed.*, 2019, **58**, 13492.
- (a) D. J. Shang, Y. L. Liu, X. Zhou, X. H. Liu and X. M. Feng, *Chem.–Eur. J.*, 2009, **15**, 3678; (b) J. Yang, P. R. Ruan, W. Yang, X. M. Feng and X. H. Liu, *Chem. Sci.*, 2019, **10**, 10305.
- For reviews on  $N,N'$ -dioxide ligands, see: (a) X. H. Liu, L. L. Lin and X. M. Feng, *Acc. Chem. Res.*, 2011, **44**, 574; (b) X. H. Liu, L. L. Lin and X. M. Feng, *Org. Chem. Front.*, 2014, **1**, 298; (c) X. H. Liu, H. F. Zheng, Y. Xia, L. L. Lin and X. M. Feng, *Acc. Chem. Res.*, 2017, **50**, 2621; (d)



- X. H. Liu, S. X. Dong, L. L. Lin and X. M. Feng, *Chin. J. Chem.*, 2018, **36**, 791.
- 11 (a) Q. Yao, Z. Wang, Y. H. Zhang, X. H. Liu, L. L. Lin and X. M. Feng, *J. Org. Chem.*, 2015, **80**, 5704; (b) Y. Zhang, Y. T. Liao, X. H. Liu, Q. Yao, Y. H. Zhou, L. L. Lin and X. M. Feng, *Chem.-Eur. J.*, 2016, **22**, 15119; (c) G. J. Wang, Y. Tang, Y. Zhang, X. H. Liu, L. L. Lin and X. M. Feng, *Chem.-Eur. J.*, 2017, **23**, 554.
- 12 For a recent example, see: A. Murre, K. Erkman, S. Kaabel, I. Järving and T. Kanger, *Synthesis*, 2019, **51**, 4183–4197.
- 13 For a recent review on magnesium catalysis, see: (a) D. X. Yang, L. Q. Wang, D. Li and R. Wang, *Chem*, 2019, **5**, 1108. For selected examples of reactions catalyzed by the *N,N'*-dioxide/Mg<sup>II</sup> complex, see: (b) X. H. Zhao, X. H. Liu, H. J. Mei, J. Guo, L. L. Lin and X. M. Feng, *Angew. Chem., Int. Ed.*, 2015, **54**, 4032; (c) X. Zhong, Q. Tang, P. F. Zhou, Z. W. Zhong, S. X. Dong, X. H. Liu and X. M. Feng, *Chem. Commun.*, 2018, **54**, 10511; (d) Q. Xiong, S. X. Dong, Y. S. Chen, X. H. Liu and X. M. Feng, *Nat. Commun.*, 2019, **10**, 2116; (e) Q. Xiong, G. L. Li, S. X. Dong, X. H. Liu and X. M. Feng, *Org. Lett.*, 2019, **12**, 8771.
- 14 Tambar and co-workers reported several fascinating examples of highly diastereoselective synthesis of *anti*-amino acid derivatives *via* palladium-catalyzed tandem allylic amination/[2,3]-rearrangement, see: (a) A. Soheili and U. K. Tambar, *J. Am. Chem. Soc.*, 2011, **133**, 12956; (b) A. Nash, A. Soheili and U. K. Tambar, *Org. Lett.*, 2013, **15**, 4770; (c) A. Soheili and U. K. Tambar, *Org. Lett.*, 2013, **15**, 5138.
- 15 Even at –20 °C, the background reaction still occurred, and the desired racemic product was isolated in 40% yield with a slightly lower *anti* : *syn* ratio (1 : 1.5).
- 16 CCDC 1960932 (**4u**)†
- 17 (a) W. W. Luo, X. Yuan, L. L. Lin, P. F. Zhou, X. H. Liu and X. M. Feng, *Chem. Sci.*, 2016, **7**, 4736; (b) D. Zhang, L. L. Lin, J. Yang, X. H. Liu and X. M. Feng, *Angew. Chem., Int. Ed.*, 2018, **57**, 12323.

