


Chemical Science

Volume 11
Number 11
21 March 2020
Pages 2865–3112

rsc.li/chemical-science



ISSN 2041-6539

EDGE ARTICLE

Fu Peng, Bo Han *et al.*
Regiodivergent construction of medium-sized heterocycles
from vinyl ethylene carbonates and allylidenemalononitriles

Cite this: *Chem. Sci.*, 2020, 11, 2888

All publication charges for this article have been paid for by the Royal Society of Chemistry

Regiodivergent construction of medium-sized heterocycles from vinyl ethylene carbonates and allylidenemalononitriles†

Xiang Zhang,^{a,c,d} Xiang Li,^b Jun-Long Li,^b Qi-Wei Wang,^{c,d} Wen-Lin Zou,^d Yan-Qing Liu,^{b,d} Zhi-Qiang Jia,^d Fu Peng^{*a} and Bo Han^{*,b}

Medium-sized heterocycles exist in a broad spectrum of biologically active natural products and medicinally important synthetic compounds. The construction of medium-sized rings remains challenging, particularly the assembly of different ring sizes from the same type of substrate. Here we report palladium-catalyzed, regiodivergent [5 + 4] and [5 + 2] annulations of vinyl ethylene carbonates and allylidenemalononitriles. We describe the production of over 50 examples of nine- and seven-membered heterocycles in high isolated yields and excellent regioselectivities. We demonstrate the synthetic utility of this approach by converting a nine-membered ring product to an interesting polycyclic caged molecule *via* a [2 + 2] transannulation. Mechanistic studies suggest that the [5 + 2] annulation proceeds through palladium-catalyzed ring-opening/re-cyclization from the [5 + 4] adducts.

Received 17th December 2019

Accepted 9th February 2020

DOI: 10.1039/c9sc06377c

rsc.li/chemical-science

Introduction

Cyclic molecular frameworks have special importance in chemical research and industry.¹ Medium-sized rings (MSR, 7–11 members),² particularly hetero-rings, exist in a large number of biologically active natural products and medicinally important synthetic molecules³ (Fig. 1). However, MSRs are challenging to prepare because of their inherent entropic factors and transannular interactions. Most established methods to generate MSRs are based on a fixed reaction site and suitable only for rings of the same size;⁴ changing the size of the ring usually requires changing the substrate design.⁵ Such a substrate-controlled strategy can be quite costly and inefficient because of the need to prepare the necessary substrate variants and optimize them in the ring-forming reactions. It could be much more efficient to develop a way to generate medium-sized rings of various sizes from the same set of

substrates, simply by altering the reaction conditions. However, to our knowledge, controlling the regioselectivity of medium-sized ring cyclization is notoriously difficult and remains underdeveloped⁶ (Scheme 1a).

Vinyl ethylene carbonates (VECs) have recently emerged as versatile building blocks for various cyclizations, because of their inherent ability to undergo decarboxylation in the presence of a palladium catalyst to generate highly reactive




Fig. 1 Selected natural products and synthetic bioactive compounds containing medium-sized oxo-heterocycles.


^aWest China School of Pharmacy, Sichuan University, Chengdu 610041, China. E-mail: pengf@scu.edu.cn

^bState Key Laboratory of Southwestern Chinese Medicine Resources, School of Pharmacy, Chengdu University of Traditional Chinese Medicine, Chengdu 611137, China. E-mail: hanbo@cdutcm.edu.cn

^cChengdu Institute of Organic Chemistry, Chinese Academy of Sciences, Chengdu 610041, China

^dAntibiotics Research and Re-evaluation Key Laboratory of Sichuan Province, Sichuan Industrial Institute of Antibiotics, Chengdu University, Chengdu 610052, China

† Electronic supplementary information (ESI) available: Experimental procedures, characterization data for new compounds and crystallographic data in CIF or other electronic formats. CCDC 1943770–1943772. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc06377c




Scheme 1 Divergent construction of medium-sized rings.

zwitterionic π -allyl palladium intermediates.^{7,8} Recently, Zhao and co-workers disclosed that π -allyl palladium species can serve as 1,5-dipoles in a highly efficient [5 + 4] annulation with 1,3-azadienes to construct nine-membered hetero-rings.⁹ Since then, palladium-catalyzed [5 + *n*] annulations involving vinylidene carbonates have been described for generating various medium-sized heterocycles¹⁰ (Scheme 1b). However, rarely have vinylidene carbonates been used for divergent annulation,^{10h,11} and to our knowledge, they have never been applied to regioselective [5 + *n*] cyclization, which could generate multiple ring sizes.

Given our experience with the assembly of biologically interesting heterocycles by exploring novel catalytic reactions,¹² we aimed to develop a convenient strategy for ring size-divergent construction of medium-sized rings. We found that by using the versatile, electron-deficient diene substrate allylidene malononitriles,¹³ we could achieve smooth [5 + 4] annulation with vinylidene carbonates in MeCN in the presence of a palladium catalyst at room temperature, generating a nine-membered product. More importantly, we could completely shift the regioselectivity to [5 + 2] cyclization by changing the solvent to THF and increasing the reaction temperature, generating a seven-membered product. In both cases, the regioselectivity was nearly perfect (Scheme 1c). In addition, the nine-membered cyclic ether adducts were able to undergo intramolecular trans-annular [2 + 2] cycloaddition¹⁴ to build a structurally interesting caged polycycle.

Results and discussion


Our investigations began with a reaction between the easily accessible diene **1a** and vinylidene carbonate **2a**. Different solvents were evaluated in the presence of Pd(PPh₃)₄ at 20 °C, and MeCN afforded the [5 + 4] adduct **3a** with a high yield and regioselectivity, while other solvents provided a mixture of nine- and seven-membered products (Table 1, entries 1–5) or **3a** in low yield (entry 6). The reaction in THF gave the highest ratio of [5 + 2] product **4a**, which encouraged us to screen the reaction conditions further in order to switch the regioselectivity. With THF as the solvent, phosphine ligands **L1–L7** were screened, but all reacted inefficiently (entry 7). To our gratification, conducting the reaction at 40 °C improved the relative amount of seven-membered cyclic ether **4a**, and increasing the temperature to 60 °C afforded **4a** as a single regioisomer in high yield (entries 8 and 9). Further increasing the temperature maintained the high regioselectivity but lowered the yield slightly (entry 10). Using other solvents at 60 °C did not improve the results in terms of yield and regioselectivity (entries 11–16).¹⁵

Table 1 Optimization studies for the annulation of allylidene malononitril **1a** and VEC **2a**^a

Entry	Catalyst	Solvent	Temp. (°C)	Yield ^b (%)	3a : 4a ^c
1 ^d	Pd(PPh ₃) ₄	Toluene	20	72	3.5 : 1
2 ^e	Pd(PPh₃)₄	MeCN	20	96(90)	>20 : 1
3 ^d	Pd(PPh ₃) ₄	DCM	20	68	3.6 : 1
4 ^d	Pd(PPh ₃) ₄	CHCl ₃	20	85	2.6 : 1
5	Pd(PPh ₃) ₄	THF	20	85	1.4 : 1
6 ^e	Pd(PPh ₃) ₄	DMF	20	16	>20 : 1
7 ^f	Pd/L1–L7	THF	20	<5	—
8	Pd(PPh ₃) ₄	THF	40	98	1 : 4.6
9	Pd(PPh₃)₄	THF	60	91(84)	<1 : 20
10	Pd(PPh ₃) ₄	THF	80	89	<1 : 20
11	Pd(PPh ₃) ₄	1,4-Dioxane	60	83	16.0 : 1
12	Pd(PPh ₃) ₄	Toluene	60	88	1 : 1.3
13	Pd(PPh ₃) ₄	MeCN	60	87	8.6 : 1
14	Pd(PPh ₃) ₄	DMF	60	81	14.8 : 1
15	Pd(PPh ₃) ₄	DCM	60	76	1 : 1.4
16	Pd(PPh ₃) ₄	CHCl ₃	60	80	5.3 : 1

^a Unless noted otherwise, the reactions were carried out with **1a** (0.10 mmol), **2a** (0.15 mmol) and the Pd catalyst (5 mol%) in solvent (1 mL) for 12 h. ^b Yield was determined by ¹H-NMR analysis with CH₂Br₂ as the internal standard; the data in parentheses refer to isolated yields. ^c The ratio of **3a** : **4a** was determined by ¹H-NMR analysis of the crude reaction mixture. ^d For 48 h. ^e For 24 h. ^f The Pd/ligand complex was pre-prepared with Pd₂(dba)₃·CHCl₃ and a ligand in THF at rt for 1 h.






Scheme 2 Large-scale reactions of regiodivergent cyclizations and further synthetic applications.

selectride triggered reductive C–O bond cleavage that opened the nine-membered ring, offering linear 1,4-diene alcohol **6** in moderate yield. The product **4a** could undergo a retro-Knoevenagel reaction under aqueous basic conditions to release the malononitrile moiety and give the ketone-containing derivative **7** in 52% yield. It could also undergo sequential retro-Knoevenagel and retro-Claisen condensation in the presence of Et_3N , $i\text{PrOH}$ and water to afford product **8** in excellent yield. In addition, we extended this divergent cyclization strategy to a reaction between **1a** and vinyloxazolidinone **9**, assembling the nine- and seven-membered azacycles **10** and **11** in satisfying yields with excellent regioselectivities (Scheme 2c).

Unexpectedly, heating the [5 + 4] adduct **3a** without the Pd catalyst in toluene generated a cage-like molecule **12a** in high yield. The structure of **12a** was confirmed by X-ray diffraction analysis. We attribute the formation of this product to heat-induced isomerization of the styrene moiety from the *E*- to *Z*-configuration, followed by transannular [2 + 2] cycloaddition (for the preliminary mechanism investigation, see the ESI†). This reaction proved tolerant of various functional groups, allowing the rapid synthesis of caged compounds **12a–12j** (Scheme 3a). With a series of synthesized molecule fused pharmacologically privileged frameworks in hand and motivated by the pharmaceutical properties of nitrile,^{16a–c} oxygen heterocycles^{1f} and caged-skeletons,^{16d–h} we preliminarily evaluated their ability to inhibit the proliferation of a panel of cancer cell lines (Scheme 3b). In these experiments, the concentrations of tested compounds and paclitaxel (PTX) were 20 μM and 5 μM , respectively. Compounds **12c/j**, **12j**, **12a** and **12d** showed




Scheme 3 The transannular [2 + 2] cycloaddition of **3** (a) and heat map of the mean inhibitory ratio of compounds **12a–12j** against a panel of cancer cell lines (b).

promising cytotoxicity against A549, PC12, SH-SY5Y and A375 cells, respectively (for the details, see ESI, Table S3†).

In order to investigate the reaction mechanism, we performed several control experiments based on the reaction of allylidene malononitril **1a** and vinyethylene carbonate **2a**. Firstly, the reaction progress was monitored by NMR analysis. As shown in Scheme 4a, under the [5 + 4] annulation reaction conditions, the nine-membered product **3a** formed gradually, without concomitant emergence of the [5 + 2] seven-membered product **4a**. In contrast, in the reaction meant to produce **4a**, the starting material **1a** was rapidly consumed and **3a** was initially generated in high NMR yield, together with trace amounts of **4a**. Subsequently, the ratio of **3a/4a** slowly decreased until **4a** was obtained as the sole regioisomer (Scheme 4b). Follow-up experiments showed that in the presence of a palladium catalyst in THF at 60 °C, **3a** converted to **4a**, but not *vice versa* (Scheme 4c). These results suggest that the nine-membered **3a** undergoes palladium-catalyzed ring-opening/re-cyclization to produce **4a**. In addition, we found that using excess vinyethylene carbonate inhibited the transformation from **3a** into **4a** under heating conditions in THF (Scheme 4d), probably because the palladium catalyst prefers to coordinate with a higher concentration of vinyethylene carbonate which blocks the palladium activation of **3a**.¹⁷

These experimental results suggest the following mechanism to rationalize the regioselectivity of the [5 + 4] and [5 + 2] annulations (Fig. 2). The palladium-catalyzed decarboxylation of vinyethylene carbonate **2a** generates an ambiphilic π -allyl palladium intermediate **I**, which undergoes vinylogous Michael





Scheme 4 Control experiments. (a) Reaction progress was monitored in MeCN at 20 °C; (b) Reaction progress was monitored in THF at 60 °C; (c) Transformation from 3a to 4a; (d) Effect of the loading of VEC on the regioisomeric ratio.




Fig. 2 Proposed mechanism.

addition with allylidenemalononitrile **1a** to form intermediate **II**. At lower temperature and in MeCN solvent, the π -allylic anion is stabilized by dicyano electron-withdrawing groups, so the corresponding α terminal carbon attacks the electrophilic π -allyl palladium moiety to deliver **3a** in a kinetically controlled manner. At higher temperature and in THF solvent, the same

pathway generates **3a**, which can revert to intermediate **II** via palladium-catalyzed ring-opening, but en route it can undergo a different ring-closing reaction between an internal γ -carbon and the π -allyl palladium moiety, delivering **4a** in a thermodynamically controlled reaction.

Conclusions

In summary, we have developed a regiodivergent cyclization of vinyl ethylene carbonates and allylidenemalononitriles for the synthesis of medium-sized heterocycles. [5 + 4] annulation proceeds smoothly in MeCN at lower temperature, delivering nine-membered oxo-heterocycles in high yields. Changing the solvent to THF and raising the temperature completely reverse the regioselectivity of the ring-closing step, giving rise to [5 + 2] annulation that generates seven-membered heterocycles. In this way, our strategy allows the selective assembly of two heterocycle sizes from the same set of substrates through simple manipulation of reaction conditions. The nine-membered products efficiently undergo a transannular [2 + 2] cycloaddition to afford intriguing caged ring systems. Mechanistic studies suggest that [5 + 2] cyclization may occur via palladium-catalyzed ring-opening/cyclization from [5 + 4] adducts. Further biological studies of these novel cyclic molecules are currently underway in our laboratory, and the results will be reported in due course.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

Financial support from the NSFC (21871031, 21702021 and 81573588), Science & Technology Department of Sichuan Province (2017JQ0032, 2017JZYD0001, and 2017JY0323), “Thousand Talents Program” of Sichuan Province, “Chengdu Talents Program” and Start-up Fund of Chengdu University is gratefully acknowledged.

Notes and references

- (a) W. Carruthers, *Cycloaddition reactions in organic synthesis*, Pergamon, Oxford, 1990; (b) *Cycloaddition reactions in organic synthesis*, ed. S. Kobayashi and K. A. Jorgensen, Wiley-VCH, New York, 2002; (c) *Handbook of cyclization reactions*, ed. S.-M. Ma, Wiley-VCH, New York, 2010; (d) *Methods and applications of cycloaddition reactions in organic syntheses*, ed. N. Nishiwaki, John Wiley & Sons, New York, 2014; (e) E. Vitaku, D. T. Smith and J. T. Njardarson, *J. Med. Chem.*, 2014, **57**, 10257; (f) M. D. Delost, D. T. Smith, B. J. Anderson and J. T. Njardarson, *J. Med. Chem.*, 2018, **61**, 10996.
- For selected reviews on medium-sized rings, see: (a) L. Yet, *Chem. Rev.*, 2000, **100**, 2963; (b) A. S. Kleinke, D. Webb and T. F. Jamison, *Tetrahedron*, 2012, **68**, 6999; (c) M. E. Maier, *Angew. Chem., Int. Ed.*, 2000, **39**, 2073; (d) I. Shiina, *Chem.*



- Rev.*, 2007, **107**, 239; (e) J. R. Donald and W. P. Unsworth, *Chem.–Eur. J.*, 2017, **23**, 8780; (f) A. Hussain, S. K. Yousuf and D. Mukherjee, *RSC Adv.*, 2014, **4**, 43241.
- 3 (a) G. Guella, I. Mancini, G. Chiasera and F. Pietra, *Helv. Chim. Acta*, 1992, **75**, 310; (b) M. Yoshida, K. Nakatani and K. Shishido, *Tetrahedron*, 2009, **65**, 5702; (c) J. B. P. A. Wijnberg, A. van Veldhuizen, H. J. Swarts, J. C. Frankland and J. A. Field, *Tetrahedron Lett.*, 1999, **40**, 5767; (d) T. J. King, S. Imre, A. Öztunc and R. H. Thomson, *Tetrahedron Lett.*, 1979, **20**, 1453; (e) H. Niwa, K. Wakamatsu and K. Yamada, *Tetrahedron Lett.*, 1989, **30**, 4543; (f) B. F. Bowden, J. C. Coll and M. C. Dai, *Aust. J. Chem.*, 1989, **42**, 665; (g) M. Satake, M. Murata and T. Yasumoto, *Tetrahedron Lett.*, 1993, **34**, 1975.
- 4 For selected recent examples for the synthesis of medium-sized rings, see: (a) C. R. Kennedy, H. Zhong, R. L. Macaulay and P. J. Chirik, *J. Am. Chem. Soc.*, 2019, **141**, 8557; (b) C. Zhu, B. Yang, B. K. Mai, S. Palazzotto, Y. Qiu, A. Gudmundsson, A. Ricke, F. Himmo and J.-E. Bäckvall, *J. Am. Chem. Soc.*, 2018, **140**, 14324; (c) L. Zhang, Y. Wang, Z.-J. Yao, S. Wang and Z.-X. Yu, *J. Am. Chem. Soc.*, 2015, **137**, 13290; (d) X. Hong, M. C. Stevens, P. Liu, P. A. Wender and K. N. Houk, *J. Am. Chem. Soc.*, 2014, **136**, 17273; (e) S. Saito, K. Maeda, R. Yamasaki, T. Kitamura, M. Nakagawa, K. Kato, I. Azumaya and H. Masu, *Angew. Chem., Int. Ed.*, 2010, **49**, 1830.
- 5 For selected examples, see: (a) Y. A. Cheng, T. Chen, C. K. Tan, J. J. Heng and Y.-Y. Yeung, *J. Am. Chem. Soc.*, 2012, **134**, 16492; (b) Y. Hu and H. Huang, *Org. Lett.*, 2017, **19**, 5070; (c) F. Medina, C. Besnard and J. Lacour, *Org. Lett.*, 2014, **16**, 3232; (d) G. Prado, A. X. Veiga, F. Fernández-Nieto, M. R. Paleo and F. J. Sardina, *Org. Lett.*, 2015, **17**, 2054; (e) N. P. Tsvetkov, A. Bayir, S. Schneider and M. Brewer, *Org. Lett.*, 2012, **14**, 264; (f) Z. Wang, S. Chen, J. Ren and Z. Wang, *Org. Lett.*, 2015, **17**, 4184; (g) I. D. G. Watson, S. Ritter and F. D. Toste, *J. Am. Chem. Soc.*, 2009, **131**, 2056; (h) Z. Wu and J. Wang, *ACS Catal.*, 2017, **7**, 7647.
- 6 There is a report on the ligand controlled divergent synthesis of medium-sized rings; however, only one example achieved the regioselective switch. For details, see: M. M. Coulter, P. K. Dornan and V. M. Dong, *J. Am. Chem. Soc.*, 2019, **131**, 6932.
- 7 For selected recent reviews on VECs, see: (a) J. E. Gómez and A. W. Kleij, *Adv. Organomet. Chem.*, 2019, **71**, 175; (b) W. Guo, J. E. Gómez, À. Cristófol, J. Xie and A. W. Kleij, *Angew. Chem., Int. Ed.*, 2018, **57**, 13735; (c) A. Khan and Y. J. Zhang, *Synlett*, 2015, **26**, 853.
- 8 For selected examples on the transitional chemistry of VECs, see: (a) A. Khan, R. Zheng, Y. Kan, J. Ye, J. Xing and Y. J. Zhang, *Angew. Chem., Int. Ed.*, 2014, **53**, 6439; (b) A. Khan, L. Yang, J. Xu, L. Y. Jin and Y. J. Zhang, *Angew. Chem., Int. Ed.*, 2014, **53**, 11257; (c) A. Cai, W. Guo, L. Martínez-Rodríguez and A. W. Kleij, *J. Am. Chem. Soc.*, 2016, **138**, 14194; (d) A. Khan, S. Khan, I. Khan, C. Zhao, Y. Mao, Y. Chen and Y. J. Zhang, *J. Am. Chem. Soc.*, 2017, **139**, 10733; (e) W. Guo, L. Martínez-Rodríguez, R. Kuniyil, E. Martin, E. C. Escudero-Adán, F. Maseras and A. W. Kleij, *J. Am. Chem. Soc.*, 2016, **138**, 11970; (f) W. Guo, L. Martínez-Rodríguez, E. Martin, E. C. Escudero-Adán and A. W. Kleij, *Angew. Chem., Int. Ed.*, 2016, **55**, 11037; (g) W. Guo, R. Kuniyil, J. E. Gómez, F. Maseras and A. W. Kleij, *J. Am. Chem. Soc.*, 2018, **140**, 3981; (h) R. Zeng, J.-L. Li, X. Zhang, Y.-Q. Liu, Z.-Q. Jia, H.-J. Leng, Q.-W. Huang, Y. Liu and Q.-Z. Li, *ACS Catal.*, 2019, **9**, 8256; (i) Y. Liu, Q.-W. Huang, Q.-Z. Li, H.-J. Leng, Q.-S. Dai, R. Zeng, Y.-Q. Liu, X. Zhang, B. Han and J.-L. Li, *Org. Lett.*, 2019, **21**, 7478.
- 9 (a) L.-C. Yang, Z.-Q. Rong, Y.-N. Wang, Z. Y. Tan, M. Wang and Y. Zhao, *Angew. Chem., Int. Ed.*, 2017, **56**, 2927; (b) Z.-Q. Rong, L.-C. Yang, S. Liu, Z. Yu, Y.-N. Wang, Z. Y. Tan, R.-Z. Huang, Y. Lan and Y. Zhao, *J. Am. Chem. Soc.*, 2017, **139**, 15304.
- 10 For selected examples on the $[5 + n]$ annulation of VECs, see: (a) P. Das, S. Gondo, P. Nagender, H. Uno, E. Tokunaga and N. Shibata, *Chem. Sci.*, 2018, **9**, 3276; (b) X. Gao, M. Xia, C. Yuan, L. Zhou, W. Sun, C. Li, B. Wu, D. Zhu, C. Zhang, B. Zheng, D. Wang and H. Guo, *ACS Catal.*, 2019, **9**, 1645; (c) C. Yuan, Y. Wu, D. Wang, Z. Zhang, C. Wang, L. Zhou, C. Zhang, B. Song and H. Guo, *Adv. Synth. Catal.*, 2018, **360**, 652; (d) B. Niu, X.-Y. Wu, Y. Wei and M. Shi, *Org. Lett.*, 2019, **21**, 4859; (e) S. Singha, T. Patra, C. G. Daniliuc and F. Glorius, *J. Am. Chem. Soc.*, 2018, **140**, 3551; (f) Y. Wei, S. Liu, M.-M. Li, Y. Li, Y. Lan, L.-Q. Lu and W.-J. Xiao, *J. Am. Chem. Soc.*, 2019, **141**, 133; (g) H.-W. Zhao, J. Du, J.-M. Guo, N.-N. Feng, L.-R. Wang, W.-Q. Ding and X.-Q. Song, *Chem. Commun.*, 2018, **54**, 9178; (h) Y. Yang and W. Yang, *Chem. Commun.*, 2018, **54**, 12182. For Pd-catalyzed $[5 + n]$ annulations with vinyloxiranes, see: (i) Y. Wu, C. Yuan, C. Wang, B. Mao, H. Jia, X. Gao, J. Liao, F. Jiang, L. Zhou, Q. Wang and H. Guo, *Org. Lett.*, 2017, **19**, 6268; (j) J.-J. Feng and J. Zhang, *J. Am. Chem. Soc.*, 2011, **133**, 7304; (k) J.-J. Feng and J. Zhang, *ACS Catal.*, 2017, **7**, 1533.
- 11 For divergent annulations with VECs, see: (a) L.-C. Yang, Z. Y. Tan, Z.-Q. Rong, R. Liu, Y.-N. Wang and Y. Zhao, *Angew. Chem., Int. Ed.*, 2018, **57**, 7860; (b) Y. Xia, Q.-F. Bao, Y. Li, L.-J. Wang, B.-S. Zhang, H.-C. Liu and Y.-M. Liang, *Chem. Commun.*, 2019, **55**, 4675.
- 12 (a) Q. Li, L. Zhou, X.-D. Shen, K.-C. Yang, X. Zhang, Q.-S. Dai, H.-J. Leng, Q.-Z. Li and J.-L. Li, *Angew. Chem., Int. Ed.*, 2018, **57**, 1913; (b) M.-C. Yang, C. Peng, H. Huang, L. Yang, X.-H. He, W. Huang, H.-L. Cui, G. He and B. Han, *Org. Lett.*, 2017, **19**, 6752; (c) J.-L. Li, L. Fu, J. Wu, K.-C. Yang, Q.-Z. Li, X.-J. Gou, C. Peng, B. Han and X.-D. Shen, *Chem. Commun.*, 2017, **53**, 6875; (d) Q.-Z. Li, X. Zhang, R. Zeng, Q.-S. Dai, Y. Liu, X.-D. Shen, H.-J. Leng, K.-C. Yang and J.-L. Li, *Org. Lett.*, 2018, **20**, 3700; (e) K.-C. Yang, Q.-Z. Li, Y. Liu, Q.-Q. He, Y. Liu, H.-J. Leng, A.-Q. Jia, S. Ramachandran and J.-L. Li, *Org. Lett.*, 2018, **20**, 7518.
- 13 (a) X.-N. Zhang, G.-Q. Chen, X.-Y. Tang, Y. Wei and M. Shi, *Angew. Chem., Int. Ed.*, 2014, **53**, 10768; (b) L. Zhang, H. Lu, G.-Q. Xu, Z.-Y. Wang and P.-F. Xu, *J. Org. Chem.*, 2017, **82**, 5782; (c) X. Zhang, Q.-F. Huang, W.-L. Zou, Q.-Z. Li,



X. Feng, Z.-Q. Jia, Y. Liu, J.-L. Li and Q.-W. Wang, *Org. Chem. Front.*, 2019, **6**, 3321.

14 For selected reviews on transannular reactions, see: (a) E. Reyes, U. Uria, L. Carrillo and J. L. Vicario, *Tetrahedron*, 2014, **70**, 9461; (b) A. Rizzo and S. R. Harutyunyan, *Org. Biomol. Chem.*, 2014, **12**, 6570; (c) S. Handa and G. Pattenden, *Contemp. Org. Synth.*, 1997, **4**, 196. For a recent example, see: (d) R. Mato, R. Manzano, E. Reyes, L. Carrillo, U. Uria and J. L. Vicario, *J. Am. Chem. Soc.*, 2019, **141**, 9495.

15 For more studies on condition screening, see the ESI.†

16 For selected reviews and examples, see: (a) F. F. Fleming, L. Yao, P. C. Ravikumar, L. Funk and B. C. Shook, *J. Med. Chem.*, 2010, **53**, 7902; (b) B. A. Klein, I. M. Robertson,

B. Reiz, T. Kampourakis, L. Li and B. D. Sykes, *ACS Med. Chem. Lett.*, 2019, **10**, 1007; (c) M. A. Cinelli, H. Li, G. Chreifi, T. L. Poulos and R. B. Silverman, *J. Med. Chem.*, 2017, **60**, 3958; (d) Q.-B. Han and H.-X. Xu, *Curr. Med. Chem.*, 2019, **16**, 3775; (e) Z. Zheng, M. Wu, J. Zhang, W. Fu, N. Xu, Y. Lao, L. Lin and H. Xu, *Frontiers in Oncology*, 2019, **9**, 654; (f) N. Anantachoke, P. Tuchinda, C. Kuhakarn, M. Pohmakotr and V. Reutrakul, *Pharm. Biol.*, 2012, **50**, 78; (g) S. Alam and F. Khan, *Sci. Rep.*, 2018, **8**, 5524; (h) C. Bao, M. Jin, B. Li, Y. Xu, J. Jin and L. Zhu, *Org. Biomol. Chem.*, 2012, **10**, 5238.

17 The regioselectivity of the annulation can be influenced by multiple factors, such as the solvent, temperature, the loading of ligand, *etc.* (for more details, see the ESI†).

