

Cite this: *Chem. Sci.*, 2020, 11, 5969

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 9th January 2020

Accepted 24th May 2020

DOI: 10.1039/d0sc00142b

rsc.li/chemical-science

Enantio- and diastereoselective diarylmethylation of 1,3-dicarbonyl compounds†


Xin Li, Songtao He and Qiuling Song *

A Cu-catalyzed enantio- and diastereoselective diarylmethylation of 1,3-dicarbonyl compounds is described. It is a successful example of constructing all-carbon quaternary stereocenters from 3° C–H nucleophiles, a challenging topic in synthetic chemistry. In the present work, two contiguous stereocenters are constructed with high levels of stereoselectivity and atom economy. The broad scope of 1,3-dicarbonyl nucleophiles and the tolerance of a wide range of functional groups make this protocol of great importance in the synthesis of chiral diarylmethane compounds.

Chiral diarylmethanes are ubiquitous in numerous natural products, medicinal molecules and bioactive compounds.¹ As a consequence, many elegant strategies have been developed to construct those structures. Among them, the asymmetric addition of an appropriate nucleophile to an aryl substituted quinone methide is considered as a direct and efficient access to chiral diarylmethanes.² During the past decade, with the development of many new kinds of nucleophiles and new modes of asymmetric catalytic systems, a variety of C–C³ and C–X⁴ bonds have been forged to furnish the target chiral diarylmethanes. Among those strategies developed, employing C–H nucleophiles is regarded as an ideal method because 100% atom economy can be achieved in that transformation. For instance, in 2013, Fan reported a beautiful work, in which malonic esters incorporated with *para*-quinone methides (*p*-QMs) enantioselectively under a chiral phase-transfer catalytic system (Scheme 1, route A).^{3a} Soon after, Jørgensen demonstrated a new organocatalytic concept to construct two contiguous stereocenters enantio- and diastereoselectively using their iconic chiral secondary amine catalysts (Scheme 1, route B).^{3b} Although great achievements have been made, the next progress in this field, *i.e.* employing 3° C–H nucleophiles to construct a more challenging all-carbon quaternary stereocenter⁵ was proven difficult, which may be attributed to the increased steric hindrance and the decreased acidity of the 3° C–H nucleophiles. To date, successful nucleophiles were restricted to oxindoles.⁶ Therefore, an efficient protocol which highly stereoselectively affords the two stereocenters with a broad 3° C–H nucleophiles scope is demanded. In this communication, we address this problem by employing a series of 1,3-dicarbonyl compounds

under a copper-chiral bisoxazoline catalytic system (Scheme 1, route C).

We commenced our investigation by using *p*-QM **1a** and 1,3-dicarbonyl compound **2a** as model substrates. After preliminary screening, a combination of 10 mol% of Cu(OTf)₂ and 12 mol% of a 1-amino-2-indanol derived bisoxazoline ligand **L1** catalyzed the reaction efficiently, affording the desired product with 98% isolated yield, 4.6 : 1 dr and 78% ee at 0 °C (Table 1, entry 1).⁷ Based on that result, a structural modification of the ligands family was carried out to improve the stereoselectivity of the reaction (entries 2–16). Gratifyingly, among a small library of 15 synthesized ligands, a 3,5-di-*tert*-butyl-benzyl substituted ligand **L8** gave an excellent outcome (98% isolated yield, 11.5 : 1 dr and 94% ee, entry 9). Meanwhile, those data showed that steric factor plays a crucial role in the performance of the ligands. For example, increasing the steric hindrance on the benzene ring of




Scheme 1 Enantioselective diarylmethylation of 1,3-dicarbonyl compounds.


Institute of Next Generation Matter Transformation, College of Materials Science & Engineering, Huaqiao University, 668 Jimei Blvd, Xiamen, Fujian, China. E-mail: qsong@hqu.edu.cn

† Electronic supplementary information (ESI) available. CCDC 1958259, 1963665 and 1963687. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d0sc00142b



Table 1 Screening of ligands^a



Entry	Ligand	T [°C]	Yield ^b [%]	dr ^c	ee ^d [%]
1	L1	0	98	4.6 : 1	78
2	L1	-10	98	4.7 : 1	81
3	L2	-10	92	4.4 : 1	83
4	L3	-10	90	4.0 : 1	79
5	L4	-10	37	4.6 : 1	75
6	L5	-10	24	3.8 : 1	73
7	L6	-10	96	4.2 : 1	83
8	L7	-10	95	3.6 : 1	77
9	L8	-10	98	11.5 : 1	94
10	L9	-10	98	6.7 : 1	86
11	L10	-10	69	5.7 : 1	88
12	L11	-10	97	5.3 : 1	85
13	L12	-10	48	3.4 : 1	83
14	L13	-10	40	4.3 : 1	74
15	L14	-10	56	4.2 : 1	70
16	L15	-10	25	3.6 : 1	90
17	L8	-15	98	14.6 : 1	98





^a Reaction conditions: *p*-quinone methide (0.05 mmol), ethyl 2-oxocyclopentanecarboxylate (0.075 mmol), Cu(OTf)₂ (10 mol%), ligand (12 mol%), THF (1.0 mL), 18 h. ^b Yield of the isolated product. ^c Determined by ¹H-NMR of the crude reaction mixture. ^d Determined by HPLC analysis and referred to the major diastereoisomer.

indanol resulted in a dramatic decrease on yield, and slightly decreases on dr and ee (entry 2 vs. entry 5). Besides, ligands bearing 4-substituted benzyl groups all gave much lower yields and poorer diastereoselectivities than their 3,5-disubstituted counterparts (entries 9–11 vs. entries 13–15). Further optimization showed that a higher stereoselectivity can be achieved when the reaction proceeded at -15 °C (entry 17).⁸

After the optimization works, a series of 1,3-dicarbonyl compounds were submitted to the standard reaction conditions to test the generalizability of the reaction toward C–H nucleophiles (Table 2). For 2 °C–H nucleophiles, ethyl 3-oxo-3-phenylpropanoate derivatives all gave good results with respect to yields, dr and ee's (3b–m). The electronic factor of the benzene rings has little influence on the reactivity and almost no influence on the stereoselectivity. For example, substrates bearing electron-withdrawing substituents on the benzene rings reacted more slowly under the same condition, thus

Table 2 Scope of 1,3-dicarbonyl nucleophiles^{a,b}


 3b, R = H, -15 °C, 99%, 10.0:1 dr, > 99% ee 3c, R = 4-F, -15 °C, 79%, > 20:1 dr, > 99% ee 3d, R = 4-Cl, -15 °C, 82%, 11.5:1 dr, > 99% ee 3e, R = 4-Br, -15 °C, 97%, 20.0:1 dr, > 99% ee 3f, R = 4-CF ₃ , -15 °C, 91%, 17.0:1 dr, 99% ee 3g, R = 4-NO ₂ , -15 °C, 74%, > 20:1 dr, 98% ee	 3h, R = 4-Me, -15 °C, 99%, 6.5:1 dr, 99% ee 3i, R = 4-OMe, -15 °C, 98%, 8.0:1 dr, 96% ee 3j, R = 3-Me, -15 °C, 99%, 8.7:1 dr, 97% ee 3k, R = 3-Cl, -15 °C, 74%, 7.8:1 dr, 96% ee 3l, R = 3-Br, 28 °C, 92%, 3.4:1 dr, 64% ee 3m, R = 2-Me, 28 °C, 99%, 9.0:1 dr, 94% ee	3n, R = Me, -15 °C, 99%, 5.4:1 dr, 88% ee 3o, R = Et, -15 °C, 89%, 5.0:1 dr, 73% ee 3p, R = ⁱ -Pr, 10 °C, 80%, 2.9:1 dr, 69% ee 3q, -15 °C, 99%, 1.8:1 dr, 83% ee (83% ee ^[c]) 3r, -15 °C, 86%, > 20:1 dr, 95% ee 3s, -15 °C, 93%, > 20:1 dr, 99% ee 3t, -15 °C, 99%, 2.3:1 dr, 98% ee 3u, -15 °C, 72%, 7.8:1 dr, 93% ee 3v, -15 °C, 99%, 11.1:1 dr, 91% ee 3w, 20 °C, 78%, 3.6:1 dr, 70% ee 3x, -15 °C, 95%, > 20:1 dr, > 99% ee 3y, -15 °C, 87%, > 20:1 dr, 94% ee 3z, -15 °C, 76%, > 20:1 dr, 96% ee 3aa, 10 °C, 75%, 1.2:1 dr, 65% ee (65% ee ^[c]) 3ab, 10 °C, 76%, 1.6:1 dr, 49% ee (30% ee ^[c])
---	--	--

^a Reaction conditions: *p*-quinone methide (0.05 mmol), 1,3-dicarbonyl compound (0.075 mmol), Cu(OTf)₂ (10 mol%), ligand (12 mol%), THF (1.0 mL), 18 h. ^b Isolated yield. ^c ee's of minor diastereoisomers.

resulting in lower conversions and yields (3c–g, 3k). Nevertheless, both electron-donating and electron-withdrawing substituents gave excellent stereoselectivities (up to >20 : 1 dr, up to >99% ee). In contrast, steric factor plays an important role in the reactivity and stereoselectivity of the substrates. For instance, *m*-bromo substituted substrate (3l) required a higher reaction temperature (ambient temperature)⁹ to afford the product but with a poorer dr and ee than those bearing smaller substituents such as methyl (3j) or chlorine atom (3k). *O*-Methyl (3m) also required a higher reaction temperature than *m*- and *p*-methyl (3h and 3j). The size of either the groups connected to the carbonyl or the ester groups determines the outcomes of the reactions. For example, with the increase of the size of ester groups, the reactivity and stereoselectivity of the nucleophiles both decreased (3n–p). Although a smaller cyclopropyl substituted ethyl acetoacetate gave poor diastereoselectivity and moderate enantioselectivity (3q), larger cyclopentyl and cyclohexyl substituents afforded the target products in good yields and excellent stereoselectivities (3r and 3s). Heterocyclic substituents are compatible with the reaction (3t and 3u).

More importantly, a diverse variety of 3° C–H nucleophiles exhibited equal reactivities and stereoselectivities. Substrates that differed in aromatic, aliphatic or acyclic carbon skeletons, both esters and lactones, all constructed the desired all-carbon quaternary stereocenters in high efficiencies (3v–z). Interestingly, the existence of an ester subunit in the substrates was



C-3 position are important as well. For methyl, the smallest substituent tested, it brings a poor discrimination between the Si- and Re-faces of the *p*-QM planes, thus resulting in the formation of unselective products (Scheme 3B, **6a**). When a 3-methyl-3'-*tert*-butyl substrate was employed, the stereoselectivity improved but was still moderate (**6b**). Isopropyl, which has a smaller size than *tert*-butyl, afforded the product slightly less stereoselectively than the *tert*-butyl did (**6c**). Interestingly, a bigger TMS group exhibited modestly better steric control (**6d**). Unlike alkyl substituents, C-3 aryl rings substituents don't obey the above-mentioned "size law", showing a deleterious effect on the reactivity and stereoselectivity (**6e**). On the basis of all of the experimental results, the high stereoselectivity of this reaction may owe to the combined interaction among the chiral catalyst, *p*-QMs, and the nucleophiles. Thus, a possible catalytic cycle and a plausible transition state which determines the absolute configuration of the products are proposed (Scheme 3C). Once the chiral Lewis acid catalyst **I** was generated from Cu(OTf)₂ and the box ligand, the subsequent ligand exchange and deprotonation of 1,3-dicarbonyl nucleophiles formed an intermediate **II**, which then underwent an asymmetric 1,6-addition with the *p*-QMs to afford the final chiral product with high levels of enantioselectivity and diastereoselectivity. The presumed transition state rationalizes the observed absolute stereochemical outcome. In this model, the chiral catalyst determines a Si-face attack on the *p*-QM planes, the ester groups of nucleophiles are opposite the aryls of *p*-QMs to minimize the steric hindrance or provide additional π -stacking interactions (for **3b–m**, **3x**).

The reaction can be carried out in gram-scale using 3 mol% catalyst loading without the destruction on yields and ee's, only at the expense of prolonged reaction times (Scheme 4A). Although these synthesized products are unstable under the acidic AlCl₃-induced de-*tert*-butylation process,¹² after reducing the carbonyls to either hydroxyls (**3a**) or methylene groups

(**3b**),¹³ the *tert*-butyls on the aryl rings can be readily removed by AlCl₃ without any loss of the optical purities (Scheme 4B).

Conclusions

In conclusion, we have developed a simple but efficient protocol to construct two contiguous stereocenters including one all-carbon quaternary stereocenter from a broad scope of *p*-QMs and 1,3-dicarbonyl compounds with high atom economy. Good yields and excellent stereoselectivities are achieved in most cases. The application of the copper-chiral bisoxazoline catalytic system to other asymmetric synthesis is under investigated in our group.

Conflicts of interest


There are no conflicts to declare.

Acknowledgements

We gratefully acknowledge the finance support from the National Natural Science Foundation (21772046 and 2193103).

Notes and references

- For reviews, see: (a) T. C. Nugent, *Chiral Amine Synthesis: Methods, Developments and Applications*, Wiley-VCH, Weinheim, Germany, 2010; (b) F. Schmidt, R. T. Stemmler, R. Rudolph and C. Bolm, *Chem. Soc. Rev.*, 2006, **35**, 454–470; (c) D. Ameen and T. J. Snape, *MedChemComm*, 2013, **4**, 893–907.
- For selected reviews on quinone methide, see: (a) *Quinone Methides*, ed. S. E. Rokita, Wiley, Hoboken, 2009; (b) T. P. Pathak and M. S. Sigman, *J. Org. Chem.*, 2011, **76**, 9210–9215; (c) N. J. Willis and C. D. Bray, *Chem.–Eur. J.*, 2012, **18**, 9160–9173; (d) Z. Wang and J. Sun, *Synthesis*, 2015, **47**, 3629–3644; (e) L. Caruana, M. Fochi and L. Bernardi, *Molecules*, 2015, **20**, 11733–11764; (f) A. Parra and M. Tortosa, *ChemCatChem*, 2015, **7**, 1524–1526; (g) W. Li, X. Xu, P. Zhang and P. Li, *Chem.–Asian J.*, 2018, **13**, 2350–2359.
- (a) W.-D. Chu, L.-F. Zhang, X. Bao, X.-H. Zhao, C. Zeng, J.-Y. Du, G.-B. Zhang, F.-X. Wang, X.-Y. Ma and C.-A. Fan, *Angew. Chem., Int. Ed.*, 2013, **52**, 9229–9233; (b) L. Caruana, F. Kniep, T. K. Johansen, P. H. Poulsen and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2014, **136**, 15929–15932; (c) K. Zhao, Y. Zhi, T. Shu, A. Valkonen, K. Rissanen and D. Enders, *Angew. Chem., Int. Ed.*, 2016, **55**, 12104–12108; (d) S. Li, Y. Liu, B. Huang, T. Zhou, H. Tao, Y. Xiao, Y. Liu and J. Zhang, *ACS Catal.*, 2017, **7**, 2805–2809; (e) F.-S. He, J.-H. Jin, Z.-T. Yang, X. Yu, J. S. Fossey and W.-P. Deng, *ACS Catal.*, 2016, **6**, 652–656; (f) L. Zhang, H. Yuan, W. Lin, Y. Cheng, P. Li and W. Li, *Org. Lett.*, 2018, **20**, 4970–4974; (g) Z. Sun, B. Sun, N. Kumagai and M. Shibasaki, *Org. Lett.*, 2018, **20**, 3070–3073; (h) X. Li, X. Xu, W. Wei, A. Lin and H. Yao, *Org. Lett.*, 2016, **18**, 428–431; (i) X.-Z. Zhang, K.-J. Gan, X.-X. Liu, Y.-H. Deng, F.-X. Wang, K.-J. Yu,



Scheme 4 Gram-scale synthesis and de-*tert*-butylation of the products.



- J. Zhang and C.-A. Fan, *Org. Lett.*, 2017, **19**, 3207–3210; (f) X.-Z. Zhang, Y.-H. Deng, X. Yan, K.-Y. Yu, F.-X. Wang, X.-Y. Ma and C.-A. Fan, *J. Org. Chem.*, 2016, **81**, 5655–5662; (k) L. Ge, X. Lu, C. Cheng, J. Chen, W. Cao, X. Wu and G. Zhao, *J. Org. Chem.*, 2016, **81**, 9315–9325; (l) T.-C. Kang, L.-P. Wu, Q.-W. Yu and X.-Y. Wu, *Chem.–Eur. J.*, 2017, **23**, 6509–6513; (m) Q. Liu, S. Li, X.-Y. Chen, K. Rissanen and D. Enders, *Org. Lett.*, 2018, **20**, 3622–3626; (n) W. Li, X. Xu, Y. Liu, H. Gao, Y. Cheng and P. Li, *Org. Lett.*, 2018, **20**, 1142–1145; (o) K. J. Romero, M. S. Galliher, M. A. R. Raycroft, J.-P. R. Chauvin, I. Bosque, D. A. Pratt and C. R. J. Stephenson, *Angew. Chem., Int. Ed.*, 2018, **57**, 17125–17129; (p) G.-B. Huang, W.-H. Huang, J. Guo, D.-L. Xu, X.-C. Qu, P.-H. Zhai, X.-H. Zheng, J. Weng and G. Lu, *Adv. Synth. Catal.*, 2019, **361**, 1241–1246; (q) Y. Wang, K. Wang, W. Cao, X. Liu and X. Feng, *Org. Lett.*, 2019, **21**, 6063–6067; (r) D. Wang, Z.-F. Song, W.-J. Wang and T. Xu, *Org. Lett.*, 2019, **21**, 3963–3967; (s) J.-R. Wang, X.-L. Jiang, Q.-Q. Hang, S. Zhang, G.-J. Mei and F. Shi, *J. Org. Chem.*, 2019, **84**, 7829–7839.
- 4 For C–B bond formation, see: (a) Y. Lou, P. Cao, T. Jia, Y. Zhang, M. Wang and J. Liao, *Angew. Chem., Int. Ed.*, 2015, **54**, 12134–12138; (b) C. Jarava-Barrera, A. Parra, A. López, F. Cruz-Acosta, D. Collado-Sanz, D. J. Cárdenas and M. Tortosa, *ACS Catal.*, 2016, **6**, 442–446; For C–S bond formation, see: (c) N. Dong, Z.-P. Zhang, X.-S. Xue, X. Li and J.-P. Cheng, *Angew. Chem., Int. Ed.*, 2016, **55**, 1460–1464; For C–N bond formation, see: (d) F. Jiang, F.-R. Yuan, L.-W. Jin, G.-J. Mei and F. Shi, *ACS Catal.*, 2018, **8**, 10234–10240.
- 5 For selected reviews on all-carbon quaternary stereocenter, see: (a) E. J. Corey and A. Guzman-Perez, *Angew. Chem., Int. Ed.*, 1998, **37**, 388–401; (b) K. W. Quasdorf and L. E. Overman, *Nature*, 2014, **516**, 181–191; (c) M. Büschleb, S. Dorich, S. Hanessian, D. Tao, K. B. Schenthal and L. E. Overman, *Angew. Chem., Int. Ed.*, 2016, **55**, 4156–4186; (d) B. M. Trost and C. Jiang, *Synthesis*, 2006, 369–396; (e) Y. Liu, S.-J. Han, W.-B. Liu and B. M. Stoltz, *Acc. Chem. Res.*, 2015, **48**, 740–751; (f) J. Feng, M. Holmes and M. J. Krische, *Chem. Rev.*, 2017, **117**, 12564–12580; (g) B. Wang and Y.-Q. Tu, *Acc. Chem. Res.*, 2011, **44**, 1207–1222; (h) R. Long, J. Huang, J. Gong and Z. Yang, *Nat. Prod. Rep.*, 2015, **32**, 1584–1601; (i) P. W. Xu, J.-S. Yu, C. Chen, Z.-Y. Cao, F. Zhou and J. Zhou, *ACS Catal.*, 2019, **9**, 1820–1882; (j) X.-P. Zeng, Z.-Y. Cao, Y.-H. Wang, F. Zhou and J. Zhou, *Chem. Rev.*, 2016, **116**, 7330–7396; (k) L. M. Repka and S. E. Reisman, *J. Org. Chem.*, 2013, **78**, 12314–12320.
- 6 (a) K. Zhao, Y. Zhi, A. Wang and D. Enders, *ACS Catal.*, 2016, **6**, 657–660; (b) Y.-H. Deng, X.-Z. Zhang, K.-Y. Yu, X. Yan, J.-Y. Du, H. Huang and C.-A. Fan, *Chem. Commun.*, 2016, **52**, 4183–4186.
- 7 See ESI† for more details.
- 8 A dramatically low conversion was obtained when the reaction conducted at $-20\text{ }^{\circ}\text{C}$. See ESI† for details.
- 9 During this work, the average ambient temperature in our lab was $28\text{ }^{\circ}\text{C}$.
- 10 *p*-QM substrates bearing strong coordinated quinolone and pyridine rings gave no conversion under the standard reaction conditions.
- 11 See ESI† for more details.
- 12 These products are very sensitive toward either strong acidic or basic conditions, and decompose into the two starting materials very quickly.
- 13 S. A. Saleh and H. I. Tashtoush, *Tetrahedron*, 1998, **54**, 14157–14177.

