Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2020, 11, 4786

dll publication charges for this article have been paid for by the Royal Society of Chemistry

Received 6th March 2020 Accepted 22nd April 2020

DOI: 10.1039/d0sc01352h

rsc.li/chemical-science

Exploiting the radical reactivity of diazaphosphinanes in hydrodehalogenations and cascade cyclizations†

Jingjing Zhang, a Jin-Dong Yang (1) ** and Jin-Pei Cheng (1) **ab

The remarkable reducibility of diazaphosphinanes has been extensively applied in various hydrogenations, based on and yet limited by their well-known hydridic reactivity. Here we exploited their unprecedented radical reactivity to implement hydrodehalogenations and cascade cyclizations originally inaccessible by hydride transfer. These reactions feature a broad substrate scope, high efficiency and simplicity of manipulation. Mechanistic studies suggested a radical chain process in which a phosphinyl radical is generated in a catalytic cycle *via* hydrogen-atom transfer from diazaphosphinanes. The radical reactivity of diazaphosphinanes disclosed here differs from their well-established hydridic reactivity, and hence, opens a new avenue for diazaphosphinane applications in organic syntheses.

Introduction

As is known, in polar hydride transfer (HT), an adequate thermodynamic compensation is needed to overcome its high kinetic barrier.¹ Hence, direct HT usually requires a significant local accumulation of positive charge on hydride acceptors, which are commonly cationic species or some highly polarized compounds. This poses a very challenging, if not insuperable, barrier for reducing less polar substrates *via* HT. Consequently, hydrodehalogenation of organic halides with a hydride donor is barely achieved *via* nucleophilic aromatic substitution.² Compared with polar HT, the radical pathway displays a kinetic superiority because of its considerably lower intrinsic barrier.³ In this regard, hydrogen-atom transfer (HAT) may provide an alternative to enable reduction with hydride donors, which is unlikely to occur through direct HT.

N-heterocyclic phosphines have recently attracted particular attention owing to their superior hydricity (Scheme 1a).⁴ Their hydridic reactivity has been extensively exploited in catalytic or stoichiometric reduction of various unsaturated compounds, such as imines,⁵ aldehydes and ketones,⁶ polar olefins⁷ and pyridines⁸ (Scheme 1b). A kinetic scale with respect to the HT tendency from N-heterocyclic phosphines was also reported.⁹ In contrast to the well-studied hydridic reactivity, understanding

Based on the hydrogen-atom donating ability of diazaphosphinanes (DAPs) derived recently, ¹⁰ we envisioned that trivalent phosphinyl radicals, which could be generated *via* HAT from DAPs, may allow a chance for radical reduction, and hence, renders a quite different reactivity and selectivity pattern from that by direct HT of DAPs. Aiming to verify this assumption, we have examined here the hydrodehalogenation of

a) Representative N-heterocyclic phosphines

b) Hydridic reactivity

Scheme 1 N-heterocyclic phosphines and their applications. (a) Representative N-heterocyclic phosphines. (b) Applications of N-heterocyclic phosphines in hydridic reduction.

tsinghua.edu.cn

of their reducibility in hydrogen-atom or electron transfer pathways remains strikingly underdeveloped.

[&]quot;Center of Basic Molecular Science, Department of Chemistry, Tsinghua University,
Beijing 100084, China. E-mail: jdyang@mail.tsinghua.edu.cn; jinpei_cheng@mail.

^bState Key Laboratory of Elemento-Organic Chemistry, College of Chemistry, Nankai University, Tianjin 300071, China

 $[\]dagger$ Electronic supplementary information (ESI) available: Details of condition optimization, product characterization and DFT calculations. See DOI: 10.1039/d0sc01352h

Edge Article Chemical Science

organic halides under properly designed conditions in favor of radical initiation. To our delight, the anticipated radical hydrodehalogenation of organohalides was indeed realized with high efficiency and wide substrate coverage, and therefore, could be a superior alternative to substitute some of the traditional approaches (e.g., halogen-metal exchange, 11 transition metal catalysis,12 nucleophilic aromatic substitution2 and radical dehalogenation¹³) where limitation exists for removing halogens. Here. used 1,3-di-tert-butyl-1,3,2we diazaphosphinane (1a, in Scheme 2a) as a potent hydrogenatom donor to implement efficient radical dehalogenation as well as cascade cyclization of organic halides, which were found to be inert when reduced by the same diazaphosphinane via direct HT.

Scheme 2b shows our design, which was proposed based on the following analysis. Comparing the P-H bond dissociation free energy (BDFE) (~78 kcal mol⁻¹)¹⁰ of 1a with that of the isobutyronitrile α -C-H bond (\sim 85 kcal mol⁻¹), ¹⁴ we can expect a feasible HAT from 1a to the isobutyronitrile radical, generated from homolysis of azobis(isobutyronitrile) (AIBN), to give the crucial phosphinyl radical 1a-[P] (step a). The in situ formed 1a-[P] with an oxidation potential (E_{ox}) of -2.39 V (vs. ferrocene in acetonitrile, see the ESI† for details) is among the most potent neutral electron donors ever reported.15 It could undergo reversible electron transfer (ET, step b) with bromobenzene¹⁶ to give the reactive bromobenzene radical anion and stable phosphenium cation (or alternatively, abstracts a bromine atom (step b') from bromobenzene). Then, subsequent spontaneous C-Br bond scission (step c) produces the phenyl radical (C-H BDFE of benzene: ~105 kcal mol⁻¹), ¹⁴ which readily captures a hydrogen-atom from 1a to regenerate 1a-[P]' (step d) and simultaneously initiate the next cycle.

Results and discussion

To evaluate the feasibility of the present design, we chose the bromobenzene 2a as the test substrate (Table 1). As seen, treatment of 2a with 1.2 equiv. of 1a and 10 mol% AIBN in

b) Our proposed catalytic cycle

Scheme 2 Our work. (a) Radical hydrodehalogenation, and (b) catalytic cycle proposed in this work.

Table 1 Condition optimization for hydrodehalogenation of bromobenzene

Entry		Conditio	n ^a			Yield ^b
1	Standard condition					90%
2	5 mol% AIBN					78%
3	1b as reductant					<10%
4	1c as reductant					<5%
5	1d as reductant					<5%
6	C as reductant					77%
7	No AIBN					<5%
8	No heat					<5%
	P-H reagents 1:					
	PH N 'tBu	PH N _{iPr}	^t Bu P H ^t Bu	Ph PH Ph	Mes N PH N Mes	
	1a	1b	1c	1d	С	

 $[^]a$ Reactions were conducted using 0.10 mmol of ${\bf 2a}$ in 0.5 mL toluene. b $^1{\rm H}$ NMR yields using 1,3,5-trimethoxybenzene as the internal standard.

toluene harvested the product benzene 3a in 90% yield (entry 1 in Table 1). Reducing the amount of AIBN to 5 mol% caused an inferior result (78%, entry 2). Further evaluations were performed by varying P-H reductants. Replacement of 1a with structurally similar 1b resulted in a much lower yield (<10%, entry 3). This is primarily because the poor reducing power of the **1b**-derived phosphinyl radical ($E_{ox} = -1.94$ V, Fig. S1†) makes electron transfer to bromobenzene sluggish. The same reason could be applied to rationalize the poor results of 1c and 1d systems (<5%, entry 4 and 5). Nevertheless, employment of the stronger reducing reagent C only gave a moderate yield (77%, entry 6), along with some unidentified byproducts. This is probably due to the phosphinyl radical derived from C being too reactive to allow a sufficient ET (to bromobenzene) to proceed before it is guenched by other components in the system. In the control reactions without either the initiator AIBN or heating, trace amounts of products were obtained (<5%, entries 7 and 8), indicating that nucleophilic aromatic substitution may exist as a background reaction. Eventually, 1.2 equiv. of 1a and 10 mol% AIBN in toluene solvent were used as the standard conditions.


With the optimized conditions derived, we next evaluated the substrate scope, which covered both aromatic and aliphatic bromides and chlorides (Scheme 3; see the ESI† for details). As shown in Scheme 3, the reduction of 2-methylthiobromobenzene produced arene 3b in a quantitative yield (98%). 3-Phenyl (2c) and 3,5-di-*tert*-butyl (2d) analogs provided good results (83% and 73%). Impressively, the sterically hindered 2,4,6-trimethyl-bromobenzene is also a compatible substrate, affording 3e in 93% yield. The present reaction worked well with the labile acetal moiety and furnished 3f in

Scheme 3 Substrate scope of hydrodebromination/dechlorination. The reactions were conducted using 0.10 mmol 2 or 2′ in 0.5 mL toluene and isolated yields were given unless otherwise specified. [a] ^1H NMR yields using 1,3,5-trimethoxybenzene as the internal standard. [b] Using toluene- d_8 as solvent for these low boiling products. [c] Reaction time: 12 hours.

a high yield (88%). Notably, the 2-carbonyl analog can be quantitatively reduced to hydrodehalogenated 3g (99%). Direct hydride transfer to the electrophilic carbonyl group, as previously reported,6 can be completely inhibited. As expected, a moderate yield (60%) was obtained for electron-rich 4methoxy-bromobenzene (2h) because of its low reduction potential. Furthermore, hydrodebromination of substituted bromonaphthalenes (2i-k) proceeded smoothly and offered moderate to excellent yields (65-96%). Other condensed cyclic bromides (21 and 2m) were also viable substrates which gave corresponding products quantitatively. Besides, electron-rich 5bromoindole could react with 1a/AIBN and 3n was produced in a moderate yield (70%) after prolonging the reaction time to 12 hours. Other heterocyclic substrates can also be reduced to yield corresponding arenes (30 and 3p in 55% and 90% yields, respectively). Alkenyl bromide can be efficiently hydrodehalogenated as well (3q, 93%). Additionally, all the alkyl bromides tested here were readily debrominated to give 3r-w in excellent yields (>98%). Moreover, alkyl chlorides 2' were also viable substrates, giving excellent yields under the standard conditions (Scheme 3).

To further apply the **1a**/AIBN system to realize a cascade cyclization, a series of substrates bearing *ortho*-allyl moieties were employed. The results are given in Scheme 4 (see the ESI† for details). As seen, **4a–d** were converted to the corresponding 5-*exo*-trig cyclization products in nearly quantitative yields (88–99%) and high chemo-selectivity (cyclization: direct hydrodehalogenation >15:1). Lengthening the side alkyl chain had a trivial effect on the reaction (**4b**, 95%). The excellent reactivity and chemoselectivity indicated that the present system could avoid the over reduction observed in other similar processes. ^{15b} For the substrate bearing the NH group, both cyclization **5e** (50%) and direct hydrodehalogenation **5e**′ (20%) products were observed with a recovery of about 20% of the starting material. Formation of **5e**′ is presumably ascribed to the competitive HAT from the NH moiety or **1a** to the aryl radical.

Next, we focused on elucidating the mechanistic details of the present system with particular interest in the process of C-Br bond activation, i.e., to identify whether the phosphinyl radical directly abstracts the bromine atom from bromobenzene like the reactions mediated by tin hydrides, 17 silanes, 13a silylated cyclohexadienes,13b and N-heterocyclic carbene-borane complexes, 13c,18 or the phosphinyl radical transfers an electron to bromobenzene to trigger the cycle as depicted in Scheme 2b. DFT calculations showed that 1a-[P]' and 1b-[P]' should have comparable ability (with an energy difference of 1.3 kcal mol⁻¹, see the ESI† for details) in abstracting the bromine atom. This failed to explain the disparate yields of 90% for 1a-[P]' and <10% for 1b-[P]' (largely from a background reaction, see the text). Besides, according to the redox potentials of 1a-[P] $(E_{ox} = -2.39 \text{ V } \nu s. \text{ Fc in MeCN})$ and bromobenzene $(E_{\rm red} = -2.8 \text{ V})$, ¹⁶ the electron transfer from **1a-[P]** to bromobenzene is a feasible reversible process, while that for 1b-[P] $(E_{\rm ox} = -1.94 \text{ V})$ is thermodynamically prohibited. These findings preferentially support an ET-initiated mechanism rather than a direct bromine abstraction.



Scheme 4 Reductive radical cyclization using 1.0 mmol of 4 in 3.0 mL toluene. Isolated yields were given.

Edge Article Chemical Science

Scheme 5 Control experiments with ¹H NMR yields (using 1,3,5-trimethoxybenzene as the internal standard; see the ESI† for details).

On the other hand, we have recently found that 1a reacted smoothly with AIBN to produce the bisphosphine [1a-P]2.10 Considering the facile interconversion between [1a-P]₂ and phosphinyl radical 1a-[P],19 bisphosphine was synthesized separately to examine whether it may have functioned as a radical reservoir during hydrodehalogenation. As depicted in Scheme 5 (eqn (1)), no product was detected when [1a-P]₂ was mixed with bromobenzene under the standard conditions. This is presumably because the relatively strong P-P bond of [1a-P]₂ made it difficult to dissociate to provide the corresponding phosphinyl radical 1a-[P] at the reaction temperature (90 °C, see the ESI† for details). Thus, the possibility of [1a-P]₂ as a reaction intermediate was excluded. Because the solvent toluene could be a common hydrogen donor, and if so, it has potential to quench the phenyl radical. To examine this, deuterium labelling experiments were conducted in toluene- d_8 (eqn (2) and (3)). The absence of deuterium incorporation indicated that the hydrogen could not come from toluene. Furthermore, replacement of 1a with its deuterated counterparts 1a-d resulted in the desired product with a 90% deuterium abundance (eqn (4)). Hence, 1a is suggested to be the hydrogen source at last.

Based on these control experiments, the plausible catalytic mechanism proposed in Scheme 2 can be verified. The *in situ* generated phosphinyl radical preferentially served as a potent electron donor to activate the bromides. The reaction is most likely to proceed through an ET-initiated radical chain process,²⁰ although a direct bromine abstraction cannot be completely excluded at the present stage.

Conclusions

In conclusion, in this work we unlocked unprecedented radical reactivity of diazaphosphinanes to achieve efficient hydrodehalogenation and cascade cyclization, which is distinguished from their well-established hydridic reactivity. The phosphinyl radical, accessed in a catalytic cycle, is believed to be responsible for activating the carbon–bromine bonds through ET. This new reaction may provide a superior approach to remove halogens from many organohalides in terms of substrate scope, reaction efficiency and chemo-selectivity. Exploitation of other radical reductions using the strategy presented here is ongoing.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful for the financial grants from the National Natural Science Foundation of China (No. 21973052, 21933008, 21602116, and 91745101), National Science & Technology Fundamental Resource Investigation Program of China (No. 2018FY201200), and Tsinghua University Initiative Scientific Research Program (No. 20181080083).

Notes and references

- (a) E.-U. Würthwein, G. Lang, L. H. Schappele and H. Mayr, J. Am. Chem. Soc., 2002, 124, 4084–4092; (b) D. W. Brinkley and J. P. Roth, J. Am. Chem. Soc., 2005, 127, 15720–15721; (c) M. Breugst, H. Zipse, J. P. Guthrie and H. Mayr, Angew. Chem., Int. Ed., 2010, 49, 5165–5169; (d) Y. Li and X.-Q. Zhu, ACS Omega, 2018, 3, 872–885.
- 2 (a) D. Y. Ong, C. Tejo, K. Xu, H. Hirao and S. Chiba, Angew. Chem., Int. Ed., 2017, 56, 1840–1844; (b) B. Rösch, T. X. Gentner, H. Elsen, C. A. Fischer, J. Langer, M. Wiesinger and S. Harder, Angew. Chem., Int. Ed., 2019, 58, 5396–5401.
- (a) S. Fukuzumi, Org. Biomol. Chem., 2003, 1, 609-620; (b)
 I. J. Rhile and J. M. Mayer, J. Am. Chem. Soc., 2004, 126, 12718-12719; (c) J. P. Roth, S. Lovell and J. M. Mayer, J. Am. Chem. Soc., 2000, 122, 5486-5498.
- 4 (*a*) D. Gudat, A. Haghverdi and M. Nieger, *Angew. Chem., Int. Ed.*, 2000, **39**, 3084–3086; (*b*) S. Burck, D. Gudat, M. Nieger and W.-W. Du Mont, *J. Am. Chem. Soc.*, 2006, **128**, 3946–3955.
- (a) M. R. Adams, C. H. Tien, B. S. N. Huchenski, M. J. Ferguson and A. W. H. Speed, Angew. Chem., Int. Ed., 2017, 56, 6268–6271; (b) M. R. Adams, T. Chieh-Hung, M. Robert and A. W. H. Speed, Angew. Chem., Int. Ed., 2017, 56, 16660–16663; (c) C.-H. Tien, M. R. Adams, M. J. Ferguson, E. R. Johnson and A. W. H. Speed, Org. Lett., 2017, 19, 5565–5568; (d) T. Lundrigan, E. N. Welsh, T. Hynes, C.-H. Tien, M. R. Adams, K. R. Roy, K. N. Robertson and A. W. H. Speed, J. Am. Chem. Soc., 2019, 141, 14083–14088.
- 6 C. C. Chong, H. Hirao and R. Kinjo, *Angew. Chem., Int. Ed.*, 2015, **54**, 190–194.
- 7 (a) C. C. Chong, B. Rao and R. Kinjo, ACS Catal., 2017, 7, 5814–5819; (b) S. Miaskiewicz, J. H. Reed, P. A. Donets, C. C. Oliveira and N. Cramer, Angew. Chem., Int. Ed., 2018, 57, 4039–4042.

8 (a) B. Rao, C. C. Chong and R. Kinjo, J. Am. Chem. Soc., 2018, 140, 652-656; (b) T. Hynes, E. N. Welsh, R. McDonald, M. J. Ferguson and A. W. H. Speed, Organometallics, 2018, 37, 841-844.

Chemical Science

- 9 J. Zhang, J.-D. Yang and J.-P. Cheng, Angew. Chem., Int. Ed., 2019, 58, 5983-5987.
- 10 J. Zhang, J.-D. Yang and J.-P. Cheng, Chem. Sci., 2020, 11, 3672-3679.
- 11 (a) P. Knochel, W. Dohle, N. Gommermann, F. F. Kneisel, F. Kopp, T. Korn, I. Sapountzis and V. A. Vu, Angew. Chem., Int. Ed., 2003, 42, 4302-4320; (b) F. Alonso, I. P. Beletskaya and M. Yus, Chem. Rev., 2002, 102, 4009-4092.
- 12 (a) B. Sahoo, A.-E. Surkus, M.-M. Pohl, J. Radnik, M. Schneider, S. Bachmann, M. Scalone, K. Junge and M. Beller, Angew. Chem. Int. Ed., 2017, 56, 11242-11247; (b) V. V. Grushin and H. Alper, Chem. Rev., 1994, 94, 1047-1062.
- 13 (a) C. Chatgilialoglu, Acc. Chem. Res., 1992, 25, 188-194; (b) A. Studer and S. Amrein, Angew. Chem., Int. Ed., 2000, 39, 3080-3082; (c) X. Pan, E. Lacôte, J. Lalevée and D. P. Curran, J. Am. Chem. Soc., 2012, 134, 5669-5674; (d) A. Dewanji, C. Mück-Lichtenfeld and A. Studer, Angew. Chem., Int. Ed., 2016, 55, 6749-6752.
- 14 J.-P. Cheng, et al., Internet Bond-energy Databank (iBonD) Home Page, http://ibond.nankai.edu.cn, accessed at Feb. 2020.

- 15 (a) S. Rohrbach, R. S. Shah, T. Tuttle and J. A. Murphy, Angew. Chem., Int. Ed., 2019, 58, 11454-11458; (b) S. S. Hanson, E. Doni, K. T. Traboulsee, G. Coulthard, J. A. Murphy and C. A. Dyker, Angew. Chem., Int. Ed., 2015, 54, 11236-11239.
- 16 (a) C. Costentin, M. Robert and J.-M. Savéant, J. Am. Chem. Soc., 2004, 126, 16051-16057; (b) L. Pause, M. Robert and J.-M. Savéant, J. Am. Chem. Soc., 1999, 121, 7158-7159.
- 17 P. A. Baguley and J. C. Walton, Angew. Chem., Int. Ed., 1998, 37, 3072-3082.
- 18 D. P. Curran, A. Solovyev, M. M. Brahmi, L. Fensterbank, M. Malacria and E. Lacôte, Angew. Chem., Int. Ed., 2011, 50, 10294-10317.
- 19 (a) R. Edge, R. J. Less, E. J. L. McInnes, K. Müther, V. Naseri, J. M. Rawson and D. S. Wright, Chem. Commun., 2009, 1691-1693; (b) O. Puntigam, D. Förster, N. A. Giffin, S. Burck, J. Bender, F. Ehret, A. D. Hendsbee, M. Nieger, J. D. Masuda and D. Gudat, Eur. J. Inorg. Chem., 2013, 2041-2050; (c) D. Förster, H. Dilger, F. Ehret, M. Nieger and D. Gudat, Eur. J. Inorg. Chem., 2012, 3989-3994; (d) M. Blum, O. Puntigam, S. Plebst, F. Ehret, J. Bender, M. Nieger and D. Gudat, Dalton Trans., 2016, 45, 1987-1997; (e) D. M. C. Ould, A. C. Rigby, L. C. Wilkins, S. J. Adams, J. A. Platts, S. J. A. Pope, E. Richards and R. L. Melen, Organometallics, 2018, 37, 712-719.
- 20 A. Studer and D. P. Curran, Nat. Chem., 2014, 6, 765-773.