

Cite this: *Chem. Sci.*, 2020, **11**, 12364

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 3rd August 2020  
Accepted 21st October 2020


DOI: 10.1039/d0sc04244g  
[rsc.li/chemical-science](http://rsc.li/chemical-science)

## Introduction

A novel reducing system, consisting of the reagent-pair, triethylsilane and potassium *tert*-butoxide was reported by Stoltz, Grubbs *et al.* in 2013.<sup>1</sup> The combination of the two reagents has since been investigated by a number of research groups<sup>2–15</sup> and provides a range of distinctive reaction types, arising through an unprecedented menu of reactive intermediates formed in the reaction, including triethylsilyl radicals **1**, silanates **2** as hydrogen atom donors to both closed shell molecules and to radicals, and as potential hydride ion donors, and *tert*-butoxy-triethylsilyl radical anions **3** as a very powerful electron donor. Exposing substrates simultaneously to multiple reactive intermediates is not routinely encountered in organic chemistry, other than in modelling of prebiotic conditions,<sup>16</sup> and so the variety of reactive intermediates produced by this reagent pair provides opportunities to witness unusual outcomes.

Thus, triethylsilyl radicals **1** are candidates for the conversions of substrates **4–7** (ref. 2–6, 8 and 13) to their products **11–14**. (Note that silylation reactions, as in formation of **13** usually occur at lower temperatures, here 45 °C). On the other hand, Jeon has established<sup>9</sup> that silanate complex **2'** (and less efficiently **2**) conducts a potassium ion-dependent H-atom transfer to afford hydrosilylation products **15** from styrenes such as **8** at 80 °C. Tuttle, Murphy *et al.* have reported that *N*-benzylindoles **9** are deprotected by electron transfer reactions with **3** acting as

electron donor.<sup>7</sup> In each of the above cases, the products can be attributed to one of the reactive intermediates. Most recently, a more complex rearrangement of *N*-aryl indoles (e.g. **10**) to



Scheme 1


<sup>a</sup>Department of Pure and Applied Chemistry, Thomas Graham Building, 295 Cathedral Street, Glasgow, G1 1XL, UK. E-mail: john.murphy@strath.ac.uk

<sup>b</sup>GlaxoSmithKline Medicines Research Centre, Gunnels Wood Road, Stevenage, Hertfordshire, SG1 2NY, UK


† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0sc04244g

‡ These authors contributed equally.





Scheme 2



Scheme 3

increased to 72% by lowering the temperature to  $70^\circ\text{C}$  (Scheme 3 and ESI†). The detection of imine 30 during the optimisation studies suggests that an iminyl anion 21 ( $\text{R} = {^n\text{Pr}}$ ) was a key intermediate in the reactions. These reactions are therefore proposed to occur by hydride ion delivery to the nitrile by intermediate 2. The mechanism of conversion of imines *e.g.* 30 to amines (in this case, 24) comes up for discussion later in this paper.

The effect of the identity of the base and silane present was then investigated. On changing the counter-ion on butoxide from potassium to sodium or lithium, no reaction was observed (see ESI†). Other potassium bases such as KHMDS, KOH, and KOEt were also unsuccessful. KH was somewhat successful with 24 being isolated in 12% yield, whereas NaH gave no reaction. These results underline the special reactivity of potassium *tert*-butoxide in this reagent system, which cannot be replicated by sodium *tert*-butoxide or lithium *tert*-butoxide. The effect of solvent on the reaction was also investigated, and solvent-free conditions were found to be optimal for the cyclisation (see ESI†).

The optimised conditions were then used to further study the scope of the reaction with substrates related to 23. Firstly, the ethoxy derivative 31 was also successful, with cyclised product 24 ( $\text{R} = \text{Et}$ ) isolated in 65% yield. Halide leaving groups were then tested (Table 1, entries 3–6). Interestingly, methoxide out-performed halide leaving groups for the formation of 24, with the halides following the general trend of  $\text{S}_{\text{N}}\text{Ar}$  reactivity ( $\text{F} > \text{Cl} > \text{Br} = \text{I}$ ). Bromo- and iodo-substituted substrates 34 and 35

dihydroacridines (in this case, 17) features sequential electron transfer from 3 and H-atom transfer from 2.<sup>14</sup> In addition to these transformations, the reagent pair  $\text{Et}_3\text{SiH}/\text{KO}^t\text{Bu}$  has found wider applications in silylation of alcohols<sup>10</sup> and amines,<sup>11</sup> as well as the silylation of terminal alkynes.<sup>15</sup> The broad range of possible pathways featuring different reactive intermediates is what makes this reagent-pair so fascinating (Scheme 1).<sup>12</sup>


During a recent study, we showed that  $\text{Et}_3\text{SiH}/\text{KO}^t\text{Bu}$  carries out reductive decyanation of benzylic nitriles (*e.g.* 18  $\rightarrow$  19, Scheme 2)<sup>7,17</sup> and our starting point for this current study was to find out more about the reactivity of the substrates and intermediates.

In 2017, Chiba *et al.* uncovered<sup>18</sup> a probe for hydride-based reduction of nitriles, where substrates *e.g.* 20 reacted with a composite of NaH and LiI to form an iminyl anion 21 that displaced the *o*-MeO group in a concerted  $\text{cS}_{\text{N}}\text{Ar}$  reaction<sup>19</sup> to form indolene 22 [ $\text{R} = (\text{CH}_2)_4$ ]. In his elegant paper, aminyl anions also underwent efficient cyclisation. We wondered whether nitriles that are subjected to the  $\text{Et}_3\text{SiH} + \text{KO}^t\text{Bu}$  reagent would behave similarly, giving evidence for formation of iminyl anion intermediates through hydride ion delivery from 2.<sup>20</sup>

## Results and discussion

Substrates 23, 25 and 27 (Scheme 2) were prepared (see ESI†) and reacted with the  $\text{Et}_3\text{SiH}/\text{KO}^t\text{Bu}$  mixture. In each case, cyclisation with displacement of the methoxy group was observed. Our initial conversion of 23  $\rightarrow$  24 occurred in 32% yield (Scheme 2), but upon optimisation, the yield of 24 was

Table 1 Testing different leaving groups



| Entry | Substrate | 24/% | 29/% | 30/%  | 37/% |
|-------|-----------|------|------|-------|------|
| 1     | 23        | 72   | 11   | —     | —    |
| 2     | 31        | 65   | 8    | —     | —    |
| 3     | 32        | 44   | —    | Trace | —    |
| 4     | 33        | 19   | —    | —     | —    |
| 5     | 34        | —    | —    | —     | 57   |
| 6     | 35        | —    | —    | —     | 78   |

did not afford any cyclised products and instead, dehalogenated compound **37** was isolated, suggesting that dehalogenation of iodides and bromides was more rapid than activation of the nitrile. Although many mechanisms can be considered, dehalogenation is a hallmark of reactions of silyl radicals **1** or can result from electron transfer chemistry of **3**.

Changing the methoxy group to benzyloxy in **36** brought about different chemistry. No displacement of the benzyloxy group was detected and, instead, compounds **38** (5%) and **39** (82%) were isolated (Scheme 4). Both products suggested an initial activation at the benzyl position, most likely *via* anion **40**. Cyclisation onto the nitrile would afford **41** which would be converted to **42** through a proton shift. The electron-rich alkene in **42** will readily undergo electron transfer and coupling to molecular oxygen to afford **43**.<sup>21–23</sup> If this can convert into a hydroperoxide, then reductive cleavage of the O–O bond can occur during the reaction. Otherwise, **43** could protonate on workup to a hydroperoxyketal, which can lose hydroperoxide anion in a hydrolysis that then leads to **39**. Alternatively, any residual anion **40** would also react with air on work up, ultimately leading to ester **38**.

Our next steps were to establish the necessary components for the cyclisation of **23** to **24**. Control reactions were now performed (Table 2). The parent reaction is shown as entry 1.

Recently, we proposed<sup>14</sup> that the reactivity of the  $\text{Et}_3\text{SiH} + \text{KO}^\ddagger\text{Bu}$  couple could be reproduced in the absence of the silane, provided that an alternative source of silyl radicals was present. To this end, entry 2 shows that when the silane was replaced by the disilane **44** (Scheme 5) in the presence of the electron donor **47**,<sup>24</sup> the radical anion of di-*p*-*tert*-butylbiphenyl, the cyclisation reaction was still observed, affording **24** (19%).


Entries 3 and 4 show that in the absence of a silyl source or an electron donor source, the reaction is not observed, while entry 5 shows that  $\text{KO}^\ddagger\text{Bu}$  alone cannot bring about the reaction.

Our explanation for entry 2 is that the electron donor **47** can cleave the disilane **44** to a silyl radical **45** and a silyl anion **46** as shown in Scheme 5.<sup>25</sup> The silyl radical can react with many species in solution. Notably, it can add to arene rings in the

Table 2 Cyclisation reaction – mechanistic studies

|       |  |       |
|-------|--|-------|
|       |  |       |
| Entry | Conditions <sup>a</sup>  | 24/%  |
| 1     | $\text{Et}_3\text{SiH} + \text{KO}^\ddagger\text{Bu}$                  | 72    |
| 2     | <b>47</b> + $\text{Me}_3\text{SiSiMe}_3 + \text{KO}^\ddagger\text{Bu}$ | 19    |
| 3     | <b>47</b> + $\text{KO}^\ddagger\text{Bu}$                              | Trace |
| 4     | $\text{Me}_3\text{SiSiMe}_3 + \text{KO}^\ddagger\text{Bu}$             | —     |
| 5     | $\text{KO}^\ddagger\text{Bu}$  | —     |


<sup>a</sup> 3 equiv. of all reagents were used.



Scheme 5



substrate to generate intermediates **49** that feature a labile H atom.<sup>2–6,13</sup> This can react with a silyl radical **45** to form trimethylsilane or with silyl anion **46** to form trimethylsilane as shown in Scheme 5. This would mean that the missing trialkylsilane reagent ( $\text{Me}_3\text{SiH}$  in this experiment) would be generated *in situ*, starting from the disilane.

A series of substrates, **51**, **20** and **55** (Scheme 6), was now prepared and tested under the optimised conditions, with surprising results. From substrate **51**, reductive decyanation to **52** was observed in 99% yield, with only a trace amount of cyclised product **53** detected. However, from the analogous substrate **20**, cyclisation to **54** was observed in 65% yield. Pyridine-containing substrate **55** afforded product **56** (24%), along with dimer **57** (15%). This compound **57** might arise by dimerisation of radical anion **58**, *e.g.* if electron transfer occurred from radical anion **3**, followed by double  $\text{cS}_\text{N}\text{Ar}$  cyclisation. Alternatively, and more probably, cyclic imine **60** could be deprotonated under the basic conditions to anion **61**,<sup>26</sup> which could then attack another molecule of **55** to give anion **62**, which affords bis-indolenine **57** by  $\text{cS}_\text{N}\text{Ar}$  cyclisation. Observation of this dimerisation solely for this substrate could




Scheme 4





Scheme 6




Scheme 8

then be attributed to enhancement of the acidity of the iminyl proton in **60** by the pyridine ring.

The results to date are consistent with bicyclic imines such as **60** as key intermediates in the formation of the final indolines, and so we were curious to probe the behaviour of related imines in the presence of the reductive silane–butoxide reagent pair.

To access imines related to **30** (Scheme 3), we considered that an iminyl anion could form by addition of a Grignard reagent to a nitrile and then undergo cyclisation, in the manner of Gademann *et al.*<sup>27</sup> To test this, substrate **23** was treated with MeMgBr at 70 °C, however no reaction occurred. Upon warming to 130 °C, however, products **63**–**65** were isolated with optimum yields arising from 4 equiv. of Grignard reagent (Scheme 7 and ESI†). Compounds **63** and **64** are indicative of the proposed mechanism for indolene formation. Compound **65** could arise by deprotonation of the iminyl-CH<sub>3</sub> group of **64** by MeMgBr, before attack onto the nitrile group of another molecule of **23**. The resulting imine anion can then undergo cSNAr to yield **65**.

The complications in Scheme 7 leading to a low yield of **64** arose from the ease of deprotonation of the methyl group in **64**. To prevent such complications, a Grignard reagent was used that cannot be deprotonated in the  $\alpha$ -position, *i.e.* PhMgBr. Interestingly, cyclisation to an inseparable mixture of indoles **66**




Scheme 7

and **67** (in 30% yield each, calculated by NMR internal standard) was observed (Scheme 8). This transformation shows loss of a propyl substituent, and aromatisation of the ring system to give indole products undoubtedly provides the driving force for this.

Grignard reagents have been previously reported in the literature to facilitate SET reactions to reducible substrates.<sup>28</sup> Therefore, we propose that electron transfer from PhMgBr must occur to the conjugated indolene **69**. First, PhMgBr adds to the nitrile of **23**, forming **68**, which can undergo cSNAr to form **69**. Electron transfer to the conjugated indolene **69** then occurs to form **70**, which can aromatise with loss of an alkyl group to form **71**, which ultimately protonates to **66** upon work-up. The formation of compound **67** is rationalised by the presence of phenyl radicals, generated upon single electron oxidation of PhMgBr, initiating a BHAS mechanism as previously reported in the literature.<sup>29</sup>

We then investigated if the reducing mixture resulting from the combination of Et<sub>3</sub>SiH and KO<sup>t</sup>Bu, could perform the same transformation of indolene to indoline and thereby give evidence of electron transfer from intermediate **3**. Compound


Table 3 Aromatisation reaction – mechanistic studies



| Entry | Conditions <sup>a</sup>                                       | 72/% | 73/% | 74/%  | 75/%  |
|-------|---|------|------|-------|-------|
| 1     | Et <sub>3</sub> SiH + KO <sup>t</sup> Bu                      | —    | 45   | 24    | —     |
| 2     | Et <sub>3</sub> SiH + KO <sup>t</sup> Bu + TEMPO <sup>b</sup> | 9    | 84   | Trace | Trace |
| 3     | K + KO <sup>t</sup> Bu  | 28   | 36   | —     | Trace |


<sup>a</sup> Entry 1 and 2: Et<sub>3</sub>SiH (3 equiv.), KO<sup>t</sup>Bu (3 equiv.), TEMPO (1 equiv.); entry 3: KO<sup>t</sup>Bu (1 equiv.), K (1.3 equiv.). <sup>b</sup> TEMPO-SiEt<sub>3</sub> was detected by GCMS (see ESI).





Scheme 9

72 was treated with  $\text{Et}_3\text{SiH}$  and  $\text{KO}^t\text{Bu}$ , and compounds 73 and 74 were isolated in 45% and 24% respectively (Table 3, entry 1). The elimination of a methyl group from 72 clearly mirrors the electron transfer reactions seen with  $\text{PhMgBr}$ . Moreover, the formation of silylated derivative 74 (an inseparable mixture of 2 regioisomers was isolated) results from the presence of triethylsilyl radicals, analogous to the phenyl radicals above. The reaction was repeated in the presence of TEMPO. The outcome was to improve the yield of indole 73 to 84% (entry 2). This outcome likely arises, at least in part, from trapping of triethylsilyl radicals by TEMPO, thereby inhibiting the formation of 74. Two further pieces of evidence support the electron transfer proposal: (i) exposure of substrate 72 to potassium metal and  $\text{KO}^t\text{Bu}$  also afforded 73 (36%), 72 (28%) and 75 (trace amounts) (entry 3); (ii) analogue 76 (Scheme 9) underwent reaction with  $\text{Et}_3\text{SiH} + \text{KO}^t\text{Bu}$  to afford principally indoline 77 (86%) together with indole 78 (trace amounts). Product 78 arises from an




Scheme 10

analogous cleavage in radical anion 79 to that seen for radical anion 70. The pentyl side-chain of 78 shows the fate of the cleaved radical, which simply abstracts a hydrogen atom from silane or hydrogen atom donor 2. The difference in outcome for substrates 72 and 76 relates to the fragmentation of their radical anions – when the radical anion of 72 fragments, a methyl group is lost and diffuses away from the substrate. In contrast, fragmentation of radical anion 79 sees the fragmented radical tethered to the indole structure in 80. Radical re-addition to the indole anion reforms radical anion 79, which then abstracts an H-atom (e.g. from triethylsilane or from species 2) to give indoline 77.

The scope of the groups that can be expelled upon aromatisation was also investigated (Scheme 10). These results show that phenyl, allyl and benzyl are feasible leaving groups.

Firstly taking substrate 81, this mirrors the reactions of imines 69, 72 and 76. Loss of a phenyl radical is more difficult than loss of an alkyl radical, but indole 84 is still formed in 35% yield. Also detected were products 85 and 87, resulting from attack on 84 by phenyl or triethylsilyl radicals and subsequent rearomatisation. Again, this mimics the addition of phenyl radicals and triethylsilyl radicals seen respectively in 67 and 74. In addition, compound 86 was detected in an inseparable mixture with compound 85 with  $^1\text{H}$  NMR data and GC-MS data consistent with those previously reported in the literature. The



Scheme 11



mechanism envisaged for the formation of **86** is somewhat analogous to that for compound **92** (see below).

For substrate **82**, the products **88** (27%) and **89** (22%) can be explained by invoking  $\text{KO}^\bullet\text{Bu}$ -induced isomerisation of a terminal allyl group to internal alkene **96** (Scheme 11). We have recently reported that allyl groups undergo base-induced isomerisation under the  $\text{Et}_3\text{SiH}/\text{KO}^\bullet\text{Bu}$  conditions.<sup>14</sup> Subjecting this compound, **96**, to electron transfer from donor **3** gives radical anion **94**. Expulsion of an allyl radical accounts for the formation of indole **88**.

The second product formed from substrate **82** is the quinoline **89** (22%). This is a really interesting product. Focusing on the 6 carbons of the allyl substituents in **82**, it appears that one carbon has been incorporated into the ring system during a ring-expansion, two carbons have been lost during the rearrangement, and the remaining three carbons end up as the methyl and ethyl substituents on neighbouring ring carbons in **89** – a rearrangement of serious complexity. Our working hypothesis is that the product **89** arises also from intermediate diene **96**. Jeon recently demonstrated H-atom addition to styrenes by reactive intermediate **2'** (inset, Scheme 11) formed from diethylsilane. A  $\text{K}^+$  ion, complexed by the aromatic ring in the styrene, held the silanate anion in **2'** (and analogues) close to the aromatic ring, and this complexation was essential for H-atom addition. Here, the feasibility of H-atom addition from **2**, rather than **2'**, to a side-chain alkene should also increase when the alkene is nearer to the aromatic ring, thereby directing H-atom addition to **96** to form radical **97**. An aza-version of a cyclopropylcarbinyl rearrangement governs the ring-expansion to radical **99**. This radical has an adjacent H-atom that is easily acidic enough to be removed by  $\text{KO}^\bullet\text{Bu}$ , affording the quinoline radical anion **100**. This undergoes reversible

cyclisation to cyclopropylcarbinyl radical **101**, which must very occasionally fragment to distal radical anion **102**; expulsion of a vinyl radical (or a vinyl anion) then affords benzylic anion **103** (or its benzylic radical counterpart) which affords **89** on workup.<sup>30</sup>

Having proposed a route to the quinoline **89** from substrate **82**, we note that two further quinolines, **92** and **93**, which arise from substrate **83**, require explanation. We have recently shown that benzylic C–H bonds can undergo abstraction of an H-atom under the conditions of these reactions, by triethylsilyl radicals **1**.<sup>13</sup> In this case, this would lead to radical **105** (Scheme 12). Cyclopropylcarbinyl radical rearrangement would lead to ring-expansion to radical **107**, which, following deprotonation, would expel a benzyl radical to yield quinoline anion **109**. Protonation from  $^\bullet\text{BuOH}$ , followed by electron transfer from **3** would give radical anion **110**. Expulsion of a phenyl radical affords anion **111** that abstracts a proton (from  $^\bullet\text{BuOH}$  or on workup) to give **93**.

We again propose **110'** as the source of the other product, **92**. Cyclisation to the neighbouring phenyl ring gives radical anion **112**. The drive to aromaticity can then oversee the expulsion of an H atom (or a proton followed by an electron) and a hydride ion to give product **92**.

2-Phenyl-substituted indolenines **81**–**83** are likely to be more receptive towards electron transfer than analogues with H or alkyl groups substituted in the 2-position, but these substrates illustrate well here the array of reactive intermediates in the  $\text{KO}^\bullet\text{Bu}/\text{Et}_3\text{SiH}$  reagent pair.

## Conclusions

The reagent pair  $\text{KO}^\bullet\text{Bu} + \text{Et}_3\text{SiH}$  provides a unique interplay of reactive intermediates to react with substrates. This study of benzyl nitriles and indolenines features products arising from (i) hydride addition from silanate complex **2**, (ii) electron transfer from **3** (iii) hydrogen atom transfer from anion **2**, and (iv) hydrogen atom abstraction by silyl radicals **1**. The range of product types observed illustrates a unique diversity of outcomes.

## Conflicts of interest


There are no conflicts to declare.

## Acknowledgements

We thank GSK and EPSRC for i-CASE awards to AJS and DD. We thank (i) the National Mass Spectrometry Service, Swansea and (ii) the Cronin group and Dr Emma Carrick (University of Glasgow) for high resolution mass spectra.

## References

- 1 A. Fedorov, A. A. Toutov, N. A. Swisher and R. H. Grubbs, *Chem. Sci.*, 2013, **4**, 1640–1645.
- 2 A. A. Toutov, W.-B. Liu, K. N. Betz, A. Fedorov, B. M. Stoltz and R. H. Grubbs, *Nature*, 2015, **518**, 80–84.



Scheme 12



3 A. A. Toutov, W.-B. Liu, K. N. Betz, B. M. Stoltz and R. H. Grubbs, *Nat. Protoc.*, 2016, **10**, 1897–1903.

4 W.-B. Liu, D. P. Schuman, Y.-F. Yang, A. A. Toutov, Y. Liang, H. F. T. Klare, N. Nesnas, M. Oestreich, D. G. Blackmond, S. C. Virgil, S. Banerjee, R. N. Zare, R. H. Grubbs, K. N. Houk and B. M. Stoltz, *J. Am. Chem. Soc.*, 2017, **139**, 6867–6879.

5 S. Banerjee, Y.-F. Yang, I. D. Jenkins, Y. Liang, A. A. Toutov, W.-B. Liu, D. P. Schuman, R. H. Grubbs, B. M. Stoltz, E. H. Krenske, K. N. Houk and R. N. Zare, *J. Am. Chem. Soc.*, 2017, **139**, 6880–6887.

6 A. A. Toutov, M. Salata, A. Fedorov, Y.-F. Yang, Y. Liang, R. Cariou, K. N. Betz, E. P. A. Couzijn, J. W. Shabaker, K. N. Houk and R. H. Grubbs, *Nat. Energy*, 2017, 17008, DOI: 10.1038/nenergy.2017.

7 A. J. Smith, A. Young, S. Rohrbach, E. F. O'Connor, M. Allison, H.-S. Wang, D. L. Poole, T. Tuttle and J. A. Murphy, *Angew. Chem., Int. Ed.*, 2017, **56**, 13747–13751.

8 (a) A. A. Toutov, K. N. Betz, A. Fedorov, B. M. Stoltz, W.-B. Liu and R. H. Grubbs, US2019/0048030A1, 2019; (b) A. A. Toutov, K. N. Betz, A. Fedorov, B. M. Stoltz, W.-B. Liu and R. H. Grubbs, US2020/0115396A1, 2020.

9 P. Asgari, Y. Hua, C. Thiamsiri, W. Prasitwatcharakorn, A. Karedath, X. Chen, S. Sardar, K. Yum, G. Leem, B. S. Pierce, K. Nam, J. Gao and J. Jeon, *Nat. Catal.*, 2019, **2**, 164–173.

10 A. A. Toutov, K. N. Betz, A. M. Romine and R. H. Grubbs, US2019/0241587A1, 2019.

11 (a) A. A. Toutov, K. N. Betz, A. M. Romine and R. H. Grubbs, US 2019/0218232A1, 2019; (b) F. Palumbo, S. Rohrbach, T. Tuttle and J. A. Murphy, *Helv. Chim. Acta*, 2019, e1900235.

12 I. D. Jenkins and E. H. Krenske, *ACS Omega*, 2020, **5**, 7053–7058.

13 J. N. Arokianathar, K. Kolodziejczak, F. Bugden, K. Clark, T. Tuttle and J. A. Murphy, *Adv. Synth. Catal.*, 2020, **362**, 2260–2267.

14 A. J. Smith, D. Dimitrova, J. N. Arokianathar, K. Kolodziejczak, A. Young, M. Alison, D. L. Pole, S. G. Leach, J. A. Parkinson, T. Tuttle and J. A. Murphy, *Chem. Sci.*, 2020, **11**, 3719–3726.

15 A. A. Toutov, K. N. Betz, D. P. Schuman, W. B. Liu, A. Fedorov, B. M. Stoltz and R. H. Grubbs, *J. Am. Chem. Soc.*, 2017, **139**, 1668–1674.

16 S. L. Miller, *Science*, 1953, **117**, 528–529.

17 For some previous papers on decyanation, see (a) J. M. R. Mattalia, *Beilstein J. Org. Chem.*, 2017, **13**, 267–284; (b) P. C. Too, G. H. Chan, Y. L. Tnay, H. Hirao and S. Chiba, *Angew. Chem., Int. Ed.*, 2016, **55**, 3719–3723; (c) E. Doni and J. A. Murphy, *Org. Chem. Front.*, 2014, **1**, 1072–1076.

18 A. Kaga, H. Hayashi, H. Hakamata, M. Oi, M. Uchiyama, R. Takita and S. Chiba, *Angew. Chem., Int. Ed.*, 2017, **56**, 11807–11811.

19 (a) E. E. Kwan, Y. Zeng, H. A. Besser and E. N. Jacobsen, *Nat. Chem.*, 2018, **10**, 917–923; (b) A. J. Lennox, *Angew. Chem., Int. Ed.*, 2018, **57**, 14898–14900; (c) S. Rohrbach, A. J. Smith, J. H. Pang, D. L. Poole, T. Tuttle, S. Chiba and J. A. Murphy, *Angew. Chem., Int. Ed.*, 2019, **58**, 16368–16388.

20 We verified that such benzylic nitrile substrates can be cyclised also with LiAlH<sub>4</sub> at 130 °C, although not at 70 °C, again indicative of cyclisation of an iminyl anion – See ESI†

21 J. A. Murphy, *J. Org. Chem.*, 2014, **79**, 3731–3746.

22 S. S. Hanson, E. Doni, K. T. Traboulsi, G. Coulthard, J. A. Murphy and C. A. Dyker, *Angew. Chem., Int. Ed.*, 2015, **54**, 11236–11239.

23 R. A. Aitken, A. D. Harper and A. M. Z. Slawin, *Synlett*, 2017, **28**, 1738–1742.

24 A. Guijarro and M. Yus, *Tetrahedron Lett.*, 1993, **34**, 3487–3490.

25 C. Däschlein and C. Strohmann, *Dalton Trans.*, 2010, **39**, 2062–2069.

26 N-Methylindoles have also been proposed to be deprotonated in the 2-position by Et<sub>3</sub>SiH + KO<sup>t</sup>Bu<sup>5</sup>

27 F. Huber, J. Roesslein and K. Gademann, *Org. Lett.*, 2019, **21**, 2560–2564.

28 (a) T. Holm, H. Baumann, H. Lönn, J. Lönngrén, H. Nyman and H. Ottosson, *Acta Chem. Scand., Ser. B*, 1983, **37**, 567–584; (b) T. Matsuyama, H. Yamataka and T. Hanafusa, *Chem. Lett.*, 1988, 1367–1370; (c) J. McKinley, A. Aponick, J. C. Raber, C. Fritz, D. Montgomery and C. T. Wigal, *J. Org. Chem.*, 1997, **62**, 4874–4876; (d) T. Ramnial, S. A. Taylor, J. A. C. Clyburne and C. J. Walsby, *Chem. Commun.*, 2007, 2066–2068; (e) N. Uchiyama, E. Shirakawa and T. Hayashi, *Chem. Commun.*, 2013, **49**, 364–366; (f) E. C. Ashby and T. L. Wiesemann, *J. Am. Chem. Soc.*, 1974, **96**, 7117–7119; (g) E. C. Ashby, *Acc. Chem. Res.*, 1988, **21**, 414–421.

29 A. Studer and D. P. Curran, *Angew. Chem., Int. Ed.*, 2011, **50**, 5018–5022.

30 An alternative direct fragmentation from **100** → **103** is also considered. Vinyl radical fragmentations often have high energy transition states, but additional driving force is expected in converting **102** → **103** as the nitrogen-containing ring in **102** is not aromatic but becomes aromatic in **103**.

