CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#) Cite this: *J. Mater. Chem. C*, 2020, 8, 1475

DOI: 10.1039/c9tc90268f

rsc.li/materials-c**Correction: Towards predicting the power conversion efficiencies of organic solar cells from donor and acceptor molecule structures**Yecheng Zhou,^{*a} Guankui Long,^b Ailin Li,^c Angus Gray-Weale,^a Yongsheng Chen^b and Tianying Yan^cCorrection for 'Towards predicting the power conversion efficiencies of organic solar cells from donor and acceptor molecule structures' by Yecheng Zhou *et al.*, *J. Mater. Chem. C*, 2018, **6**, 3276–3287.

There are errors in the equations on page 3279 of the article, under the heading '2.4 DOS determination.' The correct equations and related text are below.

DOSs are essential for the numerical simulations of CT in solar cells. However, we haven't found a theoretical method to estimate them for OSCs without any empirical or experimental parameters. Here, we put forward a method to estimate the DOS in OSCs. We assumed that each orbital can be occupied by two electrons, then $\text{DOS}(E)$ can be calculated by $\text{DOS}(E) = \sum_{\text{LUMO}(E)} 2$,

which is the summation over all LUMOs whose energies are E . The $\text{DOS}(E)$ should be Gaussian-like due to the Gaussian distribution of LUMOs. Assuming the charge occupation in the semiconductor obeys the Fermi–Dirac distribution, then, the charge density of the OSCs is given by:

$$\begin{aligned} n &= \int_{E_f}^{\infty} \text{DOS}(E) \frac{1}{1 + \exp\left(\frac{E - E_f}{k_B T}\right)} dE \\ &= \sum_{\text{LUMO}(E)} 2 \times \frac{1}{1 + \exp\left(\frac{E - E_f}{k_B T}\right)} \\ &\simeq \sum_{\text{LUMO}(E)} 2 \times \exp\left(\frac{E - E_f}{-k_B T}\right) \\ &= \sum_{\text{LUMO}(E)} 2 \times \exp\left(\frac{E - E_c}{-k_B T}\right) \times \exp\left(\frac{E_c - E_f}{-k_B T}\right) \\ &= N_c \times \exp\left(\frac{E_c - E_f}{-k_B T}\right) \end{aligned} \quad (8)$$

where E_c and E_f are the conduction band minimum (the LUMO energy level of PC₇₁BM) and the Fermi energy level, respectively. Since only LUMOs can be filled, $E - E_f$ is in the order of 0.1 eV which is several times higher than $k_B T$. The exponential term in Fermi–Dirac distribution

^a School of Chemistry, The University of Melbourne, Parkville, VIC 3010, Australia. E-mail: zhouych87@gmail.com^b Key Laboratory of Functional Polymer Materials, College of Chemistry, Nankai University, Tianjin 300071, China^c College of Materials Science and Engineering, National Institute of Advanced Materials, Nankai University, Tianjin 300071, China

is much larger than 1, then $\frac{1}{1 + \exp\left(\frac{E - E_f}{kT}\right)} \simeq \exp\left(\frac{E - E_f}{-kT}\right)$. Therefore, the DOS of the conduction band can be expressed as:

$$N_c = \sum_{\text{LUMO}(E)} 2 \times \exp\left(\frac{E - E_c}{-k_B T}\right). \quad (9)$$

The changes in the equations do not affect the interpretation of the results or the conclusions. The authors apologise for this oversight and for any confusion that it may have caused.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.